

Homework 3

Math 2031 — Professor Hamilton

Name: Imran Qasimi

Date: September 16, 2026

Section 5.2

Question 1C

Statement: $1^3 + 2^3 + 3^3 + \dots + n^3 = \frac{n^2(n+1)^2}{4}, n \in \mathbb{N}$

Proof by Induction:

Base Case: $k = 1, k \in \mathbb{N}$

LHS: $1^3 = 1$

RHS: $\frac{1^2(1+1)^2}{4} = \frac{4}{4} = 1$

Since LHS = RHS, the base case is true.

Induction Step:

Assume $1^3 + 2^3 + 3^3 + \dots + k^3 = \frac{k^2(k+1)^2}{4}$ is true for k .

We want to show the statement is true also for $(k+1)$ by showing the LHS=RHS.

Substituting $(k+1)$ into the equation:

$$1^3 + 2^3 + 3^3 + \dots + k^3 + (k+1)^3 = \frac{(k+1)^2(k+2)^2}{4}$$

Here we notice our IH:

$$\underbrace{1^3 + 2^3 + 3^3 + \dots + k^3}_{\text{Inductive Hypothesis}} + (k+1)^3 = \frac{(k+1)^2(k+2)^2}{4}$$

Replace:

$$\frac{k^2(k+1)^2}{4} + (k+1)^3 = \frac{(k+1)^2(k+2)^2}{4}$$

Factor $(k+1)^2$ from the LHS:

$$(k+1)^2 \left(\frac{k^2}{4} + k + 1 \right) = \frac{(k+1)^2(k+2)^2}{4}$$

Divide both sides by $(k+1)^2$, which is greater than 0.

$$\frac{k^2}{4} + k + 1 = \frac{(k+2)^2}{4}$$

Expanding the LHS we get:

$$\frac{k^2}{4} + k + 1 = \frac{k^2}{4} + k + 1$$

$$\therefore \forall n \in \mathbb{N}, 1^3 + 2^3 + 3^3 + \dots + n^3 = \frac{n^2(n+1)^2}{4}$$

■

Question 2A

Statement: $1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{(2n-1)(2n)(2n+1)}{6}, n \in \mathbb{N}$

Proof by Induction:

Base Case: $k = 1, k \in \mathbb{N}$

$$\text{LHS: } (2(1) - 1)^2 = 1^2 = 1$$

$$\text{RHS: } \frac{(2(1)-1)(2(1))(2(1)+1)}{6} = \frac{1 \cdot 2 \cdot 3}{6} = \frac{6}{6} = 1$$

Since LHS = RHS, the base case is true.

Induction Step:

Assume $1^2 + 3^2 + 5^2 + \dots + (2k-1)^2 = \frac{(2k-1)(2k)(2k+1)}{6}$ is true for k .

We want to show it is also true for $k+1$ by showing that the LHS=RHS.

Substituting $k+1$ into our equation:

$$1^2 + 3^2 + 5^2 + \dots + (2k-1)^2 + (2k+1)^2 = \frac{(2k+1)(2k+2)(2k+3)}{6} \text{ is true for } k.$$

Note our IH:

$$\underbrace{1^2 + 3^2 + 5^2 + \dots + (2k-1)^2}_{\text{Inductive Hypothesis}} + (2k+1)^2 = \frac{(2k+1)(2k+2)(2k+3)}{6}.$$

Substitute:

$$\frac{(2k-1)(2k)(2k+1)}{6} + (2k+1)^2 = \frac{(2k+1)(2k+2)(2k+3)}{6}$$

Factoring a $(2k+1)$ on both sides:

$$(2k+1) \cdot \left(\frac{(2k-1)(2k)}{6} + (2k+1) \right) = (2k+1) \cdot \frac{(2k+2)(2k+3)}{6}$$

Canceling the $(2k+1)$, it is greater than 0.

$$\frac{(2k-1)(2k)}{6} + (2k+1) = \frac{(2k+2)(2k+3)}{6}$$

Expanding both sides:

$$\frac{4k^2+10k+6}{6} = \frac{4k^2+10k+6}{6}$$

—

$$\therefore \forall n \in \mathbb{N}, 1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{(2n-1)(2n)(2n+1)}{6}$$

■

Question 3

Statement: For any real number a except 1, $1 + a + a^2 + a^3 + \dots + a^n = \frac{(a^{n+1}-1)}{a-1}$

Proof by Induction:

Base Case: $k = 1, k \in \mathbb{N}$

$$\text{LHS: } 1 + a^1 = a + 1$$

$$\text{RHS: } \frac{a^{1+1}-1}{a-1} = \frac{a^2-1}{1} = \frac{(a+1)(a-1)}{(a-1)} = a + 1$$

Since the LHS = RHS, the base case is true.

Induction Step:

Assume $1 + a + a^2 + a^3 + \dots + a^k = \frac{(a^{k+1}-1)}{a-1}$ is true for k .

We want to show that it is also true for $k+1$ by showing that the LHS=RHS.

Substitute $(k+1)$ into equation:

$$\underbrace{1 + a + a^2 + a^3 + \dots + a^k}_{\text{Induction Hypothesis.}} + a^{k+1} = \frac{(a^{k+2}-1)}{a-1}.$$

Substitute the IH:

$$\frac{(a^{k+1}-1)}{a-1} + a^{k+1} = \frac{(a^{k+2}-1)}{a-1}.$$

Combining the terms on our LHS:

$$\frac{a^{k+1}-1+a^{k+2}-a^{k+1}}{a-1} = \frac{(a^{k+2}-1)}{a-1}.$$

Canceling like terms and rearranging terms:

$$\frac{(a^{k+2}-1)}{a-1} = \frac{(a^{k+2}-1)}{a-1}$$

$$\therefore \forall n \in \mathbb{N} \text{ and } \forall a \in \mathbb{R} : \left(1 + a + a^2 + a^3 + \dots + a^n = \frac{(a^{n+1}-1)}{a-1} \right)$$

■

Question 8

Statement: “The Fibonnaci numbers $f_n, n = 1, 2, 3, \dots$ are defined recursively by the formulas $f_1 = 1, f_2 = 1, f_n = f_{n-1} + f_{n-2}$ for all $n \geq 3$.”

(A) $f_1 = 1, f_2 = 1, f_3 = 2, f_4 = 3, f_5 = 5, f_6 = 8, f_7 = 13, f_8 = 21, f_9 = 34, f_{10} = 55$

(B) $f_1 + f_2 = 2, f_1 + f_2 + f_3 = 4, f_1 + f_2 + f_3 + f_4 = 7$

(C) **Statement:** $f_1 + f_2 + f_3 + \dots + f_n = f_{n+2} - 1, n \geq 3$

Proof by Induction:

Base Cases: $n = 1$

LHS: $f_1 = 1$

RHS: $f_3 - 1 = 2 - 1 = 1$

Since the LHS=RHS, the statement is true for $n = 1$.

Base Cases: $n = 2$

LHS: $f_1 + f_2 = 1 + 1 = 2$

RHS: $f_4 - 1 = 3 - 1 = 2$

Since the LHS=RHS, the statement is true for $n = 2$.

Induction Step: Fix for $k \geq 3$

Let the equation $f_1 + f_2 + f_3 + \dots + f_k = f_{k+2} - 1, k \geq 3$ be true for k .

We want to show the statement is true for $(k+1)$ by demonstrating the LHS=RHS.

Substituting $(k+1)$:

$$\underbrace{f_1 + f_2 + f_3 + \dots + f_k}_{\text{Induction Hypothesis.}} + f_{k+1} = f_{k+3} - 1$$

Replace the IH:

$$f_{k+2} - 1 + f_{k+1} = f_{k+3} - 1.$$

Add 1 to both sides:

$$f_{k+2} + f_{k+1} = f_{k+3}$$

By the definition of the Fibonacci sequence:

$$f_{k+3} = f_{k+2} + f_{k+1}$$

Thus:

$$f_{k+2} + f_{k+1} = f_{k+2} + f_{k+1}$$

—

$$\therefore f_1 + f_2 + f_3 + \dots + f_n = f_{n+2} - 1, n \geq 3$$

■

Additional Problems:

Question 1

Statement: Prove that for every natural number n , $3^n \geq 1 + 2^n$

Proof by Induction:

Base Case: $k = 1$

$$\text{LHS} = 3^1 = 3$$

$$\text{RHS} = 1 + 2^1 = 1 + 2 = 3$$

Since the LHS $\geq$ RHS the statement is true for $k = 1$.

Induction Step:

Let the statement $3^k \geq 2^k + 1$ be true for k .

We want to prove the statement is true for $(k + 1)$ by showing that the LHS $\geq$ RHS.

Substituting:

$$3^{k+1} \geq 2^{k+1} + 1$$

Rewrite:

$$3 \cdot \underbrace{3^k}_{\text{Inductive Hypothesis.}} \geq 2 \cdot 2^k + 1$$

Replace with IH:

$$3 \cdot (2^k + 1) \geq 2 \cdot 2^k + 1$$

Simplify further:

$$3 \cdot 2^k + 3 \geq 2 \cdot 2^k + 1$$

Subtract 1 from both sides:

$$3 \cdot 2^k + 2 \geq 2 \cdot 2^k$$

Finalize:

$$3 \cdot 2^{k+1} \geq 2^{k+1}$$

—

$\therefore \forall n, 3^n \geq 1 + 2^n$ since we have shown that the LHS $\geq$ RHS.

■

Question 2

Statement: Prove that for every number natural number n , $n^3 + 5n + 6$ is divisible by 3.

Proof by Induction:**Base Case:** $k = 1$

We want to show that our result is an integer multiple of 3.

$$1^3 + 5(1) + 6 = 6 + 6 = 12$$

$$12 = 4(3)$$

$\therefore$ the statement is true for $k = 1$.

Induction Step:

Let the statement $k^3 + 5k + 6$ be true for k .

We want to show also that it is true for $(k + 1)$.