

Advanced Engineering Mathematics

Mathematical Techniques for Engineering

Dr Athar Kharal

atharkharal.github.io

atharkharal@gmail.com

<https://pk.linkedin.com/in/atharkharal>

Theorem

(Laplace Transform of the Derivative $f^{(n)}$ of Any Order)

$$L\left(f^{(n)}\right) = s^n L(f) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$

Theorem

(Laplace Transform of the Derivative $f^{(n)}$ of Any Order)

$$L\left(f^{(n)}\right) = s^n L(f) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$

from which we have

$$L\left(f'\right) = sL(f) - f(0) \quad , \quad L\left(f''\right) = s^2 L(f) - sf(0) - f'(0)$$

Theorem

(Laplace Transform of the Derivative $f^{(n)}$ of Any Order)

$$L\left(f^{(n)}\right) = s^n L(f) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$

from which we have

$$L(f') = sL(f) - f(0) \quad , \quad L(f'') = s^2 L(f) - sf(0) - f'(0)$$

Theorem

(Laplace Transform of Integral) *We have*

$$L\left(\int_0^t f(\tau) d\tau\right) = \frac{1}{s} F(s)$$

Theorem

(Laplace Transform of the Derivative $f^{(n)}$ of Any Order)

$$L\left(f^{(n)}\right) = s^n L(f) - s^{n-1} f(0) - s^{n-2} f'(0) - \dots - f^{(n-1)}(0)$$

from which we have

$$L(f') = sL(f) - f(0) \quad , \quad L(f'') = s^2 L(f) - sf(0) - f'(0)$$

Theorem

(Laplace Transform of Integral) We have

$$L\left(\int_0^t f(\tau) d\tau\right) = \frac{1}{s} F(s), \quad \text{thus} \quad \int_0^t f(\tau) d\tau = L^{-1}\left(\frac{1}{s} F(s)\right).$$

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative of $f(t)$ involves the $f(t)$ as one of its terms due to oscillating derivative of \sin and \cos .

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative of $f(t)$ involves the $f(t)$ as one of its terms due to oscillating derivative of \sin and \cos . That is, $f'(t) = \sin \omega t + \omega t \cos \omega t$ and $f''(t) = 2\omega \cos \omega t - \omega^2 t \sin \omega t$.

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative of $f(t)$ involves the $f(t)$ as one of its terms due to oscillating derivative of \sin and \cos . That is, $f'(t) = \sin \omega t + \omega t \cos \omega t$ and $f''(t) = 2\omega \cos \omega t - \omega^2 t \sin \omega t$. Also $f(0) = 0$, $f'(0) = 0$. Hence we have

$$L(f'') = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative of $f(t)$ involves the $f(t)$ as one of its terms due to oscillating derivative of \sin and \cos . That is, $f'(t) = \sin \omega t + \omega t \cos \omega t$ and $f''(t) = 2\omega \cos \omega t - \omega^2 t \sin \omega t$. Also $f(0) = 0$, $f'(0) = 0$. Hence we have

$$L(f'') = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

$$\text{also } L(f'') = s^2 L(f) - sf(0) - f'(0)$$

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative of $f(t)$ involves the $f(t)$ as one of its terms due to oscillating derivative of \sin and \cos . That is, $f'(t) = \sin \omega t + \omega t \cos \omega t$ and $f''(t) = 2\omega \cos \omega t - \omega^2 t \sin \omega t$. Also $f(0) = 0$, $f'(0) = 0$. Hence we have

$$L(f'') = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

$$\text{also } L(f'') = s^2 L(f) - sf(0) - f'(0)$$

$$\text{implies } s^2 L(f) - sf(0) - f'(0) = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative of $f(t)$ involves the $f(t)$ as one of its terms due to oscillating derivative of \sin and \cos . That is, $f'(t) = \sin \omega t + \omega t \cos \omega t$ and $f''(t) = 2\omega \cos \omega t - \omega^2 t \sin \omega t$. Also $f(0) = 0$, $f'(0) = 0$. Hence we have

$$L(f'') = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

$$\text{also } L(f'') = s^2 L(f) - sf(0) - f'(0)$$

$$\text{implies } s^2 L(f) - sf(0) - f'(0) = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

$$s^2 L(t \sin \omega t) - 0 - 0 = 2\omega \frac{s}{s^2 + \omega^2} - \omega^2 L(t \sin \omega t)$$

Example

To find $L(t \sin \omega t)$, let $f(t) = t \sin \omega t$. We note that second derivative of $f(t)$ involves the $f(t)$ as one of its terms due to oscillating derivative of \sin and \cos . That is, $f'(t) = \sin \omega t + \omega t \cos \omega t$ and $f''(t) = 2\omega \cos \omega t - \omega^2 t \sin \omega t$. Also $f(0) = 0$, $f'(0) = 0$. Hence we have

$$L(f'') = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

$$\text{also } L(f'') = s^2 L(f) - sf(0) - f'(0)$$

$$\text{implies } s^2 L(f) - sf(0) - f'(0) = 2\omega L(\cos \omega t) - \omega^2 L(t \sin \omega t)$$

$$s^2 L(t \sin \omega t) - 0 - 0 = 2\omega \frac{s}{s^2 + \omega^2} - \omega^2 L(t \sin \omega t)$$

$$L(t \sin \omega t) = \frac{2\omega s}{(s^2 + \omega^2)^2} \quad \blacksquare$$

Example

Let $f(t) = \cos \omega t$

Example

Let $f(t) = \cos \omega t$ Then $f(0) = 1$, $f'(0) = 0$, $f''(t) = -\omega^2 \cos \omega t$.

Example

Let $f(t) = \cos \omega t$ Then $f(0) = 1$, $f'(0) = 0$, $f''(t) = -\omega^2 \cos \omega t$. From this and formula for $L(f'')$ we have

$$s^2 L(f) - sf(0) = L(f'') = L(-\omega^2 \cos \omega t)$$

Example

Let $f(t) = \cos \omega t$ Then $f(0) = 1$, $f'(0) = 0$, $f''(t) = -\omega^2 \cos \omega t$. From this and formula for $L(f'')$ we have

$$\begin{aligned} s^2 L(f) - sf(0) &= L(f'') = L(-\omega^2 \cos \omega t) \\ s^2 L(\cos \omega t) - s &= -\omega^2 L(\cos \omega t) \\ \text{solving } L(\cos \omega t) &= \frac{s}{s^2 + \omega^2} \end{aligned}$$

Example

Let $f(t) = \cos \omega t$ Then $f(0) = 1$, $f'(0) = 0$, $f''(t) = -\omega^2 \cos \omega t$. From this and formula for $L(f'')$ we have

$$\begin{aligned} s^2 L(f) - sf(0) &= L(f'') = L(-\omega^2 \cos \omega t) \\ s^2 L(\cos \omega t) - s &= -\omega^2 L(\cos \omega t) \\ \text{solving } L(\cos \omega t) &= \frac{s}{s^2 + \omega^2} \end{aligned}$$

Similarly let $g(t) = \sin \omega t$.

Example

Let $f(t) = \cos \omega t$. Then $f(0) = 1$, $f'(0) = 0$, $f''(t) = -\omega^2 \cos \omega t$. From this and formula for $L(f'')$ we have

$$\begin{aligned} s^2 L(f) - sf(0) &= L(f'') = L(-\omega^2 \cos \omega t) \\ s^2 L(\cos \omega t) - s &= -\omega^2 L(\cos \omega t) \\ \text{solving } L(\cos \omega t) &= \frac{s}{s^2 + \omega^2} \end{aligned}$$

Similarly let $g(t) = \sin \omega t$. Then $g(0) = 0$, $g'(t) = \omega \cos \omega t$, which gives

$$\begin{aligned} sL(g) - g(0) &= L(g') = \omega L(\cos \omega t) \\ \text{Hence } L(\sin \omega t) &= \frac{\omega}{s} L(\cos \omega t) = \frac{\omega}{s^2 + \omega^2} \quad \blacksquare \end{aligned}$$

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

The presence of term $\frac{1}{s^2}$

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

The presence of term $\frac{1}{s^2}$ indicates that it is double integration of a function whose $L(f(t)) = F(s) = \frac{1}{s - 2\pi}$.

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

The presence of term $\frac{1}{s^2}$ indicates that it is double integration of a function whose $L(f(t)) = F(s) = \frac{1}{s-2\pi}$. By table of LT we know $f(t) = L^{-1} \left(\frac{1}{s-2\pi} \right) = e^{2\pi t}$.

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

The presence of term $\frac{1}{s^2}$ indicates that it is double integration of a function whose $L(f(t)) = F(s) = \frac{1}{s-2\pi}$. By table of LT we know $f(t) = L^{-1} \left(\frac{1}{s-2\pi} \right) = e^{2\pi t}$. Hence we proceed as follows:

$$\begin{aligned} & L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right) \\ &= L^{-1} \left(\frac{20}{s^2 (s - 2\pi)} \right) = 20 L^{-1} \left(\frac{1}{s^2} \frac{1}{s - 2\pi} \right) \end{aligned}$$

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

The presence of term $\frac{1}{s^2}$ indicates that it is double integration of a function whose $L(f(t)) = F(s) = \frac{1}{s-2\pi}$. By table of LT we know $f(t) = L^{-1} \left(\frac{1}{s-2\pi} \right) = e^{2\pi t}$. Hence we proceed as follows:

$$\begin{aligned} & L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right) \\ &= L^{-1} \left(\frac{20}{s^2 (s - 2\pi)} \right) = 20 L^{-1} \left(\frac{1}{s^2} \frac{1}{s - 2\pi} \right) = 20 \int_0^\tau \left(\int_0^\tau e^{2\pi\tau} d\tau \right) \\ &= 20 \int_0^\tau \left(\frac{1}{2\pi} (e^{2\pi\tau} - 1) \right) d\tau \end{aligned}$$

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

The presence of term $\frac{1}{s^2}$ indicates that it is double integration of a function whose $L(f(t)) = F(s) = \frac{1}{s-2\pi}$. By table of LT we know $f(t) = L^{-1} \left(\frac{1}{s-2\pi} \right) = e^{2\pi t}$. Hence we proceed as follows:

$$\begin{aligned} & L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right) \\ &= L^{-1} \left(\frac{20}{s^2 (s - 2\pi)} \right) = 20 L^{-1} \left(\frac{1}{s^2} \frac{1}{s - 2\pi} \right) = 20 \int_0^\tau \left(\int_0^\tau e^{2\pi\tau} d\tau \right) \\ &= 20 \int_0^\tau \left(\frac{1}{2\pi} (e^{2\pi\tau} - 1) \right) d\tau = 20 \left(-\frac{1}{4\pi^2} (2\pi\tau - e^{2\pi\tau} + 1) \right) \end{aligned}$$

Example (10e-6.2-24)

Find $L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right)$ using LT of Integral?

Note that simplification yields

$$\frac{20}{s^3 - 2\pi s^2} = \frac{20}{s^2 (s - 2\pi)} = 20 \frac{1}{s^2} \frac{1}{s - 2\pi}$$

The presence of term $\frac{1}{s^2}$ indicates that it is double integration of a function whose $L(f(t)) = F(s) = \frac{1}{s-2\pi}$. By table of LT we know $f(t) = L^{-1} \left(\frac{1}{s-2\pi} \right) = e^{2\pi t}$. Hence we proceed as follows:

$$\begin{aligned} & L^{-1} \left(\frac{20}{s^3 - 2\pi s^2} \right) \\ &= L^{-1} \left(\frac{20}{s^2 (s - 2\pi)} \right) = 20 L^{-1} \left(\frac{1}{s^2} \frac{1}{s - 2\pi} \right) = 20 \int_0^\tau \left(\int_0^\tau e^{2\pi\tau} d\tau \right) \\ &= 20 \int_0^\tau \left(\frac{1}{2\pi} (e^{2\pi\tau} - 1) \right) d\tau = 20 \left(-\frac{1}{4\pi^2} (2\pi\tau - e^{2\pi\tau} + 1) \right) \\ &= \frac{5}{\pi^2} e^{2(\pi\tau)} - \frac{10}{\pi} \tau - \frac{5}{\pi^2} \quad \blacksquare \end{aligned}$$

Example

Using Laplace transform of integral find the inverse of $\frac{1}{s(s^2+\omega^2)}$ and $\frac{1}{s^2(s^2+\omega^2)}$?

Example

Using Laplace transform of integral find the inverse of $\frac{1}{s(s^2+\omega^2)}$ and $\frac{1}{s^2(s^2+\omega^2)}$?

From the table we have $L^{-1} \left(\frac{1}{(s^2+\omega^2)} \right) = \frac{\sin \omega t}{\omega}$

Example

Using Laplace transform of integral find the inverse of $\frac{1}{s(s^2+\omega^2)}$ and $\frac{1}{s^2(s^2+\omega^2)}$?

From the table we have $L^{-1} \left(\frac{1}{(s^2+\omega^2)} \right) = \frac{\sin \omega t}{\omega}$

$$L^{-1} \left(\frac{1}{s(s^2+\omega^2)} \right) = \int_0^t \frac{\sin \omega \tau}{\omega} d\tau$$

Example

Using Laplace transform of integral find the inverse of $\frac{1}{s(s^2+\omega^2)}$ and $\frac{1}{s^2(s^2+\omega^2)}$?

From the table we have $L^{-1} \left(\frac{1}{(s^2+\omega^2)} \right) = \frac{\sin \omega t}{\omega}$

$$L^{-1} \left(\frac{1}{s(s^2+\omega^2)} \right) = \int_0^t \frac{\sin \omega \tau}{\omega} d\tau = \frac{1}{\omega^2} (1 - \cos \omega t)$$

Example

Using Laplace transform of integral find the inverse of $\frac{1}{s(s^2+\omega^2)}$ and $\frac{1}{s^2(s^2+\omega^2)}$?

From the table we have $L^{-1} \left(\frac{1}{(s^2+\omega^2)} \right) = \frac{\sin \omega t}{\omega}$

$$L^{-1} \left(\frac{1}{s(s^2+\omega^2)} \right) = \int_0^t \frac{\sin \omega \tau}{\omega} d\tau = \frac{1}{\omega^2} (1 - \cos \omega t)$$

Integrating this result again, we obtain

$$L^{-1} \left(\frac{1}{s^2(s^2+\omega^2)} \right) = \frac{1}{\omega^2} \int_0^t (1 - \cos \omega t)$$

Example

Using Laplace transform of integral find the inverse of $\frac{1}{s(s^2+\omega^2)}$ and $\frac{1}{s^2(s^2+\omega^2)}$?

From the table we have $L^{-1} \left(\frac{1}{(s^2+\omega^2)} \right) = \frac{\sin \omega t}{\omega}$

$$L^{-1} \left(\frac{1}{s(s^2+\omega^2)} \right) = \int_0^t \frac{\sin \omega \tau}{\omega} d\tau = \frac{1}{\omega^2} (1 - \cos \omega t)$$

Integrating this result again, we obtain

$$\begin{aligned} L^{-1} \left(\frac{1}{s^2(s^2+\omega^2)} \right) &= \frac{1}{\omega^2} \int_0^t (1 - \cos \omega \tau) d\tau \\ &= \left[\frac{\tau}{\omega^2} - \frac{\sin \omega \tau}{\omega^3} \right]_0^t = \frac{t}{\omega^2} - \frac{\sin \omega t}{\omega^3} \quad \blacksquare \end{aligned}$$

Differential Equations, Initial Value Problems (IVP)

Laplace transform method solves ODEs and initial value problems using following steps:

- 1 Setting up the subsidiary equation.

Differential Equations, Initial Value Problems (IVP)

Laplace transform method solves ODEs and initial value problems using following steps:

- 1 Setting up the subsidiary equation.
- 2 Solution of the subsidiary equation by algebra.

Differential Equations, Initial Value Problems (IVP)

Laplace transform method solves ODEs and initial value problems using following steps:

- ① Setting up the subsidiary equation.
- ② Solution of the subsidiary equation by algebra.
- ③ Inversion of Y to obtain $y = L^{-1}(Y)$.

Differential Equations, Initial Value Problems (IVP)


Laplace transform method solves ODEs and initial value problems using following steps:

- 1 Setting up the subsidiary equation.
- 2 Solution of the subsidiary equation by algebra.
- 3 Inversion of Y to obtain $y = L^{-1}(Y)$.

Differential Equations, Initial Value Problems (IVP)

Laplace transform method solves ODEs and initial value problems using following steps:

- ① Setting up the subsidiary equation.
- ② Solution of the subsidiary equation by algebra.
- ③ Inversion of Y to obtain $y = L^{-1}(Y)$.



Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

$$\frac{d^2}{dt^2} y(t) - y(t) = t ; \quad y(0) = 1, y'(0) = 1$$

Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

$$\frac{d^2}{dt^2} y(t) - y(t) = t ; \quad y(0) = 1, y'(0) = 1$$

$$s^2 L(y(t)) - y'(0) - sy(0) - L(y(t)) = \frac{1}{s^2}$$

Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

$$\frac{d^2}{dt^2} y(t) - y(t) = t ; \quad y(0) = 1, y'(0) = 1$$

$$s^2 L(y(t)) - y'(0) - sy(0) - L(y(t)) = \frac{1}{s^2}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + y'(0) + sy(0)}{s^2 - 1}$$

Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

$$\frac{d^2}{dt^2} y(t) - y(t) = t ; \quad y(0) = 1, y'(0) = 1$$

$$s^2 L(y(t)) - y'(0) - sy(0) - L(y(t)) = \frac{1}{s^2}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + y'(0) + sy(0)}{s^2 - 1}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + 1 + s}{s^2 - 1}$$

Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

$$\frac{d^2}{dt^2} y(t) - y(t) = t ; \quad y(0) = 1, y'(0) = 1$$

$$s^2 L(y(t)) - y'(0) - sy(0) - L(y(t)) = \frac{1}{s^2}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + y'(0) + sy(0)}{s^2 - 1}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + 1 + s}{s^2 - 1} = \frac{s^3 + s^2 + 1}{s^2(s^2 - 1)}$$

Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

$$\frac{d^2}{dt^2} y(t) - y(t) = t ; \quad y(0) = 1, y'(0) = 1$$

$$s^2 L(y(t)) - y'(0) - sy(0) - L(y(t)) = \frac{1}{s^2}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + y'(0) + sy(0)}{s^2 - 1}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + 1 + s}{s^2 - 1} = \frac{s^3 + s^2 + 1}{s^2(s^2 - 1)}$$

$$L(y(t)) = -\frac{1}{s^2} - \frac{1}{2(s+1)} + \frac{3}{2(s-1)}$$

Example

Using LT, solve the IVP : $y'' - y = t$; $y(0) = y'(0) = 1$?

$$\frac{d^2}{dt^2} y(t) - y(t) = t ; \quad y(0) = 1, y'(0) = 1$$

$$s^2 L(y(t)) - y'(0) - sy(0) - L(y(t)) = \frac{1}{s^2}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + y'(0) + sy(0)}{s^2 - 1}$$

$$L(y(t)) = \frac{\frac{1}{s^2} + 1 + s}{s^2 - 1} = \frac{s^3 + s^2 + 1}{s^2(s^2 - 1)}$$

$$L(y(t)) = -\frac{1}{s^2} - \frac{1}{2(s+1)} + \frac{3}{2(s-1)}$$

$$y(t) = -t + \cosh(t) + 2 \sinh(t) \quad \blacksquare$$

Example

Solve the IVP $y'' + 9y = 10e^{-t}$; $y(0) = y'(0) = 0$ using Laplace Transforms.

Example

Solve the IVP $y'' + 9y = 10e^{-t}$; $y(0) = y'(0) = 0$ using Laplace Transforms.

$$\frac{d^2}{dt^2}y(t) + 9y(t) = 10e^{-t}$$

Example

Solve the IVP $y'' + 9y = 10e^{-t}$; $y(0) = y'(0) = 0$ using Laplace Transforms.

$$\frac{d^2}{dt^2} y(t) + 9y(t) = 10e^{-t}$$

$$s^2 L(y(t)) - y'(0) - sy(0) + 9L(y(t)) = \frac{10}{s+1}$$

Isolating Laplace term $L(y(t)) = \frac{1}{s^2+9} \left(\frac{10}{s+1} + y'(0) + sy(0) \right)$

Example

Solve the IVP $y'' + 9y = 10e^{-t}$; $y(0) = y'(0) = 0$ using Laplace Transforms.

$$\frac{d^2}{dt^2}y(t) + 9y(t) = 10e^{-t}$$

$$s^2L(y(t)) - y'(0) - sy(0) + 9L(y(t)) = \frac{10}{s+1}$$

Isolating Laplace term $L(y(t)) = \frac{1}{s^2+9} \left(\frac{10}{s+1} + y'(0) + sy(0) \right)$

$$L(y(t)) = \frac{10}{(s+1)(s^2+9)} \quad \text{by using ICs}$$

Example

Solve the IVP $y'' + 9y = 10e^{-t}$; $y(0) = y'(0) = 0$ using Laplace Transforms.

$$\frac{d^2}{dt^2}y(t) + 9y(t) = 10e^{-t}$$

$$s^2L(y(t)) - y'(0) - sy(0) + 9L(y(t)) = \frac{10}{s+1}$$

Isolating Laplace term $L(y(t)) = \frac{1}{s^2+9} \left(\frac{10}{s+1} + y'(0) + sy(0) \right)$

$$L(y(t)) = \frac{10}{(s+1)(s^2+9)} \quad \text{by using ICs}$$

$$L(y(t)) = \frac{1}{s+1} - \frac{s}{s^2+9} - \frac{1}{s^2+9} \quad \text{by partial fractions (DIY)}$$

Example

Solve the IVP $y'' + 9y = 10e^{-t}$; $y(0) = y'(0) = 0$ using Laplace Transforms.

$$\frac{d^2}{dt^2}y(t) + 9y(t) = 10e^{-t}$$

$$s^2L(y(t)) - y'(0) - sy(0) + 9L(y(t)) = \frac{10}{s+1}$$

Isolating Laplace term $L(y(t)) = \frac{1}{s^2+9} \left(\frac{10}{s+1} + y'(0) + sy(0) \right)$

$$L(y(t)) = \frac{10}{(s+1)(s^2+9)} \quad \text{by using ICs}$$

$$L(y(t)) = \frac{1}{s+1} - \frac{s}{s^2+9} - \frac{1}{s^2+9} \quad \text{by partial fractions (DIY)}$$

$$y(t) = e^{-t} - \cos(3t) + \frac{1}{3}\sin(3t) \quad \blacksquare$$

Example (10e-6.1-10)

Solve $y'' + 3y + 2.25y = 9t^3 + 64$ with

ICs $y(0) = 1, y'(0) = 31.5$. use Laplace Transforms.

Example (10e-6.1-10)

Solve $y'' + 3y + 2.25y = 9t^3 + 64$ with

ICs $y(0) = 1, y'(0) = 31.5$. use Laplace Transforms.

$$\frac{d^2}{dt^2} y(t) + 3 \frac{d}{dt} y(t) + 2.25 y(t) = 9t^3 + 64 ; \quad y(0) = 1, y'(0) = 31.5$$

Example (10e-6.1-10)

Solve $y'' + 3y + 2.25y = 9t^3 + 64$ with

ICs $y(0) = 1, y'(0) = 31.5$. use Laplace Transforms.

$$\frac{d^2}{dt^2} y(t) + 3 \frac{d}{dt} y(t) + 2.25 y(t) = 9t^3 + 64 ; \quad y(0) = 1, y'(0) = 31.5$$

$$s^2 Y - y'(0) - sy(0) + 3sY - 3y(0) + 2.25 Y = \frac{54}{s^4} + \frac{64}{s}$$

$$Y = \frac{\frac{54}{s^4} + \frac{64}{s} + y'(0) + sy(0) + 3y(0)}{s^2 + 3s + 2.25}$$

Example (10e-6.1-10)

Solve $y'' + 3y' + 2.25y = 9t^3 + 64$ with

ICs $y(0) = 1, y'(0) = 31.5$. use Laplace Transforms.

$$\frac{d^2}{dt^2} y(t) + 3 \frac{d}{dt} y(t) + 2.25 y(t) = 9t^3 + 64 ; \quad y(0) = 1, y'(0) = 31.5$$

$$s^2 Y - y'(0) - sy(0) + 3sY - 3y(0) + 2.25 Y = \frac{54}{s^4} + \frac{64}{s}$$

$$Y = \frac{\frac{54}{s^4} + \frac{64}{s} + y'(0) + sy(0) + 3y(0)}{s^2 + 3s + 2.25}$$

$$Y = \frac{\frac{54}{s^4} + \frac{64}{s} + 34.5 + s}{s^2 + 3s + 2.25} \quad \text{using ICs}$$

$$Y = \frac{54 + 64s^3 + 34.5s^4 + s^5}{s^6 + 3s^5 + 2.25s^4} = \frac{54 + 64s^3 + 34.5s^4 + s^5}{\frac{1}{4}s^4(2s+3)^2}$$

Example (10e-6.1-10)

Solve $y'' + 3y' + 2.25y = 9t^3 + 64$ with

ICs $y(0) = 1, y'(0) = 31.5$. use Laplace Transforms.

$$\frac{d^2}{dt^2} y(t) + 3 \frac{d}{dt} y(t) + 2.25 y(t) = 9t^3 + 64 ; \quad y(0) = 1, y'(0) = 31.5$$

$$s^2 Y - y'(0) - sy(0) + 3sY - 3y(0) + 2.25 Y = \frac{54}{s^4} + \frac{64}{s}$$

$$Y = \frac{\frac{54}{s^4} + \frac{64}{s} + y'(0) + sy(0) + 3y(0)}{s^2 + 3s + 2.25}$$

$$Y = \frac{\frac{54}{s^4} + \frac{64}{s} + 34.5 + s}{s^2 + 3s + 2.25} \quad \text{using ICs}$$

$$Y = \frac{54 + 64s^3 + 34.5s^4 + s^5}{s^6 + 3s^5 + 2.25s^4} = \frac{54 + 64s^3 + 34.5s^4 + s^5}{\frac{1}{4}s^4(2s+3)^2}$$

By partial fractions(DIY) $Y = \frac{2}{2s+3} + \frac{4}{(2s+3)^2} + \frac{32}{s^2} - \frac{32}{s^3} + \frac{24}{s^4}$

$$Y = L(y(t)) = \frac{1}{s+1.5} + \frac{1}{(s+1.5)^2} + \frac{32}{s^2} - \frac{32}{s^3} + \frac{24}{s^4}$$

$$Y = L(y(t)) = \frac{1}{s+1.5} + \frac{1}{(s+1.5)^2} + \frac{32}{s^2} - \frac{32}{s^3} + \frac{24}{s^4}$$

$$y(t) = L^{-1} \left(\frac{1}{s+1.5} + \frac{1}{(s+1.5)^2} + \frac{32}{s^2} - \frac{32}{s^3} + \frac{24}{s^4} \right)$$

$$y(t) = e^{-1.5t} + te^{-1.5t} + 32t - 16t^2 + 4t^3$$

$$y(t) = 4t^3 - 16t^2 + 32t + (t+1)e^{-1.5t} \quad \blacksquare$$

Example (Shifted Data Problem: where ICs are given at non-zero point)

Example (Shifted Data Problem: where ICs are given at non-zero point)

Find solution of the IVP:

$$y'' + y = 2t \quad ; \quad y\left(\frac{1}{4}\pi\right) = \frac{1}{2}\pi, \quad y'\left(\frac{1}{4}\pi\right) = 2 - \sqrt{2}?$$

Example (Shifted Data Problem: where ICs are given at non-zero point)

Find solution of the IVP:

$$y'' + y = 2t \quad ; \quad y\left(\frac{1}{4}\pi\right) = \frac{1}{2}\pi, \quad y'\left(\frac{1}{4}\pi\right) = 2 - \sqrt{2}?$$

Solution: Put $\tilde{t} = t - \frac{\pi}{4}$

Example (Shifted Data Problem: where ICs are given at non-zero point)

Find solution of the IVP:

$$y'' + y = 2t \quad ; \quad y\left(\frac{1}{4}\pi\right) = \frac{1}{2}\pi, \quad y'\left(\frac{1}{4}\pi\right) = 2 - \sqrt{2}?$$

Solution: Put $\tilde{t} = t - \frac{\pi}{4}$ to obtain

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + y(\tilde{t}) = 2\tilde{t} + \frac{\pi}{2} \quad ; \quad y(0) = \frac{\pi}{2}, \quad y'(0) = 2 - \sqrt{2}$$

Example (Shifted Data Problem: where ICs are given at non-zero point)

Find solution of the IVP:

$$y'' + y = 2t \quad ; \quad y\left(\frac{1}{4}\pi\right) = \frac{1}{2}\pi, \quad y'\left(\frac{1}{4}\pi\right) = 2 - \sqrt{2}?$$

Solution: Put $\tilde{t} = t - \frac{\pi}{4}$ to obtain

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + y(\tilde{t}) = 2\tilde{t} + \frac{\pi}{2} \quad ; \quad y(0) = \frac{\pi}{2}, \quad y'(0) = 2 - \sqrt{2}$$

Taking LT $s^2 \tilde{Y} - y'(0) - sy(0) + \tilde{Y} = \frac{\pi}{2} \frac{1}{s} + 2 \frac{1}{s^2}$ where $\tilde{Y} = L(y(\tilde{t}))$

Example (Shifted Data Problem: where ICs are given at non-zero point)

Find solution of the IVP:

$$y'' + y = 2t \quad ; \quad y\left(\frac{1}{4}\pi\right) = \frac{1}{2}\pi, \quad y'\left(\frac{1}{4}\pi\right) = 2 - \sqrt{2}$$

Solution: Put $\tilde{t} = t - \frac{\pi}{4}$ to obtain

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + y(\tilde{t}) = 2\tilde{t} + \frac{\pi}{2} \quad ; \quad y(0) = \frac{\pi}{2}, \quad y'(0) = 2 - \sqrt{2}$$

Taking LT $s^2 \tilde{Y} - y'(0) - sy(0) + \tilde{Y} = \frac{\pi}{2} \frac{1}{s} + 2 \frac{1}{s^2}$ where $\tilde{Y} = L(y(\tilde{t}))$

Isolating $\tilde{Y} = \frac{1}{s^2+1} \left(\frac{\pi}{2} \frac{1}{s} + 2 \frac{1}{s^2} + y'(0) + sy(0) \right)$

Using ICs $\tilde{Y} = \frac{1}{s^2+1} \left(\frac{\pi}{2} \frac{1}{s} + 2 \frac{1}{s^2} + (2 - \sqrt{2}) + \frac{\pi}{2} s \right)$

$$= \frac{1}{2} \frac{s^3\pi - 2\sqrt{2}s^2 + s\pi + 4s^2 + 4}{s^2(s^2+1)}$$

By partial fractions(DIY) $\tilde{Y} = L(y(\tilde{t})) = \frac{\pi}{2} \frac{1}{s} + \frac{2}{s^2} - \frac{\sqrt{2}}{s^2+1}$

By partial fractions(DIY) $\tilde{Y} = L(y(\tilde{t})) = \frac{\pi}{2} \frac{1}{s} + \frac{2}{s^2} - \frac{\sqrt{2}}{s^2+1}$

$$y(\tilde{t}) = L^{-1}\left(\frac{\pi}{2} \frac{1}{s} + \frac{2}{s^2} - \frac{\sqrt{2}}{s^2+1}\right)$$

$$y(\tilde{t}) = \frac{\pi}{2} + 2\tilde{t} - \sqrt{2}\sin(\tilde{t})$$

By partial fractions(DIY) $\tilde{Y} = L(y(\tilde{t})) = \frac{\pi}{2} \frac{1}{s} + \frac{2}{s^2} - \frac{\sqrt{2}}{s^2+1}$

$$y(\tilde{t}) = L^{-1}\left(\frac{\pi}{2} \frac{1}{s} + \frac{2}{s^2} - \frac{\sqrt{2}}{s^2+1}\right)$$

$$y(\tilde{t}) = \frac{\pi}{2} + 2\tilde{t} - \sqrt{2}\sin(\tilde{t})$$

Putting back $\tilde{t} = t - \frac{\pi}{4}$

By partial fractions(DIY) $\tilde{Y} = L(y(\tilde{t})) = \frac{\pi}{2} \frac{1}{s} + \frac{2}{s^2} - \frac{\sqrt{2}}{s^2+1}$

$$y(\tilde{t}) = L^{-1}\left(\frac{\pi}{2} \frac{1}{s} + \frac{2}{s^2} - \frac{\sqrt{2}}{s^2+1}\right)$$

$$y(\tilde{t}) = \frac{\pi}{2} + 2\tilde{t} - \sqrt{2}\sin(\tilde{t})$$

Putting back $\tilde{t} = t - \frac{\pi}{4}$ $y(t) = 2t + \sqrt{2}\cos\left(t + \frac{\pi}{4}\right)$ ■

Example (10e-6.2-14)

The IVP $y'' + 2y' + 5y = 50t - 100$; $y(2) = -4, y'(2) = 14$
is a shifted data problem.

Example (10e-6.2-14)

The IVP $y'' + 2y' + 5y = 50t - 100$; $y(2) = -4, y'(2) = 14$
is a shifted data problem. So put \tilde{t}

Example (10e-6.2-14)

The IVP $y'' + 2y' + 5y = 50t - 100$; $y(2) = -4$, $y'(2) = 14$ is a shifted data problem. So put $\tilde{t} = t - 2$ and obtain

$$\frac{d^2}{d\tilde{t}^2} y(\tilde{t}) + 2 \frac{d}{d\tilde{t}} y(\tilde{t}) + 5 y(\tilde{t}) = 50\tilde{t} \quad ; \quad y(0) = -4, y'(0) = 14$$

Example (10e-6.2-14)

The IVP $y'' + 2y' + 5y = 50t - 100$; $y(2) = -4$, $y'(2) = 14$ is a shifted data problem. So put $\tilde{t} = t - 2$ and obtain

$$\frac{d^2}{d\tilde{t}^2} y(\tilde{t}) + 2 \frac{d}{d\tilde{t}} y(\tilde{t}) + 5 y(\tilde{t}) = 50\tilde{t} \quad ; \quad y(0) = -4, y'(0) = 14$$

$$s^2 \tilde{Y} - y'(0) - sy(0) + 2s\tilde{Y} - 2y(0) + 5\tilde{Y} = 50 \frac{1}{s^2} \quad \text{where}$$

$$\tilde{Y} = L(y(\tilde{t}))$$

$$\tilde{Y} = \frac{1}{s^2 + 2s + 5} \left(50 \frac{1}{s^2} + y'(0) + sy(0) + 2y(0) \right)$$

Example (10e-6.2-14)

The IVP $y'' + 2y' + 5y = 50t - 100$; $y(2) = -4$, $y'(2) = 14$ is a shifted data problem. So put $\tilde{t} = t - 2$ and obtain

$$\frac{d^2}{d\tilde{t}^2} y(\tilde{t}) + 2 \frac{d}{d\tilde{t}} y(\tilde{t}) + 5 y(\tilde{t}) = 50\tilde{t} \quad ; \quad y(0) = -4, y'(0) = 14$$

$$s^2 \tilde{Y} - y'(0) - sy(0) + 2s\tilde{Y} - 2y(0) + 5\tilde{Y} = 50 \frac{1}{s^2} \quad \text{where}$$

$$\tilde{Y} = L(y(\tilde{t}))$$

$$\tilde{Y} = \frac{1}{s^2 + 2s + 5} \left(50 \frac{1}{s^2} + y'(0) + sy(0) + 2y(0) \right)$$

$$\tilde{Y} = \frac{1}{s^2 + 2s + 5} \left(50 \frac{1}{s^2} + 6 - 4s \right)$$

$$\tilde{Y} = -2 \frac{2s^3 - 3s^2 - 25}{s^2(s^2 + 2s + 5)}$$

By partial fractions(DIY) $\tilde{Y} = -\frac{4}{s} + \frac{4}{(s+1)^2+4} + \frac{10}{s^2}$

$$y(\tilde{t}) = -4 + 2e^{-\tilde{t}} \sin(2\tilde{t}) + 10\tilde{t}$$

Replacing back

$$\tilde{t} = t - 2 \quad y(t) = -24 + 2e^{-t+2} \sin(2t - 4) + 10t \quad \blacksquare$$

Q (10e-6.2-15)

For the IVP $y'' + 3y' - 4y = 6e^{2t-3}$; $y(1.5) = 4, y'(1.5) = 5$
put \tilde{t}

Q (10e-6.2-15)

For the IVP $y'' + 3y' - 4y = 6e^{2t-3}$; $y(1.5) = 4, y'(1.5) = 5$
put $\tilde{t} = t - 1.5$

Q (10e-6.2-15)

For the IVP $y'' + 3y' - 4y = 6e^{2t-3}$; $y(1.5) = 4, y'(1.5) = 5$
 put $\tilde{t} = t - 1.5$ to transform the IVP as follows:

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + 3\frac{d}{d\tilde{t}}y(\tilde{t}) - 4y(\tilde{t}) = 6e^{2\tilde{t}} \quad ; \quad y(0) = 4, y'(0) = 5$$

Q (10e-6.2-15)

For the IVP $y'' + 3y' - 4y = 6e^{2t-3}$; $y(1.5) = 4, y'(1.5) = 5$
 put $\tilde{t} = t - 1.5$ to transform the IVP as follows:

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + 3\frac{d}{d\tilde{t}}y(\tilde{t}) - 4y(\tilde{t}) = 6e^{2\tilde{t}} \quad ; \quad y(0) = 4, y'(0) = 5$$

$$s^2\tilde{Y} - y'(0) - sy(0) + 3s\tilde{Y} - 3y(0) - 4\tilde{Y} = \frac{6}{s-2} \quad \text{where}$$

$$\tilde{Y} = L(y(\tilde{t}))$$

$$\tilde{Y} = \frac{1}{s^2+3s-4} \left(\frac{6}{s-2} + y'(0) + sy(0) + 3y(0) \right)$$

Using ICs $\tilde{Y} = \frac{1}{s^2+3s-4} \left(\frac{6}{s-2} + 17 + 4s \right)$

$$\tilde{Y} = \frac{4s-7}{(s-1)(s-2)}$$

By partial fractions $\tilde{Y} = \frac{3}{s-1} + \frac{1}{s-2}$

$$y(\tilde{t}) = 3e^{\tilde{t}} + e^{2\tilde{t}}$$

Replacing back $\tilde{t} = t - 1.5$, $y(t) = 3e^{(t-1.5)} + e^{2(t-1.5)}$ ■