

Advanced Engineering Mathematics

Mathematical Techniques for Engineering

Dr Athar Kharal

Humanities and Science Department
College of Aeronautical Engineering
National University of Sciences and Tecnology (NUST), Pakistan

Definition: Heaviside Unit Step Function

The unit step function or Heaviside function $u(t - a)$ is defined as:


$$u(t - a) = \begin{cases} 0 & \text{if } t < a \\ 1 & \text{if } t > a \end{cases} \quad ; \quad a \geq 0$$

Definition: Heaviside Unit Step Function

The unit step function or Heaviside function $u(t - a)$ is defined as:

$$u(t - a) = \begin{cases} 0 & \text{if } t < a \\ 1 & \text{if } t > a \end{cases} ; a \geq 0$$

Its graph is given as




Heaviside Unit Step Function: Explained

Let $f(t) = 0$ for all negative t . Then $f(t - a)u(t - a)$ with $a > 0$ is $f(t)$ shifted (translated) to the right by the amount a .

Heaviside Unit Step Function: Explained


Let $f(t) = 0$ for all negative t . Then $f(t - a)u(t - a)$ with $a > 0$ is $f(t)$ shifted (translated) to the right by the amount a .

(A) $f(t) = 5 \sin t$ (B) $f(t)u(t-2)$ (C) $f(t-2)u(t-2)$


Effects of the unit step function: (A) Given function.

(B) Switching off and on. (C) Shift.

Heaviside Unit Step Function: Explained



(A) $k[u(t-1) - 2u(t-4) + u(t-6)]$



(B) $4 \sin\left(\frac{1}{2}\pi t\right)[u(t) - u(t-2) + u(t-4) - u(t-6) + \dots]$

Use of many unit step functions.

Laplace Transform of Unit Step Function

LT of Heaviside unit step function may be found as

$$\begin{aligned}L(u(t-a)) &= \int_0^{\infty} e^{-st} u(t-a) dt \\&= \int_0^{\infty} e^{-st} (1) dt \quad \text{since } a \geq 0 \\&= \left[\frac{e^{-st}}{s} \right]_{t=a}^{\infty} \\&= \frac{e^{-as}}{s} \quad \blacksquare\end{aligned}$$

Second Shifting Theorem; t-Shifting

Theorem: (Second Shifting Theorem; t-Shifting) If $f(t)$ has the transform $F(s)$, then the 'shifted function'

$$\tilde{f}(t) = f(t - a) u(t - a)$$

Second Shifting Theorem; t-Shifting

Theorem: (Second Shifting Theorem; t-Shifting) If $f(t)$ has the transform $F(s)$, then the 'shifted function'

$$\tilde{f}(t) = f(t-a) u(t-a) = \begin{cases} 0 & \text{if } t < a \\ f(t-a) & \text{if } t > a \end{cases}$$

has the transform $e^{-as} F(s)$.

Second Shifting Theorem; t-Shifting

Theorem: (Second Shifting Theorem; t-Shifting) If $f(t)$ has the transform $F(s)$, then the 'shifted function'

$$\tilde{f}(t) = f(t-a) u(t-a) = \begin{cases} 0 & \text{if } t < a \\ f(t-a) & \text{if } t > a \end{cases}$$

has the transform $e^{-as} F(s)$. Equivalently we have

$$L(f(t-a) u(t-a)) = e^{-as} F(s)$$

Second Shifting Theorem; t-Shifting

Theorem: (Second Shifting Theorem; t-Shifting) If $f(t)$ has the transform $F(s)$, then the 'shifted function'

$$\tilde{f}(t) = f(t-a) u(t-a) = \begin{cases} 0 & \text{if } t < a \\ f(t-a) & \text{if } t > a \end{cases}$$

has the transform $e^{-as} F(s)$. Equivalently we have

$$L(f(t-a) u(t-a)) = e^{-as} F(s) \text{ and also}$$

$$L^{-1}(e^{-as} F(s)) = f(t-a) u(t-a).$$

Second Shifting Theorem; t-Shifting

Theorem: (Second Shifting Theorem; t-Shifting) If $f(t)$ has the transform $F(s)$, then the 'shifted function'

$$\tilde{f}(t) = f(t-a) u(t-a) = \begin{cases} 0 & \text{if } t < a \\ f(t-a) & \text{if } t > a \end{cases}$$

has the transform $e^{-as} F(s)$. Equivalently we have

$$L(f(t-a) u(t-a)) = e^{-as} F(s) \text{ and also}$$

$$L^{-1}(e^{-as} F(s)) = f(t-a) u(t-a).$$

Remark: Practically the t-shifting theorem amounts to this: if we know $F(s)$, we can obtain the transform of $f(t-a) u(t-a)$ by just multiplying $F(s)$ by e^{-as} .

Second Shifting Theorem; t-Shifting

Theorem: (Second Shifting Theorem; t-Shifting) If $f(t)$ has the transform $F(s)$, then the 'shifted function'

$$\tilde{f}(t) = f(t-a) u(t-a) = \begin{cases} 0 & \text{if } t < a \\ f(t-a) & \text{if } t > a \end{cases}$$

has the transform $e^{-as} F(s)$. Equivalently we have

$$L(f(t-a) u(t-a)) = e^{-as} F(s) \text{ and also}$$

$$L^{-1}(e^{-as} F(s)) = f(t-a) u(t-a).$$

Remark: Practically the t-shifting theorem amounts to this: if we know $F(s)$, we can obtain the transform of $f(t-a) u(t-a)$ by just multiplying $F(s)$ by e^{-as} .

If the conversion of $f(t)$ to $f(t-a)$ is difficult, we may use following form as well:

$$L(f(t) u(t-a)) = e^{-as} L(f(t+a))$$

Question: Find LT of the piecewise function

$$f(t) = \begin{cases} 2 & \text{if } 0 < t < 1 \\ \frac{t^2}{2} & \text{if } 1 < t < \frac{\pi}{2} \\ \cos t & \text{if } t > \frac{\pi}{2} \end{cases}$$

Question: Find LT of the piecewise function

$$f(t) = \begin{cases} 2 & \text{if } 0 < t < 1 \\ \frac{t^2}{2} & \text{if } 1 < t < \frac{\pi}{2} \\ \cos t & \text{if } t > \frac{\pi}{2} \end{cases}$$

Solution: Transform $f(t)$ in terms of unit-step functions as

$$f(t) = \underbrace{2(u(t) - u(t-1))}_{\text{First piece}}$$

Question: Find LT of the piecewise function

$$f(t) = \begin{cases} 2 & \text{if } 0 < t < 1 \\ \frac{t^2}{2} & \text{if } 1 < t < \frac{\pi}{2} \\ \cos t & \text{if } t > \frac{\pi}{2} \end{cases}$$

Solution: Transform $f(t)$ in terms of unit-step functions as

$$f(t) = \underbrace{2(u(t) - u(t-1))}_{\text{First piece}} + \underbrace{\frac{t^2}{2} \left(u(t-1) - u\left(t - \frac{\pi}{2}\right) \right)}_{\text{Second piece}}$$

Question: Find LT of the piecewise function

$$f(t) = \begin{cases} 2 & \text{if } 0 < t < 1 \\ \frac{t^2}{2} & \text{if } 1 < t < \frac{\pi}{2} \\ \cos t & \text{if } t > \frac{\pi}{2} \end{cases}$$

Solution: Transform $f(t)$ in terms of unit-step functions as

$$f(t) = \underbrace{2(u(t) - u(t-1))}_{\text{First piece}} + \underbrace{\frac{t^2}{2} \left(u(t-1) - u\left(t - \frac{\pi}{2}\right) \right)}_{\text{Second piece}} + \underbrace{(\cos t) u\left(t - \frac{\pi}{2}\right)}_{\text{Third piece}}$$

Question: Find LT of the piecewise function

$$f(t) = \begin{cases} 2 & \text{if } 0 < t < 1 \\ \frac{t^2}{2} & \text{if } 1 < t < \frac{\pi}{2} \\ \cos t & \text{if } t > \frac{\pi}{2} \end{cases}$$

Solution: Transform $f(t)$ in terms of unit-step functions as

$$f(t) = \underbrace{2(u(t) - u(t-1))}_{\text{First piece}} + \underbrace{\frac{t^2}{2} \left(u(t-1) - u\left(t - \frac{\pi}{2}\right) \right)}_{\text{Second piece}} + \underbrace{(\cos t) u\left(t - \frac{\pi}{2}\right)}_{\text{Third piece}}$$

$$f(t) = 2u(t) - 2u(t-1) + \frac{1}{2}t^2 u(t-1) - \frac{1}{2}t^2 u\left(t - \frac{\pi}{2}\right) + \cos(t) u\left(t - \frac{\pi}{2}\right)$$

Question: Find LT of the piecewise function

$$f(t) = \begin{cases} 2 & \text{if } 0 < t < 1 \\ \frac{t^2}{2} & \text{if } 1 < t < \frac{\pi}{2} \\ \cos t & \text{if } t > \frac{\pi}{2} \end{cases}$$

Solution: Transform $f(t)$ in terms of unit-step functions as

$$f(t) = \underbrace{2(u(t) - u(t-1))}_{\text{First piece}} + \underbrace{\frac{t^2}{2} \left(u(t-1) - u\left(t - \frac{\pi}{2}\right) \right)}_{\text{Second piece}} + \underbrace{(\cos t) u\left(t - \frac{\pi}{2}\right)}_{\text{Third piece}}$$

$$f(t) = 2u(t) - 2u(t-1) + \frac{1}{2}t^2 u(t-1) - \frac{1}{2}t^2 u\left(t - \frac{\pi}{2}\right) + \cos(t) u\left(t - \frac{\pi}{2}\right)$$

Applying LT on both sides

$$L(f(t)) = 2L(u(t)) - 2L(u(t-1)) + \frac{1}{2}L(t^2 u(t-1)) - \frac{1}{2}L(t^2 u\left(t - \frac{\pi}{2}\right)) + L(\cos(t) u\left(t - \frac{\pi}{2}\right))$$

Question: Find LT of the piecewise function

$$f(t) = \begin{cases} 2 & \text{if } 0 < t < 1 \\ \frac{t^2}{2} & \text{if } 1 < t < \frac{\pi}{2} \\ \cos t & \text{if } t > \frac{\pi}{2} \end{cases}$$

Solution: Transform $f(t)$ in terms of unit-step functions as

$$f(t) = \underbrace{2(u(t) - u(t-1))}_{\text{First piece}} + \underbrace{\frac{t^2}{2}(u(t-1) - u(t - \frac{\pi}{2}))}_{\text{Second piece}} + \underbrace{(\cos t)u(t - \frac{\pi}{2})}_{\text{Third piece}}$$

$$f(t) = 2u(t) - 2u(t-1) + \frac{1}{2}t^2u(t-1) - \frac{1}{2}t^2u(t - \frac{\pi}{2}) + \cos(t)u(t - \frac{\pi}{2})$$

Applying LT on both sides

$$L(f(t)) = 2L(u(t)) - 2L(u(t-1)) + \frac{1}{2}L(t^2u(t-1)) - \frac{1}{2}L(t^2u(t - \frac{\pi}{2})) + L(\cos(t)u(t - \frac{\pi}{2}))$$

Writing each term in $f(t)$ in the form $f(t-a)$, so that the LT of the form $f(t-a)u(t-a)$, of t-shifting theorem, may be applied.

Thus for $\frac{1}{2}t^2 u(t-1)$, $\frac{1}{2}t^2 = \frac{1}{2} \left((t-1)^2 + 2t - 1 \right)$

Thus for $\frac{1}{2}t^2 u(t-1)$, $\frac{1}{2}t^2 = \frac{1}{2} \left((t-1)^2 + 2t-1 \right) =$
 $\frac{1}{2} (t-1)^2 + (t-1) + 1 - \frac{1}{2} = \frac{1}{2} (t-1)^2 + (t-1) + \frac{1}{2}$

Thus for $\frac{1}{2}t^2 u(t-1)$, $\frac{1}{2}t^2 = \frac{1}{2} \left((t-1)^2 + 2t-1 \right) =$
 $\frac{1}{2} (t-1)^2 + (t-1) + 1 - \frac{1}{2} = \frac{1}{2} (t-1)^2 + (t-1) + \frac{1}{2}$

for $\frac{1}{2}t^2 u\left(t - \frac{\pi}{2}\right)$, $\frac{1}{2}t^2 = \frac{1}{2} \left(t - \frac{\pi}{2} \right)^2 + \frac{1}{2}\pi \left(t - \frac{\pi}{2} \right) + \frac{1}{8}\pi^2$

Thus for $\frac{1}{2}t^2 u(t-1)$, $\frac{1}{2}t^2 = \frac{1}{2} \left((t-1)^2 + 2t-1 \right) =$
 $\frac{1}{2} (t-1)^2 + (t-1) + 1 - \frac{1}{2} = \frac{1}{2} (t-1)^2 + (t-1) + \frac{1}{2}$

for $\frac{1}{2}t^2 u(t - \frac{\pi}{2})$, $\frac{1}{2}t^2 = \frac{1}{2} (t - \frac{\pi}{2})^2 + \frac{1}{2}\pi (t - \frac{\pi}{2}) + \frac{1}{8}\pi^2$

and for $\cos(t) u(t - \frac{\pi}{2})$, $\cos(t) = -\sin(t - \frac{1}{2}\pi)$

Thus for $\frac{1}{2}t^2 u(t-1)$, $\frac{1}{2}t^2 = \frac{1}{2}((t-1)^2 + 2t-1) =$
 $\frac{1}{2}(t-1)^2 + (t-1) + \frac{1}{2}$

for $\frac{1}{2}t^2 u(t - \frac{\pi}{2})$, $\frac{1}{2}t^2 = \frac{1}{2}(t - \frac{\pi}{2})^2 + \frac{1}{2}\pi(t - \frac{\pi}{2}) + \frac{1}{8}\pi^2$

and for $\cos(t) u(t - \frac{\pi}{2})$, $\cos(t) = -\sin(t - \frac{1}{2}\pi)$

$$L(f(t)) = \frac{2}{s} - \frac{2}{s}e^{-s} + \frac{1}{2}L(t^2)e^{-s} - \frac{1}{2}L(t^2)e^{-\frac{\pi}{2}s} + L(\cos(t))e^{-\frac{\pi}{2}s}$$

Thus for $\frac{1}{2}t^2 u(t-1)$, $\frac{1}{2}t^2 = \frac{1}{2}((t-1)^2 + 2t-1) =$
 $\frac{1}{2}(t-1)^2 + (t-1) + 1 - \frac{1}{2} = \frac{1}{2}(t-1)^2 + (t-1) + \frac{1}{2}$

for $\frac{1}{2}t^2 u(t - \frac{\pi}{2})$, $\frac{1}{2}t^2 = \frac{1}{2}(t - \frac{\pi}{2})^2 + \frac{1}{2}\pi(t - \frac{\pi}{2}) + \frac{1}{8}\pi^2$

and for $\cos(t) u(t - \frac{\pi}{2})$, $\cos(t) = -\sin(t - \frac{1}{2}\pi)$

$$L(f(t)) = \frac{2}{s} - \frac{2}{s}e^{-s} + \frac{1}{2}L(t^2)e^{-s} - \frac{1}{2}L(t^2)e^{-\frac{\pi}{2}s} + L(\cos(t))e^{-\frac{\pi}{2}s}$$

$$L(f(t)) = \frac{2}{s} - \frac{2}{s}e^{-s} + \frac{1}{2}L\left[\frac{1}{2}(t-1)^2 + (t-1) + \frac{1}{2}\right]e^{-s} -$$

$$\frac{1}{2}L\left[\frac{1}{2}(t - \frac{\pi}{2})^2 + \frac{1}{2}\pi(t - \frac{\pi}{2}) + \frac{1}{8}\pi^2\right]e^{-\frac{\pi}{2}s} +$$

$$L[-\sin(t - \frac{1}{2}\pi)]e^{-\frac{\pi}{2}s}$$

Thus for $\frac{1}{2}t^2 u(t-1)$, $\frac{1}{2}t^2 = \frac{1}{2}((t-1)^2 + 2t-1) =$
 $\frac{1}{2}(t-1)^2 + (t-1) + 1 - \frac{1}{2} = \frac{1}{2}(t-1)^2 + (t-1) + \frac{1}{2}$

for $\frac{1}{2}t^2 u(t - \frac{\pi}{2})$, $\frac{1}{2}t^2 = \frac{1}{2}(t - \frac{\pi}{2})^2 + \frac{1}{2}\pi(t - \frac{\pi}{2}) + \frac{1}{8}\pi^2$

and for $\cos(t) u(t - \frac{\pi}{2})$, $\cos(t) = -\sin(t - \frac{1}{2}\pi)$

$$L(f(t)) = \frac{2}{s} - \frac{2}{s}e^{-s} + \frac{1}{2}L(t^2)e^{-s} - \frac{1}{2}L(t^2)e^{-\frac{\pi}{2}s} + L(\cos(t))e^{-\frac{\pi}{2}s}$$

$$L(f(t)) = \frac{2}{s} - \frac{2}{s}e^{-s} + \frac{1}{2}L\left[\frac{1}{2}(t-1)^2 + (t-1) + \frac{1}{2}\right]e^{-s} -$$

$$\frac{1}{2}L\left[\frac{1}{2}(t - \frac{\pi}{2})^2 + \frac{1}{2}\pi(t - \frac{\pi}{2}) + \frac{1}{8}\pi^2\right]e^{-\frac{\pi}{2}s} +$$

$$L[-\sin(t - \frac{1}{2}\pi)]e^{-\frac{\pi}{2}s}$$

$$L(f(t)) = \frac{2}{s} - \frac{2}{s}e^{-s} + \frac{1}{2} \frac{(s^2+2s+2)}{s^3} e^{-s} -$$

$$\frac{1}{8} \left(\frac{\pi^2}{s} + \frac{4\pi}{s^2} + \frac{8}{s^3} \right) e^{-\frac{\pi}{2}s} - \frac{1}{s^2+1} e^{-\frac{\pi}{2}s} \quad \blacksquare$$

Question: (10ed-6.3-10)

$$f(t) = \begin{cases} \sinh t & \text{if } 0 < t < 2 \\ 0 & \text{if } t > 2 \end{cases}$$

Question: (10ed-6.3-10)

$$f(t) = \begin{cases} \sinh t & \text{if } 0 < t < 2 \\ 0 & \text{if } t > 2 \end{cases}$$

$$f(t) = \sinh(t) u(t) - \sinh(t) u(t-2)$$

Question: (10ed-6.3-10)

$$f(t) = \begin{cases} \sinh t & \text{if } 0 < t < 2 \\ 0 & \text{if } t > 2 \end{cases}$$

$$f(t) = \sinh(t) u(t) - \sinh(t) u(t-2)$$

$$f(t) = \sinh(t) u(t) - \sinh((t-2) + 2) u(t-2)$$

$$f(t) = \sinh(t) u(t) - (\cosh 2 \sinh t - \sinh 2 \cosh t) u(t-2)$$

Question: (10ed-6.3-10)

$$f(t) = \begin{cases} \sinh t & \text{if } 0 < t < 2 \\ 0 & \text{if } t > 2 \end{cases}$$

$$f(t) = \sinh(t) u(t) - \sinh(t) u(t-2)$$

$$f(t) = \sinh(t) u(t) - \sinh((t-2) + 2) u(t-2)$$

$$f(t) = \sinh(t) u(t) - (\cosh 2 \sinh t - \sinh 2 \cosh t) u(t-2)$$

$$L(f(t)) = L(\sinh(t) u(t)) - \cosh(2) L(\sinh(t) u(t-2)) + \sinh(2) L(\cosh(t) u(t-2))$$

$$= \frac{1}{s^2-1} - \cosh(2) \left(\frac{1}{s^2-1} \right) e^{-2s} + \sinh(2) \left(\frac{s}{s^2-1} \right) e^{-2s}$$

$$= \frac{1}{s^2-1} - \left(\frac{\sinh(2)s + \cosh(2)}{s^2-1} \right) e^{-2s} \quad \blacksquare$$

Question: (10ed-6.3-14)

$$\begin{aligned} L^{-1} \left(\frac{4(e^{-2s} - 2e^{-5s})}{s} \right) &= L^{-1} \left(\frac{1}{s} (4e^{-2s} - 8e^{-5s}) \right) \\ &= 4L^{-1} \left(\frac{1}{s} e^{-2s} \right) - 8L^{-1} \left(\frac{1}{s} e^{-5s} \right) \end{aligned}$$

Question: (10ed-6.3-14)

$$\begin{aligned} L^{-1} \left(\frac{4(e^{-2s} - 2e^{-5s})}{s} \right) &= L^{-1} \left(\frac{1}{s} (4e^{-2s} - 8e^{-5s}) \right) \\ &= 4L^{-1} \left(\frac{1}{s} e^{-2s} \right) - 8L^{-1} \left(\frac{1}{s} e^{-5s} \right) \\ f(t) &= 4u(t-2) - 8u(t-5) \quad \blacksquare \end{aligned}$$

Question: (10ed-6.3-14)

$$\begin{aligned}
 L^{-1} \left(\frac{4(e^{-2s} - 2e^{-5s})}{s} \right) &= L^{-1} \left(\frac{1}{s} (4e^{-2s} - 8e^{-5s}) \right) \\
 &= 4L^{-1} \left(\frac{1}{s} e^{-2s} \right) - 8L^{-1} \left(\frac{1}{s} e^{-5s} \right) \\
 f(t) &= 4u(t-2) - 8u(t-5) \quad \blacksquare
 \end{aligned}$$

Though not required, but this would be beneficial if student transforms above $f(t)$ into a piecewise representation.

Question: (10ed-6.3-14)

$$\begin{aligned}
 L^{-1} \left(\frac{4(e^{-2s} - 2e^{-5s})}{s} \right) &= L^{-1} \left(\frac{1}{s} (4e^{-2s} - 8e^{-5s}) \right) \\
 &= 4L^{-1} \left(\frac{1}{s} e^{-2s} \right) - 8L^{-1} \left(\frac{1}{s} e^{-5s} \right) \\
 f(t) &= 4u(t-2) - 8u(t-5) \quad \blacksquare
 \end{aligned}$$

Though not required, but this would be beneficial if student transforms above $f(t)$ into a piecewise representation. For this $f(t)$ it is given as

$$f(t) = 4u(t-2) - 8u(t-5) = \begin{cases} 4 & \text{if } 2 < t < 5 \\ -4 & \text{if } t > 5 \end{cases}$$

Question: (DIY, hints are given) Writing the Heaviside form of

$$f(t) = \begin{cases} \frac{1}{2}t & \text{if } 0 < t < \frac{1}{2} \\ 3t - 2 & \text{if } \frac{1}{2} < t < \frac{\pi}{2} \\ e^t & \text{if } t > \frac{\pi}{2} \end{cases}$$

gives

Question: (DIY, hints are given) Writing the Heaviside form of

$$f(t) = \begin{cases} \frac{1}{2}t & \text{if } 0 < t < \frac{1}{2} \\ 3t - 2 & \text{if } \frac{1}{2} < t < \frac{\pi}{2} \\ e^t & \text{if } t > \frac{\pi}{2} \end{cases}$$

gives

$$f(t) = \underbrace{\frac{1}{2}t u(t) + \frac{5}{2}t u\left(t - \frac{1}{2}\right)}_{\text{First piece}}$$

Question: (DIY, hints are given) Writing the Heaviside form of

$$f(t) = \begin{cases} \frac{1}{2}t & \text{if } 0 < t < \frac{1}{2} \\ 3t - 2 & \text{if } \frac{1}{2} < t < \frac{\pi}{2} \\ e^t & \text{if } t > \frac{\pi}{2} \end{cases}$$

gives

$$f(t) = \underbrace{\frac{1}{2}t u(t) + \frac{5}{2}t u\left(t - \frac{1}{2}\right)}_{\text{First piece}} -$$

$$\underbrace{3t u\left(t - \frac{\pi}{2}\right) + 2u\left(t - \frac{\pi}{2}\right) - 2u\left(t - \frac{1}{2}\right)}_{\text{Second piece}}$$

Question: (DIY, hints are given) Writing the Heaviside form of

$$f(t) = \begin{cases} \frac{1}{2}t & \text{if } 0 < t < \frac{1}{2} \\ 3t - 2 & \text{if } \frac{1}{2} < t < \frac{\pi}{2} \\ e^t & \text{if } t > \frac{\pi}{2} \end{cases}$$

gives

$$f(t) = \underbrace{\frac{1}{2}t u(t) + \frac{5}{2}t u\left(t - \frac{1}{2}\right)}_{\text{First piece}} -$$

$$\underbrace{3t u\left(t - \frac{\pi}{2}\right) + 2u\left(t - \frac{\pi}{2}\right) - 2u\left(t - \frac{1}{2}\right)}_{\text{Second piece}} + \underbrace{e^t u\left(t - \frac{\pi}{2}\right)}_{\text{Third piece}}$$

Question: (DIY, hints are given) Writing the Heaviside form of

$$f(t) = \begin{cases} \frac{1}{2}t & \text{if } 0 < t < \frac{1}{2} \\ 3t - 2 & \text{if } \frac{1}{2} < t < \frac{\pi}{2} \\ e^t & \text{if } t > \frac{\pi}{2} \end{cases}$$

gives


$$f(t) = \underbrace{\frac{1}{2}t u(t) + \frac{5}{2}t u\left(t - \frac{1}{2}\right)}_{\text{First piece}} -$$

$$\underbrace{3t u\left(t - \frac{\pi}{2}\right) + 2 u\left(t - \frac{\pi}{2}\right) - 2 u\left(t - \frac{1}{2}\right)}_{\text{Second piece}} + \underbrace{e^t u\left(t - \frac{\pi}{2}\right)}_{\text{Third piece}}$$

Applying t-shifting theorem to obtain the LT as

$$L(f(t)) = \frac{1}{2} \frac{1}{s^2} + \frac{5}{4} \frac{(s+2)}{s^2} e^{-\frac{1}{2}s} - \frac{3}{2} \left(\frac{2}{s^2} + \frac{\pi}{s} \right) e^{-\frac{\pi}{2}s} + \frac{2}{s} e^{-\frac{\pi}{2}s} - \frac{2}{s} e^{-\frac{1}{2}s} + \frac{e^{-\frac{\pi}{2}(s-1)}}{s-1}$$


Question: Find the response (the current) of the RLC-circuit given in figure, where $E(t)$ is sinusoidal,



acting for a short time interval only, say

$E(t) = 100 \sin(400t)$ if $0 < t < 2\pi$ and $E(t) = 0$ if $t > 2\pi$.

Question: Find the response (the current) of the RLC-circuit given in figure, where $E(t)$ is sinusoidal,



acting for a short time interval only, say

$E(t) = 100 \sin(400t)$ if $0 < t < 2\pi$ and $E(t) = 0$ if $t > 2\pi$.


Solution: The forcing function

$E(t) = \begin{cases} 100 \sin(400t) & \text{if } 0 < t < 2\pi \\ 0 & \text{if } t > 2\pi \end{cases}$ may be written in

terms of unit-step function as

$E(t) =$

Question: Find the response (the current) of the RLC-circuit given in figure, where $E(t)$ is sinusoidal,



acting for a short time interval only, say

$E(t) = 100 \sin(400t)$ if $0 < t < 2\pi$ and $E(t) = 0$ if $t > 2\pi$.


Solution: The forcing function

$E(t) = \begin{cases} 100 \sin(400t) & \text{if } 0 < t < 2\pi \\ 0 & \text{if } t > 2\pi \end{cases}$ may be written in

terms of unit-step function as

$E(t) = -100 \sin(400t) u(t - 2\pi) + 100 \sin(400t) u(t).$

Question: Find the response (the current) of the RLC-circuit given in figure, where $E(t)$ is sinusoidal,



acting for a short time interval only, say

$E(t) = 100 \sin(400t)$ if $0 < t < 2\pi$ and $E(t) = 0$ if $t > 2\pi$.

Solution: The forcing function

$E(t) = \begin{cases} 100 \sin(400t) & \text{if } 0 < t < 2\pi \\ 0 & \text{if } t > 2\pi \end{cases}$ may be written in

terms of unit-step function as

$E(t) = -100 \sin(400t) u(t - 2\pi) + 100 \sin(400t) u(t)$.

Model for current $i(t)$ is the integro-differential equation

$$0.1 \frac{d}{dt} i(t) + 11 i(t) + 100 \int_0^t i(\tau) d\tau = -100 \sin(400t) u(t - 2\pi) + 100 \sin(400t) u(t) \quad ; \quad i(0) = i'(0) = 0$$

$$0.1sY - 0.1i(0) + 11Y + 100\frac{1}{s}Y = \frac{40000}{(s^2+160000)} (1 - e^{-2\pi s})$$

$$0.1sY - 0.1i(0) + 11Y + 100\frac{1}{s}Y = \frac{40000}{(s^2+160000)} (1 - e^{-2\pi s})$$

$$Y = -\frac{(-i(0)s^2+400000 e^{-2\pi s}-160000 i(0)-400000)s}{(s^2+160000)(s^2+110 s+1000)}$$

$$Y = -\frac{(-400000+400000 e^{-2\pi s})s}{(s^2+160000)(s^2+110 s+1000)} = \frac{400000 s-400000 s e^{-2\pi s}}{(s^2+160000)(s^2+110 s+1000)}$$

$$0.1sY - 0.1i(0) + 11Y + 100\frac{1}{s}Y = \frac{40000}{(s^2+160000)} (1 - e^{-2\pi s})$$

$$Y = -\frac{(-i(0)s^2+400000 e^{-2\pi s}-160000 i(0)-400000)s}{(s^2+160000)(s^2+110s+1000)}$$

$$Y = -\frac{(-400000+400000 e^{-2\pi s})s}{(s^2+160000)(s^2+110s+1000)} = \frac{400000 s-400000 se^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)}$$

$$Y = \frac{400000 s}{(s^2+160000)(s^2+110s+1000)} - \frac{400000 se^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)}$$

$$0.1sY - 0.1i(0) + 11Y + 100\frac{1}{s}Y = \frac{40000}{(s^2+160000)} (1 - e^{-2\pi s})$$

$$Y = -\frac{(-i(0)s^2+400000 e^{-2\pi s}-160000 i(0)-400000)s}{(s^2+160000)(s^2+110s+1000)}$$

$$Y = -\frac{(-400000+400000 e^{-2\pi s})s}{(s^2+160000)(s^2+110s+1000)} = \frac{400000 s-400000 se^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)}$$

$$Y = \frac{400000 s}{(s^2+160000)(s^2+110s+1000)} - \frac{400000 se^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)}$$

By partial fractions of first term, $Y =$

$$\left(\frac{400}{153(s+100)} - \frac{4000}{14409(s+10)} - \frac{\frac{63600}{27217}s - \frac{7040000}{27217}}{s^2+160000} \right) - \left(\frac{400000 s e^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)} \right)$$

$$0.1sY - 0.1i(0) + 11Y + 100\frac{1}{s}Y = \frac{40000}{(s^2+160000)} (1 - e^{-2\pi s})$$

$$Y = -\frac{(-i(0)s^2+400000 e^{-2\pi s}-160000 i(0)-400000)s}{(s^2+160000)(s^2+110s+1000)}$$

$$Y = -\frac{(-400000+400000 e^{-2\pi s})s}{(s^2+160000)(s^2+110s+1000)} = \frac{400000 s-400000 se^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)}$$

$$Y = \frac{400000 s}{(s^2+160000)(s^2+110s+1000)} - \frac{400000 se^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)}$$

By partial fractions of first term, $Y =$

$$\left(\frac{400}{153(s+100)} - \frac{4000}{14409(s+10)} - \frac{\frac{63600}{27217}s - \frac{7040000}{27217}}{s^2+160000} \right) - \left(\frac{400000 s e^{-2\pi s}}{(s^2+160000)(s^2+110s+1000)} \right)$$

taking Inverse LT

$$i(t) = \left(\frac{400}{153} e^{-100t} - \frac{4000}{14409} e^{-10t} - \left(\frac{63600}{27217} \cos 400t - \frac{17600}{27217} \sin 400t \right) \right) \\ - u(t - 2\pi) \\ \left(\frac{400}{153} e^{200\pi-100t} - \frac{4000}{14409} e^{20\pi-10t} - \frac{63600}{27217} \cos 400t + \frac{17600}{27217} \sin 400t \right) \quad \blacksquare$$

Question: (10ed-6.3-23)

$$\frac{d^2y}{dt^2} + \frac{dy}{dt} - 2y = f(t) =$$

$$\begin{cases} 3 \sin(t) - \cos(t) & \text{if } 0 < t < 2\pi \\ 3 \sin(2t) - \cos(2t) & \text{if } t > 2\pi \end{cases}$$

where

$$y(0) = 1, y'(0) = 0$$

Solution:

Question: (10ed-6.3-23)

$$\frac{d^2y}{dt^2} + \frac{dy}{dt} - 2y = f(t) = \begin{cases} 3\sin(t) - \cos(t) & \text{if } 0 < t < 2\pi \\ 3\sin(2t) - \cos(2t) & \text{if } t > 2\pi \end{cases} \quad \text{where}$$

$$y(0) = 1, y'(0) = 0$$

Solution: Transforming the forcing function into Heaviside form:

Question: (10ed-6.3-23)

$$\frac{d^2y}{dt^2} + \frac{dy}{dt} - 2y = f(t) = \begin{cases} 3 \sin(t) - \cos(t) & \text{if } 0 < t < 2\pi \\ 3 \sin(2t) - \cos(2t) & \text{if } t > 2\pi \end{cases} \quad \text{where}$$

$$y(0) = 1, \quad y'(0) = 0$$

Solution: Transforming the forcing function into Heaviside form:

$$f(t) = -3 \sin(t) u(t - 2\pi) + 3 \sin(t) u(t) + \cos(t) u(t - 2\pi)$$

Question: (10ed-6.3-23)

$$\frac{d^2y}{dt^2} + \frac{dy}{dt} - 2y = f(t) = \begin{cases} 3\sin(t) - \cos(t) & \text{if } 0 < t < 2\pi \\ 3\sin(2t) - \cos(2t) & \text{if } t > 2\pi \end{cases} \quad \text{where}$$

$$y(0) = 1, \quad y'(0) = 0$$

Solution: Transforming the forcing function into Heaviside form:

$$f(t) = -3\sin(t)u(t-2\pi) + 3\sin(t)u(t) + \cos(t)u(t-2\pi) - \cos(t)u(t) + 3u(t-2\pi)\sin(2t) - u(t-2\pi)\cos(2t)$$

Question: (10ed-6.3-23)

$$\frac{d^2y}{dt^2} + \frac{dy}{dt} - 2y = f(t) = \begin{cases} 3 \sin(t) - \cos(t) & \text{if } 0 < t < 2\pi \\ 3 \sin(2t) - \cos(2t) & \text{if } t > 2\pi \end{cases} \quad \text{where}$$

$$y(0) = 1, y'(0) = 0$$

Solution: Transforming the forcing function into Heaviside form:

$$f(t) = -3 \sin(t) u(t - 2\pi) + 3 \sin(t) u(t) + \cos(t) u(t - 2\pi) - \cos(t) u(t) + 3u(t - 2\pi) \sin(2t) - u(t - 2\pi) \cos(2t)$$

Taking LT on both sides of the given DE

$$s^2 Y - y'(0) - sy(0) + sY - y(0) - 2Y =$$

$$\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right)$$

Question: (10ed-6.3-23)

$$\frac{d^2y}{dt^2} + \frac{dy}{dt} - 2y = f(t) = \begin{cases} 3\sin(t) - \cos(t) & \text{if } 0 < t < 2\pi \\ 3\sin(2t) - \cos(2t) & \text{if } t > 2\pi \end{cases} \quad \text{where}$$

$$y(0) = 1, y'(0) = 0$$

Solution: Transforming the forcing function into Heaviside form:

$$f(t) = -3\sin(t)u(t-2\pi) + 3\sin(t)u(t) + \cos(t)u(t-2\pi) - \cos(t)u(t) + 3u(t-2\pi)\sin(2t) - u(t-2\pi)\cos(2t)$$

Taking LT on both sides of the given DE

$$s^2Y - y'(0) - sy(0) + sY - y(0) - 2Y =$$

$$\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right)$$

$$Y = \frac{1}{s^2+s-2} \left(\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right) + y'(0) + sy(0) + y(0) \right)$$

$$Y = \frac{1}{s^2+s-2} \left(\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right) + 1 + s \right)$$

$$Y = \frac{s^4 - s^3 + 3e^{-2s\pi}s + 6s^2 - 3e^{-2s\pi} - 4s + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$Y = \frac{1}{s^2+s-2} \left(\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right) + 1 + s \right)$$

$$Y = \frac{s^4 - s^3 + 3e^{-2s\pi}s + 6s^2 - 3e^{-2s\pi} - 4s + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$L(y(t)) = Y = \frac{s^4 - s^3 + 6s^2 + (3e^{-2s\pi} - 4)s - 3e^{-2s\pi} + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$Y = \frac{1}{s^2+s-2} \left(\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right) + 1 + s \right)$$

$$Y = \frac{s^4 - s^3 + 3e^{-2s\pi}s + 6s^2 - 3e^{-2s\pi} - 4s + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$L(y(t)) = Y = \frac{s^4 - s^3 + 6s^2 + (3e^{-2s\pi} - 4)s - 3e^{-2s\pi} + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$y(t) = e^t - \sin(t) + \frac{1}{2} (2 \sin(t) - \sin(2t)) u(t - 2\pi)$$

$$Y = \frac{1}{s^2+s-2} \left(\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right) + 1 + s \right)$$

$$Y = \frac{s^4 - s^3 + 3e^{-2s\pi}s + 6s^2 - 3e^{-2s\pi} - 4s + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$L(y(t)) = Y = \frac{s^4 - s^3 + 6s^2 + (3e^{-2s\pi} - 4)s - 3e^{-2s\pi} + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$y(t) = e^t - \sin(t) + \frac{1}{2} (2 \sin(t) - \sin(2t)) u(t - 2\pi)$$

$$y(t) = \left\{ \right.$$

$$Y = \frac{1}{s^2+s-2} \left(\frac{1}{s^2+1} \left(3 - s + 3 \frac{e^{-2s\pi}(s+2)(s-1)}{s^2+4} \right) + 1 + s \right)$$

$$Y = \frac{s^4 - s^3 + 3e^{-2s\pi}s + 6s^2 - 3e^{-2s\pi} - 4s + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$L(y(t)) = Y = \frac{s^4 - s^3 + 6s^2 + (3e^{-2s\pi} - 4)s - 3e^{-2s\pi} + 8}{(s-1)(s^2+1)(s^2+4)}$$

$$y(t) = e^t - \sin(t) + \frac{1}{2} (2 \sin(t) - \sin(2t)) u(t - 2\pi)$$

$$y(t) = \begin{cases} e^t - \sin(t) & \text{if } 0 < t < 2\pi \\ e^t - \frac{1}{2} \sin(2t) & \text{if } t > 2\pi \end{cases} \quad \blacksquare$$

Question: (10e-6.3-27) Put $\tilde{t} = t - 1$ in

$$E(t) = \begin{cases} 8t^2 & 0 < t < 5 \\ 0 & t > 5 \end{cases} \quad \text{and writing it in Heaviside terms:}$$

Question: (10e-6.3-27) Put $\tilde{t} = t - 1$ in

$$E(t) = \begin{cases} 8t^2 & 0 < t < 5 \\ 0 & t > 5 \end{cases} \quad \text{and writing it in Heaviside terms:}$$

$$E(\tilde{t}) = \begin{cases} 8(\tilde{t} + 1)^2 & -1 < \tilde{t} < 4 \\ 0 & \tilde{t} > 4 \end{cases}$$

Question: (10e-6.3-27) Put $\tilde{t} = t - 1$ in

$$E(t) = \begin{cases} 8t^2 & 0 < t < 5 \\ 0 & t > 5 \end{cases} \quad \text{and writing it in Heaviside terms:}$$

$$E(\tilde{t}) = \begin{cases} 8(\tilde{t} + 1)^2 & -1 < \tilde{t} < 4 \\ 0 & \tilde{t} > 4 \end{cases}$$

$$E(\tilde{t}) = 8u(\tilde{t} + 1)\tilde{t}^2 - 8u(\tilde{t} - 4)\tilde{t}^2 + 16u(\tilde{t} + 1)\tilde{t} - 16u(\tilde{t} - 4)\tilde{t} + 8u(\tilde{t} + 1) - 8u(\tilde{t} - 4)$$

Question: (10e-6.3-27) Put $\tilde{t} = t - 1$ in

$$E(t) = \begin{cases} 8t^2 & 0 < t < 5 \\ 0 & t > 5 \end{cases} \quad \text{and writing it in Heaviside terms:}$$

$$E(\tilde{t}) = \begin{cases} 8(\tilde{t} + 1)^2 & -1 < \tilde{t} < 4 \\ 0 & \tilde{t} > 4 \end{cases}$$

$$E(\tilde{t}) = 8u(\tilde{t} + 1)\tilde{t}^2 - 8u(\tilde{t} - 4)\tilde{t}^2 + 16u(\tilde{t} + 1)\tilde{t} - 16u(\tilde{t} - 4)\tilde{t} + 8u(\tilde{t} + 1) - 8u(\tilde{t} - 4)$$

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + 4y(\tilde{t}) = E(\tilde{t}) \quad ; \quad y(0) = 1 + \cos(2), \quad y'(0) = 4 - 2\sin(2)$$

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + 4y(\tilde{t}) = 8u(\tilde{t} + 1)\tilde{t}^2 - 8u(\tilde{t} - 4)\tilde{t}^2 + 16u(\tilde{t} + 1)\tilde{t} - 16u(\tilde{t} - 4)\tilde{t} + 8u(\tilde{t} + 1) - 8u(\tilde{t} - 4)$$

Question: (10e-6.3-27) Put $\tilde{t} = t - 1$ in

$$E(t) = \begin{cases} 8t^2 & 0 < t < 5 \\ 0 & t > 5 \end{cases} \quad \text{and writing it in Heaviside terms:}$$

$$E(\tilde{t}) = \begin{cases} 8(\tilde{t}+1)^2 & -1 < \tilde{t} < 4 \\ 0 & \tilde{t} > 4 \end{cases}$$

$$E(\tilde{t}) = 8u(\tilde{t}+1)\tilde{t}^2 - 8u(\tilde{t}-4)\tilde{t}^2 + 16u(\tilde{t}+1)\tilde{t} - 16u(\tilde{t}-4)\tilde{t} + 8u(\tilde{t}+1) - 8u(\tilde{t}-4)$$

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + 4y(\tilde{t}) = E(\tilde{t}) \quad ; \quad y(0) = 1 + \cos(2), \quad y'(0) = 4 - 2\sin(2)$$

$$\frac{d^2}{d\tilde{t}^2}y(\tilde{t}) + 4y(\tilde{t}) = 8u(\tilde{t}+1)\tilde{t}^2 - 8u(\tilde{t}-4)\tilde{t}^2 + 16u(\tilde{t}+1)\tilde{t} - 16u(\tilde{t}-4)\tilde{t} + 8u(\tilde{t}+1) - 8u(\tilde{t}-4)$$

$$s^2\tilde{Y} - y'(0) - sy(0) + 4\tilde{Y} = 16\frac{1}{s^2} + 8\frac{1}{s} + 8\frac{-e^{-4s}(25s^2+10s+2)+2}{s^3}$$

$$s^2 \tilde{Y} - 4 + 2 \sin(2) - s(1 + \cos(2)) + 4 \tilde{Y} = 16 \frac{1}{s^2} + 8 \frac{1}{s} + 8 \frac{-e^{-4s}(25s^2 + 10s + 2) + 2}{s^3} \quad \text{using ICs}$$

$$\tilde{Y} = \frac{1}{s^2 + 4} \left(\frac{16}{s^2} + \frac{8}{s} + 8 \frac{-e^{-4s}(25s^2 + 10s + 2) + 2}{s^3} + 4 - 2 \sin(2) + s(1 + \cos(2)) \right)$$

$$s^2 \tilde{Y} - 4 + 2 \sin(2) - s(1 + \cos(2)) + 4 \tilde{Y} = 16 \frac{1}{s^2} + 8 \frac{1}{s} + 8 \frac{-e^{-4s}(25s^2 + 10s + 2) + 2}{s^3} \quad \text{using ICs}$$

$$\tilde{Y} = \frac{1}{s^2 + 4} \left(\frac{16}{s^2} + \frac{8}{s} + 8 \frac{-e^{-4s}(25s^2 + 10s + 2) + 2}{s^3} + 4 - 2 \sin(2) + s(1 + \cos(2)) \right)$$

$$\tilde{Y} = \frac{-s^4 \cos(2) + 2 \sin(2)s^3 - s^4 + 200e^{-4s}s^2 - 4s^3 + 80e^{-4s}s - 8s^2 + 16e^{-4s} - 16s - 16}{s^3(s^2 + 4)}$$

$$\tilde{Y} = \frac{-s^4 \cos(2) + 2 \sin(2)s^3 - s^4 + 200e^{-4s}s^2 - 4s^3 + 80e^{-4s}s - 8s^2 + 16e^{-4s} - 16s - 16}{s^3(s^2 + 4)}$$

$$y(\tilde{t}) = 1 + \cos(2) \cos(2\tilde{t}) - \sin(2) \sin(2\tilde{t}) + 2\tilde{t}^2 + 4\tilde{t} - u(\tilde{t} - 4) \\ (100 (\sin(\tilde{t} - 4))^2 + 2\tilde{t}^2 + \cos(2\tilde{t} - 8) - 10 \sin(2\tilde{t} - 8) + 4\tilde{t} - 49)$$

$$y(\tilde{t}) = \begin{cases} 1 + \cos(2) \cos(2\tilde{t}) - \sin(2) \sin(2\tilde{t}) + 2\tilde{t}^2 + 4\tilde{t} & \tilde{t} < 4 \\ -50 + \cos(2) \cos(2\tilde{t}) - \sin(2) \sin(2\tilde{t}) + 100 (\cos(\tilde{t} - 4))^2 - \cos(2\tilde{t} - 8) + 10 \sin(2\tilde{t} - 8) & \tilde{t} > 4 \end{cases}$$

$$y(\tilde{t}) = \begin{cases} 1 + \cos(2) \cos(2\tilde{t}) - \sin(2) \sin(2\tilde{t}) + 2\tilde{t}^2 + 4\tilde{t} & \tilde{t} < 4 \\ -50 + \cos(2) \cos(2\tilde{t}) - \sin(2) \sin(2\tilde{t}) + 100(\cos(\tilde{t}-4))^2 - \cos(2\tilde{t}-8) + 10 \sin(2\tilde{t}-8) & \tilde{t} > 4 \end{cases}$$

putting back $\tilde{t} = t - 1$, $y(t) =$

$$\begin{cases} -3 + \cos(2) \cos(2t-2) - \sin(2) \sin(2t-2) + 2(t-1)^2 + 4t & t < 5 \\ -50 + \cos(2) \cos(2t-2) - \sin(2) \sin(2t-2) + 100(\cos(t-5))^2 - \cos(2t-10) + 10 \sin(2t-10) & t > 5 \end{cases}$$

$$y(t) = \begin{cases} \cos(2t) + 2t^2 - 1 & \text{if } t < 5 \\ \cos(2t) + 49 \cos(2t-10) + 10 \sin(2t-10) & \text{if } t > 5 \end{cases}$$



Question: (10ed-6.3-30) $0.5 \frac{d}{dt} i(t) + 10i(t) = E(t) =$

$$\begin{cases} 200t & , \quad 0 < t < 2 \\ 0 & , \quad t > 2 \end{cases} ; \quad i(0) = 0$$

Question: (10ed-6.3-30) $0.5 \frac{d}{dt} i(t) + 10i(t) = E(t) =$

$$\begin{cases} 200t & , \quad 0 < t < 2 \\ 0 & , \quad t > 2 \end{cases} ; \quad i(0) = 0$$

$$E(t) = 200t \, u(t) - 200t \, u(t - 2)$$

Question: (10ed-6.3-30) $0.5 \frac{d}{dt} i(t) + 10i(t) = E(t) =$
$$\begin{cases} 200t & , \quad 0 < t < 2 \\ 0 & , \quad t > 2 \end{cases} ; \quad i(0) = 0$$

$$E(t) = 200t u(t) - 200t u(t - 2)$$

$$0.5 \frac{d}{dt} i(t) + 10i(t) = 200t u(t) - 200t u(t - 2)$$

$$0.5sY - 0.5i(0) + 10Y = 200(1 - e^{-2s}(2s + 1))s^{-2}$$

Question: (10ed-6.3-30) $0.5 \frac{d}{dt} i(t) + 10i(t) = E(t) =$
$$\begin{cases} 200t & , \quad 0 < t < 2 \\ 0 & , \quad t > 2 \end{cases} ; \quad i(0) = 0$$

$$E(t) = 200t u(t) - 200t u(t - 2)$$

$$0.5 \frac{d}{dt} i(t) + 10i(t) = 200t u(t) - 200t u(t - 2)$$

$$0.5sY - 0.5i(0) + 10Y = 200(1 - e^{-2s}(2s + 1))s^{-2}$$

$$0.5sY + 10Y = 200(1 - e^{-2s}(2s + 1))s^{-2}$$

$$Y = -400 \frac{2e^{-2s}s + e^{-2s} - 1}{s^2(s + 20)}$$

Question: (10ed-6.3-30) $0.5 \frac{d}{dt} i(t) + 10i(t) = E(t) =$

$$\begin{cases} 200t & , \quad 0 < t < 2 \\ 0 & , \quad t > 2 \end{cases} ; \quad i(0) = 0$$

$$E(t) = 200t u(t) - 200t u(t-2)$$

$$0.5 \frac{d}{dt} i(t) + 10i(t) = 200t u(t) - 200t u(t-2)$$

$$0.5sY - 0.5i(0) + 10Y = 200(1 - e^{-2s}(2s+1))s^{-2}$$

$$0.5sY + 10Y = 200(1 - e^{-2s}(2s+1))s^{-2}$$

$$Y = -400 \frac{2e^{-2s}s + e^{-2s} - 1}{s^2(s+20)}$$

$$i(t) = -1 + 20t u(2-t) + e^{-20t} + 2 u(t-2) \left(e^{(20-10t)} \sinh(10t-20) + 20 e^{(-20t+40)} \right)$$

$$i(t) = \begin{cases} -1 + e^{-20t} + 20t & , \quad t < 2 \\ -1 + e^{-20t} + 2 e^{(20-10t)} \sinh(10t-20) + 40 e^{(-20t+40)} & , \quad t > 2 \end{cases}$$

Question: (10ed-6.3-40)


$$\frac{d}{dt}i(t) + 2i(t) + 10 \int_0^t i(\tau) d\tau = E(t) =$$

$$\begin{cases} 255 \sin(t) & 0 < t < 2\pi \\ 0 & t > 2\pi \end{cases} ; \quad i(0) = 0$$

Question: (10ed-6.3-40)

$$\frac{d}{dt}i(t) + 2i(t) + 10 \int_0^t i(\tau) d\tau = E(t) =$$


$$\begin{cases} 255 \sin(t) & 0 < t < 2\pi \\ 0 & t > 2\pi \end{cases} ; \quad i(0) = 0$$



$$E(t) = 255 \sin(t) u(t) - 255 \sin(t) u(-2\pi + t)$$

Question: (10ed-6.3-40)

$$\frac{d}{dt}i(t) + 2i(t) + 10 \int_0^t i(\tau) d\tau = E(t) =$$



$$\begin{cases} 255 \sin(t) & 0 < t < 2\pi \\ 0 & t > 2\pi \end{cases} ; \quad i(0) = 0$$

$$E(t) = 255 \sin(t) u(t) - 255 \sin(t) u(-2\pi + t)$$

$$\frac{d}{dt}i(t) + 2i(t) + 10 \int_0^t i(\tau) d\tau =$$

$$255 \sin(t) u(t) - 255 \sin(t) u(-2\pi + t)$$

$$s Y - i(0) + 2Y + 10 \frac{Y}{s} = 255 \frac{1 - e^{-2s\pi}}{s^2 + 1}$$

$$Y = 1 \left(255 \frac{1 - e^{-2s\pi}}{s^2 + 1} + i(0) \right) (s + 2 + 10s^{-1})^{-1}$$

$$Y = 255 \frac{1 - e^{-2s\pi}}{(s^2 + 1) \left(s + 2 + \frac{10}{s} \right)}$$

$$Y = -255 \frac{(-1 + e^{-2s\pi})s}{(s^2 + 1)(s^2 + 2s + 10)}$$

$$Y = 255 \frac{1 - e^{-2s\pi}}{(s^2 + 1) \left(s + 2 + \frac{10}{s}\right)}$$

$$Y = -255 \frac{(-1 + e^{-2s\pi})s}{(s^2 + 1)(s^2 + 2s + 10)}$$

$$i(t) = 3 u(2\pi - t) (9 \cos(t) + 2 \sin(t)) +$$

$$(e^{2\pi - t} u(-2\pi + t) - e^{-t}) (27 \cos(3t) + 11 \sin(3t))$$

$$i(t) =$$

$$\begin{cases} 27 \cos(t) + 6 \sin(t) - (27 \cos(3t) + 11 \sin(3t)) e^{-t} & t < 2\pi \\ (27 \cos(3t) + 11 \sin(3t))(-e^{-t} + e^{(2\pi - t)}) & t > 2\pi \end{cases}$$

$$Y = 255 \frac{1 - e^{-2s\pi}}{(s^2 + 1) \left(s + 2 + \frac{10}{s}\right)}$$

$$Y = -255 \frac{(-1 + e^{-2s\pi})s}{(s^2 + 1)(s^2 + 2s + 10)}$$

$$i(t) = 3 u(2\pi - t) (9 \cos(t) + 2 \sin(t)) +$$

$$(e^{2\pi-t} u(-2\pi + t) - e^{-t}) (27 \cos(3t) + 11 \sin(3t))$$

$$i(t) =$$

$$\begin{cases} 27 \cos(t) + 6 \sin(t) - (27 \cos(3t) + 11 \sin(3t)) e^{-t} & t < 2\pi \\ (27 \cos(3t) + 11 \sin(3t))(-e^{-t} + e^{(2\pi-t)}) & t > 2\pi \end{cases}$$

On plotting the current $i(t)$ we have

