

Advanced Engineering Mathematics

Vector Differential Calculus

Dr Athar Kharal atharkharal@gmail.com

Humanities and Science Department
College of Aeronautical Engineering
National University of Sciences and Technology (NUST), Pakistan

Vector and Scalar Functions and their Fields

Definition

For any point $P(x, y, z)$, the **scalar function** f is defined as $f(P) = f(x, y, z)$.

Vector and Scalar Functions and their Fields

Definition

For any point $P(x, y, z)$, the **scalar function** f is defined as $f(P) = f(x, y, z)$. And the **vector function** \mathbf{v} is defined as $\mathbf{v} = \mathbf{v}(P) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$.

Vector and Scalar Functions and their Fields

Definition

For any point $P(x, y, z)$, the **scalar function** f is defined as $f(P) = f(x, y, z)$. And the **vector function** \mathbf{v} is defined as $\mathbf{v} = \mathbf{v}(P) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$.

Definition

Field is a region in which every point has a defined scalar or vector attached to it through a scalar or vector function.

Vector and Scalar Functions and their Fields

Definition

For any point $P(x, y, z)$, the **scalar function** f is defined as $f(P) = f(x, y, z)$. And the **vector function** \mathbf{v} is defined as $\mathbf{v} = \mathbf{v}(P) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$.

Definition

Field is a region in which every point has a defined scalar or vector attached to it through a scalar or vector function. The field is accordingly named as **scalar field** or **vector field**.

Vector and Scalar Functions and their Fields

Definition

For any point $P(x, y, z)$, the **scalar function** f is defined as $f(P) = f(x, y, z)$. And the **vector function** \mathbf{v} is defined as $\mathbf{v} = \mathbf{v}(P) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$.

Definition

Field is a region in which every point has a defined scalar or vector attached to it through a scalar or vector function. The field is accordingly named as **scalar field** or **vector field**.

Examples of scalar fields are: Temperature field of a body and pressure field of air in Earth's atmosphere.

Vector and Scalar Functions and their Fields


Definition

For any point $P(x, y, z)$, the **scalar function** f is defined as $f(P) = f(x, y, z)$. And the **vector function** \mathbf{v} is defined as $\mathbf{v} = \mathbf{v}(P) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$.

Definition

Field is a region in which every point has a defined scalar or vector attached to it through a scalar or vector function. The field is accordingly named as **scalar field** or **vector field**.

Examples of scalar fields are: Temperature field of a body and pressure field of air in Earth's atmosphere.



Examples of vector field are: gravitational field,


Examples of vector field are: gravitational field, electromagnetic field,

Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft,

Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft, field of tangent vectors of a curve


Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft, field of tangent vectors of a curve and field of normal vectors of a surface.

Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft, field of tangent vectors of a curve and field of normal vectors of a surface.




Field of tangents

Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft, field of tangent vectors of a curve and field of normal vectors of a surface.




Field of tangents




Field of normals


Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft, field of tangent vectors of a curve and field of normal vectors of a surface.




Field of tangents




Field of normals





Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft, field of tangent vectors of a curve and field of normal vectors of a surface.




Field of tangents




Field of normals






Examples of vector field are: gravitational field, electromagnetic field, flow field around an aircraft, field of tangent vectors of a curve and field of normal vectors of a surface.






Field of tangents





Field of normals









NASA's Seasat used radar to take 350,000 wind measurements over the world's oceans. In the figure, the arrows show wind direction; their length and the color contouring indicate speed:



NASA's Seasat used radar to take 350,000 wind measurements over the world's oceans. In the figure, the arrows show wind direction; their length and the color contouring indicate speed: hence a vector field!



NASA's Seasat used radar to take 350,000 wind measurements over the world's oceans. In the figure, the arrows show wind direction; their length and the color contouring indicate speed: hence a vector field! Notice the heavy storm south of Greenland.

Calculus of vector functions is called Vector Calculus.

Calculus of vector functions is called Vector Calculus. It has two main branches Vector Differential Calculus and Vector Integral Calculus.

Calculus of vector functions is called Vector Calculus. It has two main branches Vector Differential Calculus and Vector Integral Calculus. Except few adjustments in notations and scheme of representation of various mathematical objects, it is similar to its classical counterparts in Calculus of real functions.

Calculus of vector functions is called Vector Calculus. It has two main branches Vector Differential Calculus and Vector Integral Calculus. Except few adjustments in notations and scheme of representation of various mathematical objects, it is similar to its classical counterparts in Calculus of real functions.

Definition

The derivative of a vector function $\mathbf{v}(t) = [v_1(t), v_2(t), v_3(t)]$ is given as $\mathbf{v}'(t) = [v_1'(t), v_2'(t), v_3'(t)]$.

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function $T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function

$T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Solution: For isotherms put $T(x, y) = \frac{y}{x^2 + y^2} = c$, where c is a constant.

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function

$T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Solution: For isotherms put $T(x, y) = \frac{y}{x^2 + y^2} = c$, where c is a constant. In such questions we try to unearth an equation of a familiar curve. Here

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function

$T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Solution: For isotherms put $T(x, y) = \frac{y}{x^2 + y^2} = c$, where c is a constant. In such questions we try to unearth an equation of a familiar curve. Here denominator indicates something near to a circle, perhaps.

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function

$T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Solution: For isotherms put $T(x, y) = \frac{y}{x^2 + y^2} = c$, where c is a constant. In such questions we try to unearth an equation of a familiar curve. Here denominator indicates something near to a circle, perhaps. So we proceed as follows:

$$\frac{y}{x^2 + y^2} = c \Rightarrow \frac{x^2 + y^2}{y} = \frac{1}{c}$$

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function

$T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Solution: For isotherms put $T(x, y) = \frac{y}{x^2 + y^2} = c$, where c is a constant. In such questions we try to unearth an equation of a familiar curve. Here denominator indicates something near to a circle, perhaps. So we proceed as follows:

$$\frac{y}{x^2 + y^2} = c \Rightarrow \frac{x^2 + y^2}{y} = \frac{1}{c}$$

$$x^2 + y^2 = \frac{y}{c} \Rightarrow x^2 + y^2 - \frac{y}{c} = 0$$

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function

$T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Solution: For isotherms put $T(x, y) = \frac{y}{x^2 + y^2} = c$, where c is a constant. In such questions we try to unearth an equation of a familiar curve. Here denominator indicates something near to a circle, perhaps. So we proceed as follows:

$$\frac{y}{x^2 + y^2} = c \Rightarrow \frac{x^2 + y^2}{y} = \frac{1}{c}$$

$$x^2 + y^2 = \frac{y}{c} \Rightarrow x^2 + y^2 - \frac{y}{c} = 0$$

Completing square

$$x^2 + y^2 - \frac{y}{c} + \left(\frac{1}{2c}\right)^2 - \left(\frac{1}{2c}\right)^2 = 0$$

$$x^2 + \left(y - \frac{1}{2c}\right)^2 = \frac{1}{4c^2}$$

Hence the isotherms for this scalar field are circles with centre $(0, \frac{1}{2c})$ and radius $\frac{1}{2c}$.

Question: (10ed-9.4-5) Let the temperature T in a body be independent of z so that it is given by a scalar function

$T = T(x, y)$. Identify the isotherms $T(x, y) = \text{const}$ for $T(x, y) = \frac{y}{x^2 + y^2}$. Also sketch some of them.

Solution: For isotherms put $T(x, y) = \frac{y}{x^2 + y^2} = c$, where c is a constant. In such questions we try to unearth an equation of a familiar curve. Here denominator indicates something near to a circle, perhaps. So we proceed as follows:

$$\frac{y}{x^2 + y^2} = c \Rightarrow \frac{x^2 + y^2}{y} = \frac{1}{c}$$

$$x^2 + y^2 = \frac{y}{c} \Rightarrow x^2 + y^2 - \frac{y}{c} = 0$$


Completing square

$$x^2 + y^2 - \frac{y}{c} + \left(\frac{1}{2c}\right)^2 - \left(\frac{1}{2c}\right)^2 = 0$$

$$x^2 + \left(y - \frac{1}{2c}\right)^2 = \frac{1}{4c^2}$$

Hence the isotherms for this scalar field are circles with centre $(0, \frac{1}{2c})$ and radius $\frac{1}{2c}$.

Isotherms for $c = \frac{1}{2}, 3$ and 5 are sketched below:



Question: (10ed-9.4-12) What kind of surfaces are the level surfaces $f(x, y, z) = \text{const}$, if $f(x, y, z) = z - \sqrt{x^2 + y^2}$?

Question: (10ed-9.4-12) What kind of surfaces are the level surfaces $f(x, y, z) = \text{const}$, if $f(x, y, z) = z - \sqrt{x^2 + y^2}$?

Solution: For finding the level surfaces

$f(x, y, z) = z - \sqrt{x^2 + y^2} = c$ where c is a constant.

$$z - \sqrt{x^2 + y^2} = c \Rightarrow z - c = \sqrt{x^2 + y^2}$$

is a typical equation of

Question: (10ed-9.4-12) What kind of surfaces are the level surfaces $f(x, y, z) = \text{const}$, if $f(x, y, z) = z - \sqrt{x^2 + y^2}$?

Solution: For finding the level surfaces

$f(x, y, z) = z - \sqrt{x^2 + y^2} = c$ where c is a constant.

$$z - \sqrt{x^2 + y^2} = c \Rightarrow z - c = \sqrt{x^2 + y^2}$$

is a typical equation of one branch of hyperbola with origin at $(0, 0, c)$.


Question: (10ed-9.4-12) What kind of surfaces are the level surfaces $f(x, y, z) = \text{const}$, if $f(x, y, z) = z - \sqrt{x^2 + y^2}$?

Solution: For finding the level surfaces

$f(x, y, z) = z - \sqrt{x^2 + y^2} = c$ where c is a constant.

$$z - \sqrt{x^2 + y^2} = c \Rightarrow z - c = \sqrt{x^2 + y^2}$$

is a typical equation of one branch of hyperbola with origin at $(0, 0, c)$. For $c = \frac{1}{2}, 2$ and 6 we have the sketches as



Question: (10ed-9.4-15) Sketch the vector field given by $\mathbf{v} = \mathbf{i} - \mathbf{j}$.

Question: (10ed-9.4-15) Sketch the vector field given by $\mathbf{v} = \mathbf{i} - \mathbf{j}$.

Solution: Consider some arbitrary points $(1, 2)$, $(7, 4)$ and $(2, 6)$.

Then

$$\text{at point } (1, 2), \mathbf{v} = \mathbf{i} - 2\mathbf{j} \text{ and } |\mathbf{v}| = \sqrt{5} = 2.2$$

$$\text{at point } (7, 4), \mathbf{v} = 7\mathbf{i} - 4\mathbf{j} \text{ and } |\mathbf{v}| = \sqrt{65} = 8.1$$

$$\text{at point } (2, 6), \mathbf{v} = 2\mathbf{i} - 6\mathbf{j} \text{ and } |\mathbf{v}| = \sqrt{40} = 6.3$$

Question: (10ed-9.4-15) Sketch the vector field given by $\mathbf{v} = \mathbf{i} - \mathbf{j}$.

Solution: Consider some arbitrary points $(1, 2)$, $(7, 4)$ and $(2, 6)$.


Then

at point $(1, 2)$, $\mathbf{v} = \mathbf{i} - 2\mathbf{j}$ and $|\mathbf{v}| = \sqrt{5} = 2.2$

at point $(7, 4)$, $\mathbf{v} = 7\mathbf{i} - 4\mathbf{j}$ and $|\mathbf{v}| = \sqrt{65} = 8.1$

at point $(2, 6)$, $\mathbf{v} = 2\mathbf{i} - 6\mathbf{j}$ and $|\mathbf{v}| = \sqrt{40} = 6.3$

Doing the same for enough points and then on each point draw an arrow in the direction of \mathbf{v} with length $|\mathbf{v}|$ we get



Question: (10ed-9.4-24) Find the partial derivative of
 $\mathbf{v}_1 = [e^x \cos y, e^x \sin y]$ and $\mathbf{v}_2 = [\cos x \cosh y, -\sin x \sinh y]$.

Question: (10ed-9.4-24) Find the partial derivative of

$\mathbf{v}_1 = [e^x \cos y, e^x \sin y]$ and $\mathbf{v}_2 = [\cos x \cosh y, -\sin x \sinh y]$.

Solution:

$$\frac{\partial \mathbf{v}_1}{\partial x} = \left[\frac{\partial}{\partial x} (e^x \cos y), \frac{\partial}{\partial x} (e^x \sin y) \right] = [e^x \cos y, e^x \sin y]$$

$$\frac{\partial \mathbf{v}_1}{\partial y} = \left[\frac{\partial}{\partial y} (e^x \cos y), \frac{\partial}{\partial y} (e^x \sin y) \right] = [-e^x \sin y, e^x \cos y]$$

Question: (10ed-9.4-24) Find the partial derivative of

$\mathbf{v}_1 = [e^x \cos y, e^x \sin y]$ and $\mathbf{v}_2 = [\cos x \cosh y, -\sin x \sinh y]$.

Solution:

$$\frac{\partial \mathbf{v}_1}{\partial x} = \left[\frac{\partial}{\partial x} (e^x \cos y), \frac{\partial}{\partial x} (e^x \sin y) \right] = [e^x \cos y, e^x \sin y]$$

$$\frac{\partial \mathbf{v}_1}{\partial y} = \left[\frac{\partial}{\partial y} (e^x \cos y), \frac{\partial}{\partial y} (e^x \sin y) \right] = [-e^x \sin y, e^x \cos y]$$

For \mathbf{v}_2 ,

$$\frac{\partial \mathbf{v}_2}{\partial x} = \left[\frac{\partial}{\partial x} (\cos x \cosh y), \frac{\partial}{\partial x} (-\sin x \sinh y) \right]$$

$$= [-\sin x \cosh y, -\cos x \sinh y]$$



$$\frac{\partial \mathbf{v}_2}{\partial y} = \left[\frac{\partial}{\partial y} (\cos x \cosh y), \frac{\partial}{\partial y} (-\sin x \sinh y) \right]$$

$$= [\cos x \sinh y, -\sin x \cosh y] \quad \blacksquare$$

Curves and Arc Length

A parametric representation of a space curve is given as



$$\mathbf{r}(t) = [x(t), y(t), z(t)] = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}.$$



Curves and Arc Length

A parametric representation of a space curve is given as

$$\mathbf{r}(t) = [x(t), y(t), z(t)] = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}.$$





Such a representation has two distinct advantages:

Curves and Arc Length

A parametric representation of a space curve is given as

$$\mathbf{r}(t) = [x(t), y(t), z(t)] = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}.$$





Such a representation has two distinct advantages:

- 1 the coordinates x, y, z play an equal role, i.e. all three are dependent variables,

Curves and Arc Length

A parametric representation of a space curve is given as

$$\mathbf{r}(t) = [x(t), y(t), z(t)] = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}.$$




Such a representation has two distinct advantages:

- 1 the coordinates x, y, z play an equal role, i.e. all three are dependent variables,
- 2 this representation induces an orientation, i.e. a beginning and an end equivalently a sense of direction, of the curve.

Following are few examples of space curves with their parametric representations:


Following are few examples of space curves with their parametric representations:

$$\mathbf{r}(t) = [h + a \cos t, k + a \sin t, 0] = (h + a \cos(t))\mathbf{i} + (k + a \sin(t))\mathbf{j}; \text{ radius} = a, \text{ centre} = (h, k)$$




Following are few examples of space curves with their parametric representations:

$$\mathbf{r}(t) = [h + a \cos t, k + a \sin t, 0] = (h + a \cos(t))\mathbf{i} + (k + a \sin(t))\mathbf{j}; \text{ radius} = a, \text{ centre} = (h, k)$$




$$\mathbf{r}(t) = [a \cos t, b \sin t, 0] = a \cos(t)\mathbf{i} + b \sin(t)\mathbf{j}; a \neq b$$




Following are few examples of space curves with their parametric representations:


$$\mathbf{r}(t) = [h + a \cos t, k + a \sin t, 0] = (h + a \cos(t))\mathbf{i} + (k + a \sin(t))\mathbf{j}; \text{ radius} = a, \text{ centre} = (h, k)$$



$$\mathbf{r}(t) = [a \cos t, b \sin t, 0] = a \cos(t)\mathbf{i} + b \sin(t)\mathbf{j}; a \neq b$$




$$\mathbf{r}(t) = \mathbf{a} + \mathbf{b}t = [a_1 + b_1 t, a_2 + b_2 t, a_3 + b_3 t]$$




Following are few examples of space curves with their parametric representations:


$$\mathbf{r}(t) = [h + a \cos t, k + a \sin t, 0] = (h + a \cos(t))\mathbf{i} + (k + a \sin(t))\mathbf{j}; \text{ radius} = a, \text{ centre} = (h, k)$$




$$\mathbf{r}(t) = [a \cos t, b \sin t, 0] = a \cos(t)\mathbf{i} + b \sin(t)\mathbf{j}; a \neq b$$




$$\mathbf{r}(t) = \mathbf{a} + \mathbf{b}t = [a_1 + b_1t, a_2 + b_2t, a_3 + b_3t]$$



$$\mathbf{r}(t) = [a \cos t, a \sin t, ct] = a \cos(t)\mathbf{i} + a \sin(t)\mathbf{j} + ct\mathbf{k}$$





Definitions

We have:

- The tangent vector to the curve $\mathbf{r}(t)$ at point P is $\mathbf{r}'(t)$.

Definitions


We have:

- The tangent vector to the curve $\mathbf{r}(t)$ at point P is $\mathbf{r}'(t)$.
- Unit tangent is computed as $\mathbf{u} = \frac{1}{|\mathbf{r}'|} \mathbf{r}'$. Both \mathbf{r}' and \mathbf{u} are in the direction of increasing t (a blessing of parametric notation!).

Definitions

We have:


- The tangent vector to the curve $\mathbf{r}(t)$ at point P is $\mathbf{r}'(t)$.
- Unit tangent is computed as $\mathbf{u} = \frac{1}{|\mathbf{r}'|} \mathbf{r}'$. Both \mathbf{r}' and \mathbf{u} are in the direction of increasing t (a blessing of parametric notation!).
- Vector equation of the tangent line passing through point P on curve $\mathbf{r}(t)$ is given as $\mathbf{q}(w) = \mathbf{r} + w\mathbf{r}'$



Definitions

We have:

- The tangent vector to the curve $\mathbf{r}(t)$ at point P is $\mathbf{r}'(t)$.
- Unit tangent is computed as $\mathbf{u} = \frac{1}{|\mathbf{r}'|} \mathbf{r}'$. Both \mathbf{r}' and \mathbf{u} are in the direction of increasing t (a blessing of parametric notation!).
- Vector equation of the tangent line passing through point P on curve $\mathbf{r}(t)$ is given as $\mathbf{q}(w) = \mathbf{r} + w\mathbf{r}'$



- Length of the curve $\mathbf{r}(t)$ from $t = a$ to an arbitrary point in t is given as $s(t) = \int_a^t \sqrt{\mathbf{r}' \cdot \mathbf{r}'} d\tau$, where $\mathbf{r}' = \frac{d\mathbf{r}}{d\tau}$.

- If a curve is representing the path of a moving body, as usually is the case in Mechanics, then velocity and acceleration of the body are given as $\mathbf{v}(t) = \mathbf{r}'(t)$ and $\mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t)$.

Definitions

- If a curve is representing the path of a moving body, as usually is the case in Mechanics, then velocity and acceleration of the body are given as $\mathbf{v}(t) = \mathbf{r}'(t)$ and $\mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t)$.
- Acceleration vector has its tangential and normal components i.e. $\mathbf{a} = \mathbf{a}_{\text{tan}} + \mathbf{a}_{\text{norm}}$, which are obtained as $\mathbf{a}_{\text{tan}} = \frac{\mathbf{a} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}} \mathbf{v}$ and $\mathbf{a}_{\text{norm}} = \mathbf{a} - \mathbf{a}_{\text{tan}}$.

Definitions


- If a curve is representing the path of a moving body, as usually is the case in Mechanics, then velocity and acceleration of the body are given as $\mathbf{v}(t) = \mathbf{r}'(t)$ and $\mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t)$.
- Acceleration vector has its tangential and normal components i.e. $\mathbf{a} = \mathbf{a}_{\text{tan}} + \mathbf{a}_{\text{norm}}$, which are obtained as $\mathbf{a}_{\text{tan}} = \frac{\mathbf{a} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}} \mathbf{v}$ and $\mathbf{a}_{\text{norm}} = \mathbf{a} - \mathbf{a}_{\text{tan}}$.
- At any given point on a space curve we have three defining unit vectors:
 - a unit tangent,

Definitions

- If a curve is representing the path of a moving body, as usually is the case in Mechanics, then velocity and acceleration of the body are given as $\mathbf{v}(t) = \mathbf{r}'(t)$ and $\mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t)$.
- Acceleration vector has its tangential and normal components i.e. $\mathbf{a} = \mathbf{a}_{\text{tan}} + \mathbf{a}_{\text{norm}}$, which are obtained as $\mathbf{a}_{\text{tan}} = \frac{\mathbf{a} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}} \mathbf{v}$ and $\mathbf{a}_{\text{norm}} = \mathbf{a} - \mathbf{a}_{\text{tan}}$.
- At any given point on a space curve we have three defining unit vectors:
 - a **unit tangent**,
 - a **unit normal** which is perpendicular to unit tangent but lies in the same plane as that of tangent, and

Definitions

- If a curve is representing the path of a moving body, as usually is the case in Mechanics, then velocity and acceleration of the body are given as $\mathbf{v}(t) = \mathbf{r}'(t)$ and $\mathbf{a}(t) = \mathbf{v}'(t) = \mathbf{r}''(t)$.
- Acceleration vector has its tangential and normal components i.e. $\mathbf{a} = \mathbf{a}_{\text{tan}} + \mathbf{a}_{\text{norm}}$, which are obtained as $\mathbf{a}_{\text{tan}} = \frac{\mathbf{a} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}} \mathbf{v}$ and $\mathbf{a}_{\text{norm}} = \mathbf{a} - \mathbf{a}_{\text{tan}}$.
- At any given point on a space curve we have three defining unit vectors:
 - a **unit tangent**,
 - a **unit normal** which is perpendicular to unit tangent but lies in the same plane as that of tangent, and
 - a **unit binormal**, which is perpendicular to both i.e. unit tangent and unit normal vectors.



Question: (10ed-9.5-4,10) Sketch the curves
 $[-2, 2 + 5 \cos t, -1 + 5 \sin t]$ and $[t, 2, \frac{1}{t}]$?

Question: (10ed-9.5-4,10) Sketch the curves

$[-2, 2 + 5 \cos t, -1 + 5 \sin t]$ and $[t, 2, \frac{1}{t}]$?

Solution: We compute the table as

t	x-coord=-2	y-coord= $2 + 5 \cos t$	z-coord= $-1 + 5 \sin t$
0	-2	7	-1
2	-2	-0.008	3.5
3	-2	-2.9	-0.29
5	-2	3.4	-5.8


Question: (10ed-9.5-4,10) Sketch the curves

$[-2, 2 + 5 \cos t, -1 + 5 \sin t]$ and $[t, 2, \frac{1}{t}]$?

Solution: We compute the table as

t	x -coord= -2	y -coord= $2 + 5 \cos t$	z -coord= $-1 + 5 \sin t$
0	-2	7	-1
2	-2	-0.008	3.5
3	-2	-2.9	-0.29
5	-2	3.4	-5.8

Carefully plotting yields and by a similar computation for $[t, 2, \frac{1}{t}]$, we have:



Question: (10ed-9.5-11) Find parametric representation of a circle in the plane $z = 2$ with centre $(1, -1)$ and passing through origin.

Question: (10ed-9.5-11) Find parametric representation of a circle in the plane $z = 2$ with centre $(1, -1)$ and passing through origin.


Solution: Equation of a circle with centre $(1, -1)$ is given as $(x - 1)^2 + (y + 1)^2 = r^2$. As $(0, 0)$ is on the circle so we have $(0 - 1)^2 + (0 + 1)^2 = r^2 \Rightarrow r = \sqrt{2}$

Parametric equation of circle with centre (h, k) and radius r is given as

Question: (10ed-9.5-11) Find parametric representation of a circle in the plane $z = 2$ with centre $(1, -1)$ and passing through origin.

Solution: Equation of a circle with centre $(1, -1)$ is given as $(x - 1)^2 + (y + 1)^2 = r^2$. As $(0, 0)$ is on the circle so we have $(0 - 1)^2 + (0 + 1)^2 = r^2 \Rightarrow r = \sqrt{2}$

Parametric equation of circle with centre (h, k) and radius r is given as $[h + r \cos t, k + r \sin t]$. Hence the required parametric equation is $[1 + \sqrt{2} \cos t, -1 + \sqrt{2} \sin t]$. Sketch:





Question: (10ed-9.5-16) Find parametric representation of the intersection of the circular cylinder of radius 1 about z -axis and the plane $z = y$.

Question: (10ed-9.5-16) Find parametric representation of the intersection of the circular cylinder of radius 1 about z -axis and the plane $z = y$.

Solution: Equation of the cylinder is $x^2 + y^2 = 1$ and it calls for putting $x = \cos t$ and $y = \sin t$. This gives the equation of plane as $z = \sin t$. Hence the parametric equation of the circle of intersection is given as

$$[\cos t, \sin t, \sin t]$$

Sketches are given as



Question: (10ed-9.5-18) Helix: $x^2 + y^2 = 25, z = 2 \arctan \left(\frac{y}{x} \right)$.

Write its parametric equation.

Question: (10ed-9.5-18) Helix: $x^2 + y^2 = 25, z = 2 \arctan \left(\frac{y}{x} \right)$.

Write its parametric equation.


Solution: Put $x = 5 \cos t$ and

$y = 5 \sin t \Rightarrow z = 2 \arctan \left(\frac{5 \sin t}{5 \cos t} \right) = 2 \arctan (\tan t) = 2t$ Hence the parametric equation of the helix is given as

$$[5 \cos t, 5 \sin t, 2t]$$

Sketch:

$$[5 \cos(t), 5 \sin(t), 2t]$$



Question: (10ed-9.5-27) Given a curve $C : \mathbf{r}(t) = \left[t, \frac{4}{t}, 0 \right]$, find tangent vector $\mathbf{r}'(t)$, a unit tangent vector $\mathbf{u}'(t)$ and tangent of C at $P(4, 1, 0)$.

Question: (10ed-9.5-27) Given a curve $C : \mathbf{r}(t) = \left[t, \frac{4}{t}, 0 \right]$, find tangent vector $\mathbf{r}'(t)$, a unit tangent vector $\mathbf{u}'(t)$ and tangent of C at $P(4, 1, 0)$.

Solution: Tangent vector: $\mathbf{r}'(t) =$

Question: (10ed-9.5-27) Given a curve $C : \mathbf{r}(t) = \left[t, \frac{4}{t}, 0 \right]$, find tangent vector $\mathbf{r}'(t)$, a unit tangent vector $\mathbf{u}'(t)$ and tangent of C at $P(4, 1, 0)$.

Solution: Tangent vector: $\mathbf{r}'(t) = \frac{d}{dt} \left(\left[t, \frac{4}{t}, 0 \right] \right) = \left[1, -\frac{4}{t^2}, 0 \right]$

Question: (10ed-9.5-27) Given a curve $C : \mathbf{r}(t) = \left[t, \frac{4}{t}, 0 \right]$, find tangent vector $\mathbf{r}'(t)$, a unit tangent vector $\mathbf{u}'(t)$ and tangent of C at $P(4, 1, 0)$.

Solution: Tangent vector: $\mathbf{r}'(t) = \frac{d}{dt} \left(\left[t, \frac{4}{t}, 0 \right] \right) = \left[1, -\frac{4}{t^2}, 0 \right]$

Unit tangent vector: $\mathbf{u}' = \frac{1}{|\mathbf{r}'(t)|} \mathbf{r}'(t)$

Question: (10ed-9.5-27) Given a curve $C : \mathbf{r}(t) = \left[t, \frac{4}{t}, 0 \right]$, find tangent vector $\mathbf{r}'(t)$, a unit tangent vector $\mathbf{u}'(t)$ and tangent of C at $P(4, 1, 0)$.

Solution: Tangent vector: $\mathbf{r}'(t) = \frac{d}{dt} \left(\left[t, \frac{4}{t}, 0 \right] \right) = \left[1, -\frac{4}{t^2}, 0 \right]$

Unit tangent vector: $\mathbf{u}' = \frac{1}{|\mathbf{r}'(t)|} \mathbf{r}'(t) = \frac{1}{\left| \left[1, -\frac{4}{t^2}, 0 \right] \right|} \left[1, -\frac{4}{t^2}, 0 \right] =$

$$\frac{1}{\sqrt{\frac{16}{t^4} + 1}} \left[1, -\frac{4}{t^2}, 0 \right] = \left[\frac{1}{\sqrt{\frac{16}{t^4} + 1}}, -\frac{4}{t^2 \sqrt{\frac{16}{t^4} + 1}}, 0 \right]$$

Question: (10ed-9.5-27) Given a curve $C : \mathbf{r}(t) = [t, \frac{4}{t}, 0]$, find tangent vector $\mathbf{r}'(t)$, a unit tangent vector $\mathbf{u}'(t)$ and tangent of C at $P(4, 1, 0)$.

Solution: Tangent vector: $\mathbf{r}'(t) = \frac{d}{dt} ([t, \frac{4}{t}, 0]) = [1, -\frac{4}{t^2}, 0]$

Unit tangent vector: $\mathbf{u}' = \frac{1}{|\mathbf{r}'(t)|} \mathbf{r}'(t) = \frac{1}{|[1, -\frac{4}{t^2}, 0]|} [1, -\frac{4}{t^2}, 0] =$

$$\frac{1}{\sqrt{\frac{16}{t^4} + 1}} [1, -\frac{4}{t^2}, 0] = \left[\frac{1}{\sqrt{\frac{16}{t^4} + 1}}, -\frac{4}{t^2 \sqrt{\frac{16}{t^4} + 1}}, 0 \right]$$

Tangent line from $P : q(w) = [4 + w,]$

Question: (10ed-9.5-30) Find the length and sketch the curve given by $\mathbf{r}(t) = [4 \cos t, 4 \sin t, 5t]$ from $(4, 0, 0)$ to $(4, 0, 10\pi)$?

Solution: We have the formula $s(t) = \int_a^t \sqrt{\mathbf{r}' \cdot \mathbf{r}'} d\tau$.

Question: (10ed-9.5-30) Find the length and sketch the curve given by $\mathbf{r}(t) = [4 \cos t, 4 \sin t, 5t]$ from $(4, 0, 0)$ to $(4, 0, 10\pi)$?

Solution: We have the formula $s(t) = \int_a^t \sqrt{\mathbf{r}' \cdot \mathbf{r}'} d\tau$.

$$\mathbf{r}'(t) = \left[\frac{d}{dt}(4 \cos t), \frac{d}{dt}(4 \sin t), \frac{d}{dt}(5t) \right] = [-4 \sin t, 4 \cos t, 5] \Rightarrow \mathbf{r}' \cdot \mathbf{r}' = 16 \cos^2 t + 16 \sin^2 t + 25$$

As the point $(4, 0, 0)$ is on the curve, hence for some

$$t, \mathbf{r}(t) = (4 \cos t, 4 \sin t, 5t) = (4, 0, 0) \Rightarrow 5t = 0 \Rightarrow t = 0.$$

Also the second point $(4, 0, 10\pi)$ is on the curve, hence for some

$$t, \mathbf{r}(t) = (4 \cos t, 4 \sin t, 5t) = (4, 0, 10\pi) \Rightarrow 5t = 10\pi \Rightarrow t = 2\pi.$$

Hence putting values in the formula

$$s = \int_0^{2\pi} \sqrt{16 \cos^2 t + 16 \sin^2 t + 25} d\tau = 2\sqrt{41}\pi \quad \blacksquare$$

Question: (10ed-9.5-35) Find speed, velocity and tangential and normal acceleration for the parabola $\mathbf{r}(t) = [t, 4t^2, 0]$?

Question: (10ed-9.5-35) Find speed, velocity and tangential and normal acceleration for the parabola $\mathbf{r}(t) = [t, 4t^2, 0]$?


Solution: $\mathbf{v} = \mathbf{r}'(t) = [1, 8t, 0]$

$\mathbf{a} = \mathbf{v}' = \mathbf{r}''(t) = [0, 8, 0]$

$\mathbf{a}_{\tan} = \frac{\mathbf{a} \cdot \mathbf{v}}{\mathbf{v} \cdot \mathbf{v}} \mathbf{v} = \frac{[0, 8, 0] \cdot [1, 8t, 0]}{[1, 8t, 0] \cdot [1, 8t, 0]} [1, 8t, 0] = \frac{64t}{64t^2 + 1} [1, 8t, 0]$

$\mathbf{a}_{\text{norm}} = \mathbf{a} - \mathbf{a}_{\tan} = [0, 8, 0] - \frac{64t}{64t^2 + 1} [1, 8t, 0] =$
 $\left[\frac{-64t}{64t^2 + 1}, 8 - \frac{512t^2}{64t^2 + 1}, 0 \right]$

Sketch:



$[t, 4t^2, 0]$

Question: (10ed-9.5-46) A satellite in a circular orbit 450 miles above Earth's surface and completes 1 revolution in 100 min. Find the acceleration of gravity at the orbit from these data and from the radius of Earth (3960 miles)?

Question: (10ed-9.5-46) A satellite in a circular orbit 450 miles above Earth's surface and completes 1 revolution in 100 min. Find the acceleration of gravity at the orbit from these data and from the radius of Earth (3960 miles)?

Solution: $R = 3960 + 450 = 4410$ mi.

$$2\pi R = 100 |\mathbf{v}| \text{ and } \mathbf{v} = 277.1 \text{ mi/min}$$

$$g = |\mathbf{a}| = \omega^2 R = \frac{|\mathbf{v}|^2}{R} = 17.41 \text{ mi/min}^2 = 25.53 \text{ ft/sec}^2 = 7.78 \text{ m/sec}^2 \quad \blacksquare$$

Gradient of a Scalar Field

Definition

Gradient of a scalar function $f(x, y, z)$ is denoted as $\text{grad } f$ or ∇f (read as **nabla** f) and is defined as

$$\text{grad } f = \nabla f = \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right]$$

Gradient of a Scalar Field

Definition

Gradient of a scalar function $f(x, y, z)$ is denoted as $\text{grad } f$ or ∇f (read as **nabla** f) and is defined as

$$\text{grad } f = \nabla f = \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} + \frac{\partial f}{\partial z} \mathbf{k}$$

Gradient of a Scalar Field

Definition

Gradient of a scalar function $f(x, y, z)$ is denoted as $\text{grad } f$ or ∇f (read as **nabla** f) and is defined as

$$\text{grad } f = \nabla f = \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} + \frac{\partial f}{\partial z} \mathbf{k}$$

We also write the **differential operator** as $\nabla = \frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k}$.

Major Usages of Gradient

- 1 For rate of change of $f(x, y, z)$ in any direction in space, technically called Directional Derivative.

Major Usages of Gradient

- ① For rate of change of $f(x, y, z)$ in any direction in space, technically called Directional Derivative.
 - ① The directional derivative $D_{\mathbf{a}}f$ of a scalar function $f(x, y, z)$ at a point P in the direction of a vector \mathbf{a} is given as
$$D_{\mathbf{a}}f = \frac{1}{|\mathbf{a}|} \mathbf{a} \cdot \text{grad } f.$$

Major Usages of Gradient


- 1 For rate of change of $f(x, y, z)$ in any direction in space, technically called Directional Derivative.
 - 1 The directional derivative $D_{\mathbf{a}}f$ of a scalar function $f(x, y, z)$ at a point P in the direction of a vector \mathbf{a} is given as
$$D_{\mathbf{a}}f = \frac{1}{|\mathbf{a}|} \mathbf{a} \cdot \text{grad } f.$$
- 2 $\text{grad } f = \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right]$ points in the direction of maximum increase of $f(x, y, z)$.

Major Usages of Gradient

- 1 For rate of change of $f(x, y, z)$ in any direction in space, technically called Directional Derivative.
 - 1 The directional derivative $D_{\mathbf{a}}f$ of a scalar function $f(x, y, z)$ at a point P in the direction of a vector \mathbf{a} is given as
$$D_{\mathbf{a}}f = \frac{1}{|\mathbf{a}|} \mathbf{a} \cdot \text{grad } f.$$
- 2 $\text{grad } f = \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right]$ points in the direction of maximum increase of $f(x, y, z)$.
- 3 For a curve $C = [x(t), y(t), z(t)]$ lying on a level surface $f(x, y, z) = c = \text{const}$,

Major Usages of Gradient

- 1 For rate of change of $f(x, y, z)$ in any direction in space, technically called Directional Derivative.
 - 1 The directional derivative $D_{\mathbf{a}}f$ of a scalar function $f(x, y, z)$ at a point P in the direction of a vector \mathbf{a} is given as
$$D_{\mathbf{a}}f = \frac{1}{|\mathbf{a}|} \mathbf{a} \cdot \text{grad } f.$$
- 2 $\text{grad } f = \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right]$ points in the direction of maximum increase of $f(x, y, z)$.
- 3 For a curve $C = [x(t), y(t), z(t)]$ lying on a level surface $f(x, y, z) = c = \text{const}$, $\text{grad } f$ is normal vector of S at P .



Potential

One of the most important usage of gradient is in obtaining vector field from a scalar field.

Potential

One of the most important usage of gradient is in obtaining vector field from a scalar field. A vector field obtained in this manner is relatively easily studied using $f(x, y, z)$ only.

Potential

One of the most important usage of gradient is in obtaining vector field from a scalar field. A vector field obtained in this manner is relatively easily studied using $f(x, y, z)$ only. For such vector fields, $f(x, y, z)$ is said to be its **potential**.

Potential

One of the most important usage of gradient is in obtaining vector field from a scalar field. A vector field obtained in this manner is relatively easily studied using $f(x, y, z)$ only. For such vector fields, $f(x, y, z)$ is said to be its **potential**. Furthermore such vector field is said to be **conservative** if no energy is lost or gained in displacing a body from one point to another and then back.

Potential

One of the most important usage of gradient is in obtaining vector field from a scalar field. A vector field obtained in this manner is relatively easily studied using $f(x, y, z)$ only. For such vector fields, $f(x, y, z)$ is said to be its **potential**. Furthermore such vector field is said to be **conservative** if no energy is lost or gained in displacing a body from one point to another and then back.

Definition

The potential $f(x, y, z)$ of a conservative vector field satisfies the Laplace's equation, given as

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = 0$$

Potential

One of the most important usage of gradient is in obtaining vector field from a scalar field. A vector field obtained in this manner is relatively easily studied using $f(x, y, z)$ only. For such vector fields, $f(x, y, z)$ is said to be its **potential**. Furthermore such vector field is said to be **conservative** if no energy is lost or gained in displacing a body from one point to another and then back.

Definition

The potential $f(x, y, z)$ of a conservative vector field satisfies the Laplace's equation, given as

$$\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = 0$$

It is universally agreed that Laplace equation is The Most Important partial differential equation in today's Physics and its numerous applications.

Question: (10ed-9.7-4) Find $\text{grad } f$ where
 $f = (x - 2)^2 + (2y + 4)^2$. Graph some level curves $f = \text{const}$
Indicate ∇f by arrows at some points of these curves?

Question: (10ed-9.7-4) Find $\text{grad } f$ where

$f = (x - 2)^2 + (2y + 4)^2$. Graph some level curves $f = \text{const}$

Indicate ∇f by arrows at some points of these curves?

Solution:

$$\begin{aligned} & \text{grad} \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial}{\partial x} \left((x - 2)^2 + (2y + 4)^2 \right), \right. \\ & \quad \frac{\partial}{\partial y} \left((x - 2)^2 + (2y + 4)^2 \right), \\ & \quad \left. \frac{\partial}{\partial z} \left((x - 2)^2 + (2y + 4)^2 \right) \right] \end{aligned}$$

Question: (10ed-9.7-4) Find **grad** f where $f = (x - 2)^2 + (2y + 4)^2$. Graph some level curves $f = \text{const}$. Indicate ∇f by arrows at some points of these curves?

Solution:

$$\begin{aligned} & \text{grad} \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial}{\partial x} \left((x - 2)^2 + (2y + 4)^2 \right), \right. \\ & \quad \left. \frac{\partial}{\partial y} \left((x - 2)^2 + (2y + 4)^2 \right), \right. \\ & \quad \left. \frac{\partial}{\partial z} \left((x - 2)^2 + (2y + 4)^2 \right) \right] \\ &= [2x - 4, 8y + 16, 0] \end{aligned}$$

Question: (10ed-9.7-4) Find **grad** f where

$f = (x - 2)^2 + (2y + 4)^2$. Graph some level curves $f = \text{const}$

Indicate ∇f by arrows at some points of these curves?

Solution:

$$\begin{aligned} & \text{grad} \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial}{\partial x} \left((x - 2)^2 + (2y + 4)^2 \right), \right. \\ & \quad \frac{\partial}{\partial y} \left((x - 2)^2 + (2y + 4)^2 \right), \\ & \quad \left. \frac{\partial}{\partial z} \left((x - 2)^2 + (2y + 4)^2 \right) \right] \\ &= [2x - 4, 8y + 16, 0] \end{aligned}$$

We choose points $(-1, 2)$, $(2, 1)$ and $(5, -5)$ for computation of gradient vectors. Sketch is given below:

Question: (10ed-9.7-4) Find $\text{grad } f$ where

$f = (x - 2)^2 + (2y + 4)^2$. Graph some level curves $f = \text{const}$


Indicate ∇f by arrows at some points of these curves?

Solution:

$$\begin{aligned} & \text{grad} \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] \left((x - 2)^2 + (2y + 4)^2 \right) \\ &= \left[\frac{\partial}{\partial x} \left((x - 2)^2 + (2y + 4)^2 \right), \right. \\ & \quad \frac{\partial}{\partial y} \left((x - 2)^2 + (2y + 4)^2 \right), \\ & \quad \left. \frac{\partial}{\partial z} \left((x - 2)^2 + (2y + 4)^2 \right) \right] \\ &= [2x - 4, 8y + 16, 0] \end{aligned}$$

We choose points $(-1, 2)$, $(2, 1)$ and $(5, -5)$ for computation of gradient vectors. Sketch is given below:

Gradient Vectors



For the function $f(x, y) = (x - 2)^2 + (2y + 4)^2$, level curves, their projections to the xy -plane, and gradient vectors at the point(s) $[(-1, 2), (2, 1), (5, -5)]$.

Question: (10ed-9.7-10) Prove that

$$\nabla^2 (fg) = g \nabla^2 f + 2 \nabla f \cdot \nabla g + f \nabla^2 g?$$

Question: (10ed-9.7-10) Prove that

$$\nabla^2 (fg) = g \nabla^2 f + 2 \nabla f \cdot \nabla g + f \nabla^2 g?$$

Proof:

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg)$$

Question: (10ed-9.7-10) Prove that
 $\nabla^2 (fg) = g \nabla^2 f + 2 \nabla f \cdot \nabla g + f \nabla^2 g$?

Proof:

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg)$$

$$\text{Consider } \frac{\partial^2}{\partial x^2} (fg) = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} (fg) \right) = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} g + f \frac{\partial g}{\partial x} \right) = \\ \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + f \frac{\partial^2 g}{\partial x^2} \right)$$

Question: (10ed-9.7-10) Prove that

$$\nabla^2 (fg) = g \nabla^2 f + 2 \nabla f \cdot \nabla g + f \nabla^2 g?$$

Proof:

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg)$$

$$\text{Consider } \frac{\partial^2}{\partial x^2} (fg) = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} (fg) \right) = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} g + f \frac{\partial g}{\partial x} \right) = \\ \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + f \frac{\partial^2 g}{\partial x^2} \right)$$

Similar computation for $\frac{\partial^2}{\partial y^2} (fg)$ and $\frac{\partial^2}{\partial z^2} (fg)$ yield

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg)$$

Question: (10ed-9.7-10) Prove that

$$\nabla^2 (fg) = g \nabla^2 f + 2 \nabla f \cdot \nabla g + f \nabla^2 g?$$

Proof:

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg)$$

$$\text{Consider } \frac{\partial^2}{\partial x^2} (fg) = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} (fg) \right) = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} g + f \frac{\partial g}{\partial x} \right) = \\ \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + f \frac{\partial^2 g}{\partial x^2} \right)$$

Similar computation for $\frac{\partial^2}{\partial y^2} (fg)$ and $\frac{\partial^2}{\partial z^2} (fg)$ yield

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg) \\ = \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + f \frac{\partial^2 g}{\partial x^2} \right) + \left(\frac{\partial^2 f}{\partial y^2} g + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} \right) + \\ \left(\frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + f \frac{\partial^2 g}{\partial y^2} \right) + \left(\frac{\partial^2 f}{\partial z^2} g + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \left(\frac{\partial f}{\partial z} \frac{\partial g}{\partial z} + f \frac{\partial^2 g}{\partial z^2} \right)$$

Question: (10ed-9.7-10) Prove that

$$\nabla^2 (fg) = g \nabla^2 f + 2 \nabla f \cdot \nabla g + f \nabla^2 g?$$

Proof:

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg)$$

$$\text{Consider } \frac{\partial^2}{\partial x^2} (fg) = \frac{\partial}{\partial x} \left(\frac{\partial}{\partial x} (fg) \right) = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} g + f \frac{\partial g}{\partial x} \right) = \\ \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + f \frac{\partial^2 g}{\partial x^2} \right)$$

Similar computation for $\frac{\partial^2}{\partial y^2} (fg)$ and $\frac{\partial^2}{\partial z^2} (fg)$ yield

$$\nabla^2 (fg) = \frac{\partial^2}{\partial x^2} (fg) + \frac{\partial^2}{\partial y^2} (fg) + \frac{\partial^2}{\partial z^2} (fg) \\ = \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + f \frac{\partial^2 g}{\partial x^2} \right) + \left(\frac{\partial^2 f}{\partial y^2} g + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} \right) + \\ \left(\frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + f \frac{\partial^2 g}{\partial y^2} \right) + \left(\frac{\partial^2 f}{\partial z^2} g + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \left(\frac{\partial f}{\partial z} \frac{\partial g}{\partial z} + f \frac{\partial^2 g}{\partial z^2} \right) \\ = \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial f}{\partial x} \frac{\partial g}{\partial x} \right) + \left(\frac{\partial^2 f}{\partial y^2} g + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} \right) + \left(\frac{\partial^2 f}{\partial z^2} g + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \\ \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + f \frac{\partial^2 g}{\partial x^2} \right) + \left(\frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + f \frac{\partial^2 g}{\partial y^2} \right) + \left(\frac{\partial f}{\partial z} \frac{\partial g}{\partial z} + f \frac{\partial^2 g}{\partial z^2} \right)$$

$$= \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial^2 f}{\partial y^2} g + \frac{\partial^2 f}{\partial z^2} g \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) +$$

$$\left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \left(f \frac{\partial^2 g}{\partial x^2} + f \frac{\partial^2 g}{\partial y^2} + f \frac{\partial^2 g}{\partial z^2} \right)$$

$$\begin{aligned}
&= \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial^2 f}{\partial y^2} g + \frac{\partial^2 f}{\partial z^2} g \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \\
&\quad \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \left(f \frac{\partial^2 g}{\partial x^2} + f \frac{\partial^2 g}{\partial y^2} + f \frac{\partial^2 g}{\partial z^2} \right) \\
&= g \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \right) + 2 \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \\
&\quad f \left(\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 g}{\partial z^2} \right) \\
&= g \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \right) + 2 \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] \cdot \left[\frac{\partial g}{\partial x}, \frac{\partial g}{\partial y}, \frac{\partial g}{\partial z} \right] + \\
&\quad f \left(\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 g}{\partial z^2} \right)
\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{\partial^2 f}{\partial x^2} g + \frac{\partial^2 f}{\partial y^2} g + \frac{\partial^2 f}{\partial z^2} g \right) + \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \\
&\quad \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \left(f \frac{\partial^2 g}{\partial x^2} + f \frac{\partial^2 g}{\partial y^2} + f \frac{\partial^2 g}{\partial z^2} \right) \\
&= g \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \right) + 2 \left(\frac{\partial f}{\partial x} \frac{\partial g}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial g}{\partial y} + \frac{\partial f}{\partial z} \frac{\partial g}{\partial z} \right) + \\
&\quad f \left(\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 g}{\partial z^2} \right) \\
&= g \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} \right) + 2 \left[\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right] \cdot \left[\frac{\partial g}{\partial x}, \frac{\partial g}{\partial y}, \frac{\partial g}{\partial z} \right] + \\
&\quad f \left(\frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2} + \frac{\partial^2 g}{\partial z^2} \right) \\
&= g \nabla^2 f + 2 \nabla f \cdot \nabla g + f \nabla^2 g \quad \blacksquare
\end{aligned}$$

Question: (10ed-9.7-14) The force in an electric field given by $f(x, y, z) = (x^2 + y^2 + z^2)^{-\frac{1}{2}}$ has the direction of the gradient. Find ∇f and its value at $P(12, 0, 16)$?

Question: (10ed-9.7-14) The force in an electric field given by $f(x, y, z) = (x^2 + y^2 + z^2)^{-\frac{1}{2}}$ has the direction of the gradient. Find ∇f and its value at $P(12, 0, 16)$?

Solution: $\nabla f = \nabla \left((x^2 + y^2 + z^2)^{-\frac{1}{2}} \right) =$

$$\left[\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right] \left((x^2 + y^2 + z^2)^{-\frac{1}{2}} \right) = \frac{-1}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} [x, y, z]$$


$$\nabla f|_P = \left(\frac{-1}{(x^2 + y^2 + z^2)^{\frac{3}{2}}} [x, y, z] \right)_{x=12, y=0, z=16} = \frac{1}{500} \left[\frac{-3}{4}, 0, -1 \right] =$$

$$\frac{-3}{2000} \mathbf{i} - \frac{1}{500} \mathbf{k} \quad \blacksquare$$

Question: (10ed-9.7-16) For what points $P(x,y,z)$ does ∇f with $f = 25x^2 + 9y^2 + 16z^2$ have the direction from P to the origin?

Question: (10ed-9.7-16) For what points $P(x,y,z)$ does ∇f with $f = 25x^2 + 9y^2 + 16z^2$ have the direction from P to the origin?



Solution: $\nabla (25x^2 + 9y^2 + 16z^2) = [50x, 18y, 32z]$. Presence of integer multiples of x, y and z only in the gradient indicates that all points on any of the three axes would have direction from P to origin. Sketch of gradient vector field is given as



Question: (10ed-9.7-20) Given the velocity potential $f = x \left(1 + (x^2 + y^2)^{-1} \right)$, find the velocity $\mathbf{v} = \nabla f$ of the field and its value $\mathbf{v}(P)$ at P . Sketch $\mathbf{v}(P)$ and the curve $f = \text{const}$ passing through $P(1, 1)$?

Question: (10ed-9.7-20) Given the velocity potential $f = x \left(1 + (x^2 + y^2)^{-1} \right)$, find the velocity $\mathbf{v} = \nabla f$ of the field and its value $\mathbf{v}(P)$ at P . Sketch $\mathbf{v}(P)$ and the curve $f = \text{const}$ passing through $P(1,1)$?

Solution: $\mathbf{v} = \nabla f = \nabla \left(x \left(1 + (x^2 + y^2)^{-1} \right) \right) = \left[\frac{1}{x^2 + y^2} - \frac{2x^2}{(x^2 + y^2)^2} + 1, -2x \frac{y}{(x^2 + y^2)^2}, 0 \right]$ and this velocity field is sketched below (two different views) as follows:




Equation of curve is of the form $x \left(1 + (x^2 + y^2)^{-1} \right) = c$, since the curve has to pass through $P(1, 1)$, hence to find c put $x = 1, y = 1$ in f i.e. $x \left(1 + (x^2 + y^2)^{-1} \right) |_{(1,1)} = \frac{3}{2} = c$.

Equation of curve passing through $P(1, 1)$ is

$$x \left(1 + (x^2 + y^2)^{-1} \right) = \frac{3}{2}.$$

Equation of curve is of the form $x \left(1 + (x^2 + y^2)^{-1} \right) = c$, since the curve has to pass through $P(1, 1)$, hence to find c put $x = 1, y = 1$ in f i.e. $x \left(1 + (x^2 + y^2)^{-1} \right) |_{(1,1)} = \frac{3}{2} = c$.

Equation of curve passing through $P(1, 1)$ is $x \left(1 + (x^2 + y^2)^{-1} \right) = \frac{3}{2}$. Sketch of the required curve is an implicit graph and is given as:



Question: (10ed-9.7-25) Experiments show that in a temperature field, heat flows in the direction of maximum decrease of temperature $T = \frac{z}{(x^2+y^2)}$. Find this direction in general and at the given point $P(0, 1, 2)$. Sketch that direction at P as an arrow.

Question: (10ed-9.7-25) Experiments show that in a temperature field, heat flows in the direction of maximum decrease of temperature $T = \frac{z}{(x^2+y^2)}$. Find this direction in general and at the given point $P(0, 1, 2)$. Sketch that direction at P as an arrow.

Solution: As ∇f points in the direction of maximum increase hence

Direction of maximum

decrease = $-\nabla T$

Question: (10ed-9.7-25) Experiments show that in a temperature field, heat flows in the direction of maximum decrease of temperature $T = \frac{z}{(x^2+y^2)}$. Find this direction in general and at the given point $P(0, 1, 2)$. Sketch that direction at P as an arrow.

Solution: As ∇f points in the direction of maximum increase hence

Direction of maximum

$$\text{decrease} = -\nabla T = -\nabla \left(\frac{z}{(x^2+y^2)} \right) = \left[\frac{2xz}{(x^2+y^2)^2}, \frac{2yz}{(x^2+y^2)^2}, -\frac{1}{x^2+y^2} \right]$$

Question: (10ed-9.7-25) Experiments show that in a temperature field, heat flows in the direction of maximum decrease of temperature $T = \frac{z}{(x^2+y^2)}$. Find this direction in general and at the given point $P(0, 1, 2)$. Sketch that direction at P as an arrow.

Solution: As ∇f points in the direction of maximum increase hence


Direction of maximum

$$\text{decrease} = -\nabla T = -\nabla \left(\frac{z}{(x^2+y^2)} \right) = \left[\frac{2xz}{(x^2+y^2)^2}, \frac{2yz}{(x^2+y^2)^2}, -\frac{1}{x^2+y^2} \right]$$

Direction of maximum decrease at

$$P = \left[\frac{2xz}{(x^2+y^2)^2}, \frac{2yz}{(x^2+y^2)^2}, -\frac{1}{x^2+y^2} \right]_{x=0, y=1, z=2} = [0, 4, -1]$$

Sketch of general direction of the field and at point P are given as



Question: (10ed-9.7-33) Find the normal vector of the surface $6x^2 + 2y^2 + z^2 = 225$ at the point $P(5, 5, 5)$?

Question: (10ed-9.7-33) Find the normal vector of the surface $6x^2 + 2y^2 + z^2 = 225$ at the point $P(5, 5, 5)$?


Solution: Since on a surface $f(x, y, z) = \text{const}$ normal vector is given by ∇f , hence we compute

$\nabla(6x^2 + 2y^2 + z^2) = [12x, 4y, 2z]$ and the normal at the given point $= [12x, 4y, 2z]_{x=5, y=5, z=5} = [60, 20, 10]$.

Question: (10ed-9.7-33) Find the normal vector of the surface $6x^2 + 2y^2 + z^2 = 225$ at the point $P(5, 5, 5)$?

Solution: Since on a surface $f(x, y, z) = \text{const}$ normal vector is given by ∇f , hence we compute

$\nabla(6x^2 + 2y^2 + z^2) = [12x, 4y, 2z]$ and the normal at the given point = $[12x, 4y, 2z]_{x=5, y=5, z=5} = [60, 20, 10]$. We sketch the surface and the normal as



Question: (10ed-9.7-39) Find directional derivative of $f(x, y, z) = \frac{1}{\sqrt{x^2 + y^2 + z^2}}$ at $P(3, 0, 4)$ in the direction of $\mathbf{a} = [1, 1, 1]$. Also sketch it.

Question: (10ed-9.7-39) Find directional derivative of $f(x, y, z) = \frac{1}{\sqrt{x^2 + y^2 + z^2}}$ at $P(3, 0, 4)$ in the direction of $\mathbf{a} = [1, 1, 1]$. Also sketch it.

Solution: We have $D_{\mathbf{a}}f = \frac{1}{|\mathbf{a}|} \mathbf{a} \cdot \text{grad } f$.

Question: (10ed-9.7-39) Find directional derivative of $f(x, y, z) = \frac{1}{\sqrt{x^2+y^2+z^2}}$ at $P(3, 0, 4)$ in the direction of $\mathbf{a} = [1, 1, 1]$. Also sketch it.

Solution: We have $D_{\mathbf{a}}f = \frac{1}{|\mathbf{a}|} \mathbf{a} \cdot \text{grad } f$. So we compute

$$\begin{aligned}\text{grad } f &= \nabla \left(\frac{1}{\sqrt{x^2+y^2+z^2}} \right) = \\ &\left[-\frac{x}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{y}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{z}{(x^2+y^2+z^2)^{\frac{3}{2}}} \right] \\ D_{\mathbf{a}}f &= \\ &\left[\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right] \cdot \left[-\frac{x}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{y}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{z}{(x^2+y^2+z^2)^{\frac{3}{2}}} \right] \\ D_{\mathbf{a}}f|_{(3,0,4)} &= \\ &\left[\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right] \cdot \left[-\frac{x}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{y}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{z}{(x^2+y^2+z^2)^{\frac{3}{2}}} \right]_{(3,0,4)} \\ &= \left[\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right] \cdot \left[-\frac{3}{125}, 0, -\frac{4}{125} \right] \\ &= -\frac{7}{375} \sqrt{3} = -.032\end{aligned}$$


Question: (10ed-9.7-39) Find directional derivative of $f(x, y, z) = \frac{1}{\sqrt{x^2+y^2+z^2}}$ at $P(3, 0, 4)$ in the direction of $\mathbf{a} = [1, 1, 1]$. Also sketch it.

Solution: We have $D_{\mathbf{a}}f = \frac{1}{|\mathbf{a}|} \mathbf{a} \cdot \text{grad } f$. So we compute

$$\begin{aligned} \text{grad } f &= \nabla \left(\frac{1}{\sqrt{x^2+y^2+z^2}} \right) = \\ &= \left[-\frac{x}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{y}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{z}{(x^2+y^2+z^2)^{\frac{3}{2}}} \right] \\ D_{\mathbf{a}}f &= \left[\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right] \cdot \left[-\frac{x}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{y}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{z}{(x^2+y^2+z^2)^{\frac{3}{2}}} \right] \\ D_{\mathbf{a}}f|_{(3,0,4)} &= \left[\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right] \cdot \left[-\frac{x}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{y}{(x^2+y^2+z^2)^{\frac{3}{2}}}, -\frac{z}{(x^2+y^2+z^2)^{\frac{3}{2}}} \right]_{(3,0,4)} \\ &= \left[\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right] \cdot \left[-\frac{3}{125}, 0, -\frac{4}{125} \right] \\ &= -\frac{7}{375} \sqrt{3} = -.032 \end{aligned}$$

Hence the required vector is

$\left[-\frac{3}{125}, 0, -\frac{4}{125} \right]$ at the point $P(3, 0, 4)$



Question: Find a potential $f = \text{grad } f$ for the given
 $\mathbf{v}(x, y, z) = [ye^x, e^x, z^2]$?

Question: Find a potential $f = \text{grad } f$ for the given

$$\mathbf{v}(x, y, z) = [ye^x, e^x, z^2]?$$

Solution: By examining $\mathbf{v} = [ye^x, e^x, z^2]$, we conclude that

$$f(x, y, z) = ye^x + \frac{1}{3}z^3,$$

Question: Find a potential $f = \text{grad } f$ for the given

$$\mathbf{v}(x, y, z) = [ye^x, e^x, z^2]?$$

Solution: By examining $\mathbf{v} = [ye^x, e^x, z^2]$, we conclude that $f(x, y, z) = ye^x + \frac{1}{3}z^3$, because

$$\begin{aligned}\text{grad } f &= \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) f(x, y, z) \\ &= \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \right) \left(ye^x + \frac{1}{3}z^3 \right) \\ &= [ye^x, e^x, z^2] \quad \blacksquare\end{aligned}$$

Question: Find a unit normal vector \mathbf{n} of the cone of revolution $z^2 = 4(x^2 + y^2)$ at the point $P : (1, 0, 2)$?

Question: Find a unit normal vector \mathbf{n} of the cone of revolution $z^2 = 4(x^2 + y^2)$ at the point $P : (1, 0, 2)$?

Solution: The cone is the level surface $f = 0$ of $f(x, y, z) = 4(x^2 + y^2) - z^2$. Thus

$$\nabla f = [8x, 8y, -2z], \quad \nabla f(P) = [8, 0, -4]$$

$$\mathbf{n} = \frac{1}{|\nabla f(P)|} \nabla f(P) = \left[\frac{2}{\sqrt{5}}, 0, \frac{-1}{\sqrt{5}} \right]$$

\mathbf{n} points downward since it has a negative z -component. The other unit normal vector of the cone at P is $-\mathbf{n}$. ■

Definition: Divergence of a Vector Field

For a differentiable vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$, the divergence is denoted and defined as $\text{div } \mathbf{v} = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} + \frac{\partial v_3}{\partial z}$.

Definition: Divergence of a Vector Field

For a differentiable vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$, the divergence is denoted and defined as $\text{div } \mathbf{v} = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} + \frac{\partial v_3}{\partial z}$.

$\text{div } \mathbf{v}$ is also denoted as $\nabla \cdot \mathbf{v}$ because $\text{div } \mathbf{v} = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} + \frac{\partial v_3}{\partial z} = \left[\frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right] \cdot [v_1 \mathbf{i} + v_2 \mathbf{j} + v_3 \mathbf{k}] = \nabla \cdot \mathbf{v}$

Definition: Divergence of a Vector Field

For a differentiable vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$, the divergence is denoted and defined as $\text{div } \mathbf{v} = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} + \frac{\partial v_3}{\partial z}$.

$\text{div } \mathbf{v}$ is also denoted as $\nabla \cdot \mathbf{v}$ because $\text{div } \mathbf{v} = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} + \frac{\partial v_3}{\partial z} = \left[\frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right] \cdot [v_1 \mathbf{i} + v_2 \mathbf{j} + v_3 \mathbf{k}] = \nabla \cdot \mathbf{v}$

If $\mathbf{v} = \text{grad}(f)$ and f is twice differentiable scalar function $f(x, y, z)$, we have

$$\text{div } \mathbf{v} = \text{div}(\text{grad } f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = \nabla^2 f.$$

Definition: Divergence of a Vector Field


For a differentiable vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$, the divergence is denoted and defined as $\text{div } \mathbf{v} = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} + \frac{\partial v_3}{\partial z}$.

$\text{div } \mathbf{v}$ is also denoted as $\nabla \cdot \mathbf{v}$ because $\text{div } \mathbf{v} = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} + \frac{\partial v_3}{\partial z} = \left[\frac{\partial}{\partial x} \mathbf{i} + \frac{\partial}{\partial y} \mathbf{j} + \frac{\partial}{\partial z} \mathbf{k} \right] \cdot [v_1 \mathbf{i} + v_2 \mathbf{j} + v_3 \mathbf{k}] = \nabla \cdot \mathbf{v}$

If $\mathbf{v} = \text{grad}(f)$ and f is twice differentiable scalar function $f(x, y, z)$, we have

$$\text{div } \mathbf{v} = \text{div}(\text{grad } f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = \nabla^2 f.$$



Visualization of Divergence

From a scalar field we can obtain a vector field by the gradient.

Usage of Divergence

From a scalar field we can obtain a vector field by the gradient.
Conversely, from a vector field we can obtain a scalar field by the divergence.

Usage of Divergence

From a scalar field we can obtain a vector field by the gradient.
Conversely, from a vector field we can obtain a scalar field by the divergence.



We now intend to present a physical interpretation of the notion of divergence.

We now intend to present a physical interpretation of the notion of divergence. For this we need following definitions:

Definition

Flux is the total loss of mass leaving an object per unit of time.

We now intend to present a physical interpretation of the notion of divergence. For this we need following definitions:

Definition

Flux is the total loss of mass leaving an object per unit of time.

Definition

Compressible fluid is the fluid whose density ρ (mass per unit volume) depends upon coordinates x, y, z (and possibly on time t). Examples are gases and vapors. Water is an incompressible fluid.

We now intend to present a physical interpretation of the notion of divergence. For this we need following definitions:

Definition

Flux is the total loss of mass leaving an object per unit of time.

Definition

Compressible fluid is the fluid whose density ρ (mass per unit volume) depends upon coordinates x, y, z (and possibly on time t). Examples are gases and vapors. Water is an incompressible fluid.

Definition

If density ρ is independent of time t , the flow is said to be **steady**.


Question: Give a physical interpretation of divergence? OR Derive equation of continuity for fluids?

Question: Give a physical interpretation of divergence? OR Derive equation of continuity for fluids?

Derivation: Consider motion of a compressible fluid in region R with no source or sink in R . Consider flow of the fluid through the box B with volume $\Delta V = \Delta x \Delta y \Delta z$ (here Δ is denoting a small quantity and not the Laplacian).

Question: Give a physical interpretation of divergence? OR Derive equation of continuity for fluids?


Derivation: Consider motion of a compressible fluid in region R with no source or sink in R . Consider flow of the fluid through the box B with volume $\Delta V = \Delta x \Delta y \Delta z$ (here Δ is denoting a small quantity and not the Laplacian).



Physical interpretation of the divergence

Question: Give a physical interpretation of divergence? OR Derive equation of continuity for fluids?

Derivation: Consider motion of a compressible fluid in region R with no source or sink in R . Consider flow of the fluid through the box B with volume $\Delta V = \Delta x \Delta y \Delta z$ (here Δ is denoting a small quantity and not the Laplacian).



Physical interpretation of the divergence

Let $\mathbf{v} = [v_1, v_2, v_3] = v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}$ be the velocity vector of the motion. We set

$$\mathbf{u} = \rho \mathbf{v} = [u_1, u_2, u_3] = u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}$$

Consider the flow through the xz face whose area is $\Delta x \Delta z$. Since the vectors $v_1 \mathbf{i}$ and $v_3 \mathbf{k}$ are parallel to xz face, the components v_1 and v_3 contribute nothing to this flow.

Consider the flow through the xz face whose area is $\Delta x \Delta z$. Since the vectors $v_1 \mathbf{i}$ and $v_3 \mathbf{k}$ are parallel to xz face, the components v_1 and v_3 contribute nothing to this flow. Hence the mass of fluid entering through xz face during a short time interval Δt is

$$(\rho v_2) \Delta x \Delta z \Delta t = (u_2)_y \Delta x \Delta z \Delta t$$

and the mass leaving from opposite face is $(u_2)_{y+\Delta y} \Delta x \Delta z \Delta t$. Hence the difference

$$\Delta u_2 \Delta x \Delta y \Delta z = \frac{\Delta u_2}{\Delta y} \Delta y \Delta t \quad \text{where} \quad \Delta u_2 = (u_2)_{y+\Delta y} - (u_2)_y$$

is the approximate loss of mass. Other two faces also give similar expressions and the total loss of mass in B during the time interval Δt is approximately

$$\left(\frac{\Delta u_1}{\Delta x} + \frac{\Delta u_2}{\Delta y} + \frac{\Delta u_3}{\Delta z} \right) \Delta V \Delta t \quad (1)$$

where $\Delta u_1 = (u_1)_{x+\Delta x} - (u_1)_x$ and $\Delta u_3 = (u_3)_{z+\Delta z} - (u_3)_z$.

This loss of mass in B is caused by the time rate of change of the density and is thus equals to

$$-\frac{\partial \rho}{\partial t} \Delta V \Delta t \quad (2)$$

Equating (1) and (2) and letting small changes approach to zero we get

$$\begin{aligned} \text{div } \mathbf{u} &= \text{div } (\rho \mathbf{v}) = -\frac{\partial \rho}{\partial t} \\ \frac{\partial \rho}{\partial t} + \text{div } (\rho \mathbf{v}) &= 0 \end{aligned}$$

is the '**equation of continuity** of a compressible fluid flow' also called 'condition for the coservation of mass'.


If the flow is steady then $\frac{\partial \rho}{\partial t} = 0$ and the equation becomes $\text{div } (\rho \mathbf{v}) = 0$ and if the density is constant i.e. the fluid is incompressible then $\text{div } \mathbf{v} = 0$. ■

Question: (10ed-9.8-5) Find $\text{div } \mathbf{v}$ at $P(-1, 3, -2)$ where $\mathbf{v} = [x^2yz, xy^2z, xyz^2]$?

Question: (10ed-9.8-5) Find $\text{div } \mathbf{v}$ at $P(-1, 3, -2)$ where $\mathbf{v} = [x^2yz, xy^2z, xyz^2]$?

Solution: $\text{div } \mathbf{v} = \text{div}([x^2yz, xy^2z, xyz^2]) = 6xyz \Rightarrow$
 $\text{div } \mathbf{v}|_{P(-1,3,-2)} = 6(-1 \times 3 \times -2) = 36$

The vector field $\mathbf{v} = [x^2yz, xy^2z, xyz^2]$ around a box is sketched below:



The vector field arrows, the surface through which the field passes, and vectors normal to the surface.



Question: (10ed-9.8-17) Find $\nabla^2 f$ by the formula $\nabla^2 f = \text{div}(\text{grad } f)$, where $f = \ln(x^2 + y^2)$?

Question: (10ed-9.8-17) Find $\nabla^2 f$ by the formula

$\nabla^2 f = \text{div}(\text{grad } f)$, where $f = \ln(x^2 + y^2)$?

Solution: $\nabla^2 f = \text{div}(\text{grad } f) = \text{div}(\text{grad}(\ln(x^2 + y^2))) =$
 $\text{div}\left(\left[\frac{2x}{x^2+y^2}, \frac{2y}{x^2+y^2}, 0\right]\right) = \frac{4}{x^2+y^2} - \frac{4y^2}{(x^2+y^2)^2} - \frac{4x^2}{(x^2+y^2)^2} = 0$ ■

Definition: Curl of a Vector Field

Curl of a vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$ is defined as


$$\text{curl } \mathbf{v} = \nabla \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$

Definition: Curl of a Vector Field

Curl of a vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$ is defined as

$$\text{curl } \mathbf{v} = \nabla \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$




Definition: Curl of a Vector Field

Curl of a vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$ is defined as

$$\text{curl } \mathbf{v} = \nabla \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$




From a scalar field we can obtain a vector field by the gradient.

Definition: Curl of a Vector Field

Curl of a vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$ is defined as

$$\text{curl } \mathbf{v} = \nabla \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$




From a scalar field we can obtain a vector field by the gradient.
Conversely, from a vector field we can obtain a scalar field by the divergence

Definition: Curl of a Vector Field

Curl of a vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$ is defined as

$$\text{curl } \mathbf{v} = \nabla \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$



From a scalar field we can obtain a vector field by the gradient. Conversely, from a vector field we can obtain a scalar field by the divergence or yet another vector field by the curl.




Definition: Curl of a Vector Field

Curl of a vector function

$\mathbf{v}(x, y, z) = [v_1(x, y, z), v_2(x, y, z), v_3(x, y, z)]$ is defined as

$$\text{curl } \mathbf{v} = \nabla \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_1 & v_2 & v_3 \end{vmatrix}$$



From a scalar field we can obtain a vector field by the gradient. Conversely, from a vector field we can obtain a scalar field by the divergence or yet another vector field by the curl.



We also have:

- ① **Gradient fields are irrotational.** That is, if a continuously differentiable vector function is the gradient of a scalar function, then its curl is the zero vector,

$$\text{curl}(\text{grad } f) = \mathbf{0}$$

We also have:

- 1 **Gradient fields are irrotational.** That is, if a continuously differentiable vector function is the gradient of a scalar function, then its curl is the zero vector,

$$\text{curl}(\text{grad } f) = \mathbf{0}$$

- 2 Divergence of the curl of a twice continuously differentiable vector function \mathbf{v} is zero,

$$\text{div}(\text{curl } f) = 0.$$


Question: (10ed-9.9-5) Find $\text{curl } \mathbf{v}$ for $\mathbf{v} = xyz [x^2, y^2, z^2]$?

Question: (10ed-9.9-5) Find $\text{curl } \mathbf{v}$ for $\mathbf{v} = xyz [x^2, y^2, z^2]^T$?

Solution: $\text{curl } \mathbf{v} = \text{curl} (xyz [x^2, y^2, z^2]^T)$

$$= \text{curl} ([x^3yz, xy^3z, xyz^3]) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^3yz & xy^3z & xyz^3 \end{vmatrix} =$$
$$(xz^3 - xy^3) \mathbf{i} + (x^3y - yz^3) \mathbf{j} + (y^3z - x^3z) \mathbf{k}$$

The vector field and its curl are sketched below:



Vector field $[x^3yz, xy^3z, xyz^3]$ and its Curl



Question: (10ed-9.9-10) Let $\mathbf{v} = [\sec x, \csc x, 0]$ be the velocity vector of a steady fluid flow. Is the flow irrotational? Incompressible? Find the streamlines (the paths of the particles).

Question: (10ed-9.9-10) Let $\mathbf{v} = [\sec x, \csc x, 0]$ be the velocity vector of a steady fluid flow. Is the flow irrotational? Incompressible? Find the streamlines (the paths of the particles).

Solution: $\text{curl}([\sec x, \csc x, 0]) = -\frac{\cos x}{\sin^2 x} \mathbf{k} \neq \mathbf{0} \Rightarrow$ Not irrotational.

Question: (10ed-9.9-10) Let $\mathbf{v} = [\sec x, \csc x, 0]$ be the velocity vector of a steady fluid flow. Is the flow irrotational? Incompressible? Find the streamlines (the paths of the particles).

Solution: $\text{curl}([\sec x, \csc x, 0]) = -\frac{\cos x}{\sin^2 x} \mathbf{k} \neq \mathbf{0} \Rightarrow$ Not irrotational.

$\text{div}([\sec x, \csc x, 0]) = \frac{1}{\cos^2 x} \sin x \neq 0 \Rightarrow$ Compressible. ■

Question: (10ed-9.9-10) Let $\mathbf{v} = [\sec x, \csc x, 0]$ be the velocity vector of a steady fluid flow. Is the flow irrotational? Incompressible? Find the streamlines (the paths of the particles).

Solution: $\text{curl}([\sec x, \csc x, 0]) = -\frac{\cos x}{\sin^2 x} \mathbf{k} \neq \mathbf{0} \Rightarrow$ Not irrotational.

$\text{div}([\sec x, \csc x, 0]) = \frac{1}{\cos^2 x} \sin x \neq 0 \Rightarrow$ Compressible. ■

Question: (10ed-9.9-11) Let $\mathbf{v} = [y, -2x, 0]$ be the velocity vector of a steady fluid flow. Is the flow irrotational? Incompressible? Find the streamlines (the paths of the particles)?

Question: (10ed-9.9-10) Let $\mathbf{v} = [\sec x, \csc x, 0]$ be the velocity vector of a steady fluid flow. Is the flow irrotational? Incompressible? Find the streamlines (the paths of the particles).

Solution: $\text{curl}([\sec x, \csc x, 0]) = -\frac{\cos x}{\sin^2 x} \mathbf{k} \neq \mathbf{0} \Rightarrow$ Not irrotational.

$\text{div}([\sec x, \csc x, 0]) = \frac{1}{\cos^2 x} \sin x \neq 0 \Rightarrow$ Compressible. ■

Question: (10ed-9.9-11) Let $\mathbf{v} = [y, -2x, 0]$ be the velocity vector of a steady fluid flow. Is the flow irrotational? Incompressible? Find the streamlines (the paths of the particles)?

Solution: $\text{curl}(\mathbf{v}) = \text{curl}([y, -2x, 0]) = -3\mathbf{k} \neq \mathbf{0} \Rightarrow$ Not irrotational.

$\text{div}([y, -2x, 0]) = 0 \Rightarrow$ incompressible. ■