

Advanced Engineering Mathematics

Mathematical Techniques for Engineering

Dr Athar Kharal

atharkharal.github.io

atharkharal@gmail.com

<https://pk.linkedin.com/in/atharkharal>

Dirac's Delta Function

Definition

Consider a function $f_k(t - a) = \begin{cases} \frac{1}{k} & , \quad a \leq t \leq a + k \\ 0 & , \quad \text{otherwise} \end{cases}$,

Dirac's Delta Function

Definition

Consider a function $f_k(t - a) = \begin{cases} \frac{1}{k} & , \quad a \leq t \leq a + k \\ 0 & , \quad \text{otherwise} \end{cases}$, which has unit area

Dirac's Delta Function

Definition

Consider a function $f_k(t - a) = \begin{cases} \frac{1}{k} & , \quad a \leq t \leq a + k \\ 0 & , \quad \text{otherwise} \end{cases}$, which has unit area as


$$\int_0^{\infty} f_k(t - a) dt = \int_a^{a+k} \frac{1}{k} dt = 1.$$

Dirac's Delta Function

Definition

Consider a function $f_k(t - a) = \begin{cases} \frac{1}{k} & , \quad a \leq t \leq a + k \\ 0 & , \quad \text{otherwise} \end{cases}$, which has unit area as

$$\int_0^{\infty} f_k(t - a) dt = \int_a^{a+k} \frac{1}{k} dt = 1.$$




Dirac's Delta Function

Definition

Consider a function $f_k(t - a) = \begin{cases} \frac{1}{k} & , \quad a \leq t \leq a + k \\ 0 & , \quad \text{otherwise} \end{cases}$, which has unit area as

$$\int_0^{\infty} f_k(t - a) dt = \int_a^{a+k} \frac{1}{k} dt = 1.$$



Then Dirac's delta function (aka, unit impulse function) is defined as

$$\delta(t - a) = \lim_{k \rightarrow 0} f_k(t - a).$$

Paul Adrien Maurice Dirac

Dirac (8 August **1902** – 20 October **1984**) was an English theoretical physicist

Paul Adrien Maurice Dirac

Dirac (8 August **1902** – 20 October **1984**) was an English theoretical physicist who made fundamental contributions to the early development of both quantum mechanics and quantum electrodynamics.

Paul Adrien Maurice Dirac

Dirac (8 August **1902** – 20 October **1984**) was an English theoretical physicist who made fundamental contributions to the early development of both quantum mechanics and quantum electrodynamics. Among other discoveries, he formulated the **Dirac equation**, which describes the behaviour of fermions and **predicted the existence of antimatter**.

Paul Adrien Maurice Dirac

Dirac (8 August **1902** – 20 October **1984**) was an English theoretical physicist who made fundamental contributions to the early development of both quantum mechanics and quantum electrodynamics. Among other discoveries, he formulated the **Dirac equation**, which describes the behaviour of fermions and **predicted the existence of antimatter**. Dirac shared the **Nobel Prize** in Physics for 1933 **with Erwin Schrödinger**, "for the discovery of new productive forms of atomic theory"

Paul Adrien Maurice Dirac

Dirac (8 August **1902** – 20 October **1984**) was an English theoretical physicist who made fundamental contributions to the early development of both quantum mechanics and quantum electrodynamics. Among other discoveries, he formulated the **Dirac equation**, which describes the behaviour of fermions and **predicted the existence of antimatter**. Dirac shared the **Nobel Prize** in Physics for 1933 **with Erwin Schrödinger**, "for the discovery of new productive forms of atomic theory"

He also did work that forms the basis of modern attempts to reconcile general relativity with quantum mechanics resulting into **Superstrings Theory** and lastly, the **Baby Universes of Stephen Hawking**.

Paul Adrien Maurice Dirac

Dirac (8 August **1902** – 20 October **1984**) was an English theoretical physicist who made fundamental contributions to the early development of both quantum mechanics and quantum electrodynamics. Among other discoveries, he formulated the **Dirac equation**, which describes the behaviour of fermions and **predicted the existence of antimatter**. Dirac shared the **Nobel Prize** in Physics for 1933 **with Erwin Schrödinger**, "for the discovery of new productive forms of atomic theory"

He also did work that forms the basis of modern attempts to reconcile general relativity with quantum mechanics resulting into **Superstrings Theory** and lastly, the **Baby Universes of Stephen Hawking**.



LT of Dirac's Delta Function

LT of Dirac's Delta function may be obtained by writing it in Heaviside terms as

$$\delta(t-a) = \frac{1}{k} [u(t-a) - u(t-(a+k))]$$

LT of Dirac's Delta Function

LT of Dirac's Delta function may be obtained by writing it in Heaviside terms as

$$\delta(t-a) = \frac{1}{k} [u(t-a) - u(t-(a+k))]$$

$$L(f_k(t-a)) =$$

LT of Dirac's Delta Function

LT of Dirac's Delta function may be obtained by writing it in Heaviside terms as

$$\begin{aligned}\delta(t-a) &= \frac{1}{k} [u(t-a) - u(t-(a+k))] \\ L(f_k(t-a)) &= \frac{1}{ks} [e^{-as} - e^{-(a+k)s}] = e^{-as} \frac{1 - e^{-ks}}{ks}\end{aligned}$$

LT of Dirac's Delta Function

LT of Dirac's Delta function may be obtained by writing it in Heaviside terms as

$$\delta(t-a) = \frac{1}{k} [u(t-a) - u(t-(a+k))]$$

$$L(f_k(t-a)) = \frac{1}{ks} [e^{-as} - e^{-(a+k)s}] = e^{-as} \frac{1 - e^{-ks}}{ks}$$

$$L\left(\lim_{k \rightarrow 0} f_k(t-a)\right) = \lim_{k \rightarrow 0} e^{-as} \frac{1 - e^{-ks}}{ks}$$

LT of Dirac's Delta Function

LT of Dirac's Delta function may be obtained by writing it in Heaviside terms as

$$\delta(t-a) = \frac{1}{k} [u(t-a) - u(t-(a+k))]$$

$$L(f_k(t-a)) = \frac{1}{ks} [e^{-as} - e^{-(a+k)s}] = e^{-as} \frac{1 - e^{-ks}}{ks}$$

$$\begin{aligned} L\left(\lim_{k \rightarrow 0} f_k(t-a)\right) &= \lim_{k \rightarrow 0} e^{-as} \frac{1 - e^{-ks}}{ks} \\ &= e^{-as} \lim_{k \rightarrow 0} \frac{\frac{d}{dk}(1 - e^{-ks})}{\frac{d}{dk}(ks)} \quad (\text{l'Hopital}) \end{aligned}$$


$$= e^{-as} \left[\frac{se^{-ks}}{s} \right]_{k=0} = e^{-as}$$

$$L(\delta(t-a)) = e^{-as} \quad \blacksquare$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if


$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?




Solution: Since for this circuit

$$Li' + Ri + \frac{q}{C} = 1i' + 20i + 10000 q = \delta(t),$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?




Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = 1i' + 20i + 10000 q = \delta(t)$, but the question is about voltage,

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?




Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = 1i' + 20i + 10000 q = \delta(t)$, but the question is about voltage, hence the equation may be re-written as

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Solution: Since for this circuit


$Li' + Ri + \frac{q}{C} = \delta(t)$, but the question is about voltage, hence the equation may be re-written as follows:

$$\frac{d^2}{dt^2} q(t) + 20 \frac{d}{dt} q(t) + 10000 q(t) = \delta(t) ; \quad q(0) = q'(0) = 0$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = \delta(t)$, but the question is about voltage, hence the equation may be re-written as follows:


$$\frac{d^2}{dt^2} q(t) + 20 \frac{d}{dt} q(t) + 10000 q(t) = \delta(t) ; \quad q(0) = q'(0) = 0$$

$$s^2 Q - q'(0) - sq(0) + 20sQ - 20q(0) + 10000Q = 1$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = \delta(t)$, but the question is about voltage, hence the equation may be re-written as follows:

$$\frac{d^2}{dt^2} q(t) + 20 \frac{d}{dt} q(t) + 10000 q(t) = \delta(t) ; \quad q(0) = q'(0) = 0$$


$$s^2 Q - q'(0) - sq(0) + 20sQ - 20q(0) + 10000Q = 1$$

$$Q = \frac{1 + q'(0) + sq(0) + 20q(0)}{s^2 + 20s + 10000} = \frac{1}{s^2 + 20s + 10000} = \frac{1}{(s+10)^2 + 9900}$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = \delta(t)$, but the question is about voltage, hence the equation may be re-written as follows:

$$\frac{d^2}{dt^2} q(t) + 20 \frac{d}{dt} q(t) + 10000 q(t) = \delta(t) ; \quad q(0) = q'(0) = 0$$

$$s^2 Q - q'(0) - s q(0) + 20 s Q - 20 q(0) + 10000 Q = 1$$


$$Q = \frac{1 + q'(0) + s q(0) + 20 q(0)}{s^2 + 20 s + 10000} = \frac{1}{s^2 + 20 s + 10000} = \frac{1}{(s+10)^2 + 9900}$$

$$\text{By inverse LT, } q(t) = \frac{\sqrt{11} e^{-10t} \sin(30 \sqrt{11} t)}{330}$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = \delta(t)$, but the question is about voltage, hence the equation may be re-written as follows:

$$\frac{d^2}{dt^2} q(t) + 20 \frac{d}{dt} q(t) + 10000 q(t) = \delta(t) ; \quad q(0) = q'(0) = 0$$

$$s^2 Q - q'(0) - s q(0) + 20 s Q - 20 q(0) + 10000 Q = 1$$

$$Q = \frac{1 + q'(0) + s q(0) + 20 q(0)}{s^2 + 20 s + 10000} = \frac{1}{s^2 + 20 s + 10000} = \frac{1}{(s+10)^2 + 9900}$$


By inverse LT, $q(t) = \frac{\sqrt{11} e^{-10t} \sin(30 \sqrt{11} t)}{330}$

$$v(t)$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = \delta(t)$, but the question is about voltage, hence the equation may be re-written as follows:

$$\frac{d^2}{dt^2} q(t) + 20 \frac{d}{dt} q(t) + 10000 q(t) = \delta(t) ; \quad q(0) = q'(0) = 0$$

$$s^2 Q - q'(0) - s q(0) + 20 s Q - 20 q(0) + 10000 Q = 1$$

$$Q = \frac{1 + q'(0) + s q(0) + 20 q(0)}{s^2 + 20 s + 10000} = \frac{1}{s^2 + 20 s + 10000} = \frac{1}{(s+10)^2 + 9900}$$


$$\text{By inverse LT, } q(t) = \frac{\sqrt{11} e^{-10t} \sin(30 \sqrt{11} t)}{330}$$

$$v(t) = \frac{q(t)}{C} \Big|_{C=10^{-4}}$$

Question:

Find output voltage response of the four-terminal RLC circuit given in figure if

$R = 20\Omega$, $L = 1H$, $C = 10^{-4}F$, the input is an impulse, current and charge are zero at time $t = 0$?



Solution: Since for this circuit

$Li' + Ri + \frac{q}{C} = \delta(t)$, but the question is about voltage, hence the equation may be re-written as follows:

$$\frac{d^2}{dt^2} q(t) + 20 \frac{d}{dt} q(t) + 10000 q(t) = \delta(t) ; \quad q(0) = q'(0) = 0$$

$$s^2 Q - q'(0) - s q(0) + 20 s Q - 20 q(0) + 10000 Q = 1$$

$$Q = \frac{1 + q'(0) + s q(0) + 20 q(0)}{s^2 + 20 s + 10000} = \frac{1}{s^2 + 20 s + 10000} = \frac{1}{(s+10)^2 + 9900}$$

By inverse LT, $q(t) = \frac{\sqrt{11} e^{-10t} \sin(30 \sqrt{11} t)}{330}$

$$v(t) = \left. \frac{q(t)}{C} \right|_{C=10^{-4}} = \frac{1000}{33} \sqrt{11} e^{-10t} \sin(30 \sqrt{11} t)$$

Question: (10e-6.4-10)

$$y'' + 5y' + 6y = \delta\left(t - \frac{1}{2}\pi\right) + u(t - \pi) \cos t \quad ; \quad y(0) = y'(0) = 0$$

Question: (10e-6.4-10)

$$y'' + 5y' + 6y = \delta\left(t - \frac{1}{2}\pi\right) + u(t - \pi) \cos t \quad ; \quad y(0) = y'(0) = 0$$

$$s^2 Y - y'(0) - sy(0) + 5sY - 5y(0) + 6Y = e^{-\frac{\pi}{2}s} - \frac{s}{s^2+1}e^{-s\pi}$$

Using ICs and isolating Y , we have

$$Y = \frac{1}{s^2+5s+6} \left(e^{-\frac{\pi}{2}s} - \frac{s}{s^2+1}e^{-s\pi} \right)$$

Question: (10e-6.4-10)

$$y'' + 5y' + 6y = \delta\left(t - \frac{1}{2}\pi\right) + u(t - \pi) \cos t \quad ; \quad y(0) = y'(0) = 0$$

$$s^2 Y - y'(0) - sy(0) + 5sY - 5y(0) + 6Y = e^{-\frac{\pi}{2}s} - \frac{s}{s^2+1}e^{-s\pi}$$

Using ICs and isolating Y , we have

$$Y = \frac{1}{s^2+5s+6} \left(e^{-\frac{\pi}{2}s} - \frac{s}{s^2+1} e^{-s\pi} \right)$$

$$Y = \left(\frac{1}{(s+3)(s+2)} \right) e^{-\frac{\pi}{2}s} - \left(\frac{s}{(s+3)(s+2)(s^2+1)} \right) e^{-s\pi}$$

Question: (10e-6.4-10)

$$y'' + 5y' + 6y = \delta\left(t - \frac{1}{2}\pi\right) + u(t - \pi) \cos t \quad ; \quad y(0) = y'(0) = 0$$

$$s^2 Y - y(0) - sy'(0) + 5sY - 5y(0) + 6Y = e^{-\frac{\pi}{2}s} - \frac{s}{s^2+1} e^{-s\pi}$$

Using ICs and isolating Y , we have

$$Y = \frac{1}{s^2+5s+6} \left(e^{-\frac{\pi}{2}s} - \frac{s}{s^2+1} e^{-s\pi} \right)$$

$$Y = \left(\frac{1}{(s+3)(s+2)} \right) e^{-\frac{\pi}{2}s} - \left(\frac{s}{(s+3)(s+2)(s^2+1)} \right) e^{-s\pi}$$

Using Partial Fractions

$$Y = \left(\frac{1}{(s+2)} - \frac{1}{(s+3)} \right) e^{-\frac{\pi}{2}s} - \left(\frac{-2}{5} \frac{1}{(s+2)} + \frac{1}{10} \frac{s+1}{s^2+1} + \frac{3}{10} \frac{1}{(s+3)} \right) e^{-s\pi}$$

$$y(t) = L^{-1} \left(\frac{e^{-\frac{\pi}{2}s}}{s+2} \right) - L^{-1} \left(\frac{e^{-\frac{\pi}{2}s}}{s+3} \right) + \frac{2}{5} L^{-1} \left(\frac{1}{s+2} e^{-s\pi} \right) - \frac{1}{10} L^{-1} \left(\frac{s}{1+s^2} e^{-s\pi} \right) - \frac{1}{10} L^{-1} \left(\frac{1}{1+s^2} e^{-s\pi} \right) - \frac{3}{10} L^{-1} \left(\frac{1}{s+3} e^{-s\pi} \right)$$

$$y(t) = L^{-1} \left(\frac{e^{-\frac{\pi}{2}s}}{s+2} \right) - L^{-1} \left(\frac{e^{-\frac{\pi}{2}s}}{s+3} \right) + \frac{2}{5} L^{-1} \left(\frac{1}{s+2} e^{-s\pi} \right) - \frac{1}{10} L^{-1} \left(\frac{s}{1+s^2} e^{-s\pi} \right) - \frac{1}{10} L^{-1} \left(\frac{1}{1+s^2} e^{-s\pi} \right) - \frac{3}{10} L^{-1} \left(\frac{1}{s+3} e^{-s\pi} \right)$$

$$y(t) = e^{\pi-2t} u\left(t - \frac{1}{2}\pi\right) - e^{\frac{3}{2}\pi-3t} u\left(t - \frac{1}{2}\pi\right) + \frac{2}{5} e^{2\pi-2t} u(t - \pi) - \frac{1}{10} (-\cos t) u(t - \pi) - \frac{1}{10} (-\sin t) u(t - \pi) - \frac{3}{10} e^{3\pi-3t} u(t - \pi)$$

$$y(t) = \frac{1}{10} u(t - \pi) (-3e^{-3t+3\pi} + \sin(t) + \cos(t) + 4e^{-2t+2\pi}) + u(t - \pi/2) (-e^{-3t+3/2\pi} + e^{-2t+\pi}) \quad \blacksquare$$

Question: (10ed-6.4-12)

$$\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t) + 5y(t) = 25t - 100\delta(t - \pi), \quad y(0) = -2, \quad y'(0) = 5$$

Question: (10ed-6.4-12)

$$\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t) + 5y(t) = 25t - 100\delta(t - \pi), \quad y(0) = -2, \quad y'(0) = 5$$

$$s^2 Y - y'(0) - sy(0) + 2sY - 2y(0) + 5Y = 25\frac{1}{s^2} - 100e^{-s\pi}$$

Question: (10ed-6.4-12)

$$\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t) + 5y(t) = 25t - 100\delta(t - \pi), \quad y(0) = -2, \quad y'(0) = 5$$

$$s^2 Y - y'(0) - sy(0) + 2sY - 2y(0) + 5Y = 25\frac{1}{s^2} - 100e^{-s\pi}$$

$$Y = \frac{1}{s^2 + 2s + 5} \left(25\frac{1}{s^2} - 100e^{-s\pi} + y'(0) + sy(0) + 2y(0) \right)$$

$$Y = \frac{1}{s^2 + 2s + 5} \left(25\frac{1}{s^2} - 100e^{-s\pi} + 1 - 2s \right)$$

$$Y = -\frac{100e^{-s\pi}s^2 + 2s^3 - s^2 - 25}{s^2(s^2 + 2s + 5)}$$

Question: (10ed-6.4-12)

$$\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t) + 5y(t) = 25t - 100\delta(t - \pi), \quad y(0) = -2, \quad y'(0) = 5$$

$$s^2 Y - y'(0) - sy(0) + 2sY - 2y(0) + 5Y = 25\frac{1}{s^2} - 100e^{-s\pi}$$

$$Y = \frac{1}{s^2 + 2s + 5} \left(25\frac{1}{s^2} - 100e^{-s\pi} + y'(0) + sy(0) + 2y(0) \right)$$

$$Y = \frac{1}{s^2 + 2s + 5} \left(25\frac{1}{s^2} - 100e^{-s\pi} + 1 - 2s \right)$$

$$Y = -\frac{100e^{-s\pi}s^2 + 2s^3 - s^2 - 25}{s^2(s^2 + 2s + 5)}$$

$$Y = -100\frac{1}{s^2 + 2s + 5}e^{-s\pi} - 2\frac{s}{s^2 + 2s + 5} + \frac{1}{s^2 + 2s + 5} + \frac{25}{s^2(s^2 + 2s + 5)}$$

$$y(t) = -50u(t - \pi)e^{-t+\pi}\sin(2t) + 5t - 2$$

Question: (10ed-6.4-12)

$$\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t) + 5y(t) = 25t - 100\delta(t - \pi), \quad y(0) = -2, \quad y'(0) = 5$$

$$s^2 Y - y'(0) - sy(0) + 2sY - 2y(0) + 5Y = 25\frac{1}{s^2} - 100e^{-s\pi}$$

$$Y = \frac{1}{s^2 + 2s + 5} (25\frac{1}{s^2} - 100e^{-s\pi} + y'(0) + sy(0) + 2y(0))$$

$$Y = \frac{1}{s^2 + 2s + 5} (25\frac{1}{s^2} - 100e^{-s\pi} + 1 - 2s)$$

$$Y = -\frac{100e^{-s\pi}s^2 + 2s^3 - s^2 - 25}{s^2(s^2 + 2s + 5)}$$

$$Y = -100\frac{1}{s^2 + 2s + 5}e^{-s\pi} - 2\frac{s}{s^2 + 2s + 5} + \frac{1}{s^2 + 2s + 5} + \frac{25}{s^2(s^2 + 2s + 5)}$$

$$y(t) = -50u(t - \pi)e^{-t+\pi}\sin(2t) + 5t - 2$$

$$y(t) = \begin{cases} 5t - 2 & , \quad t < \pi \\ 5t - 2 - 50e^{\pi-t}\sin(2t) & , \quad t > \pi \end{cases}$$



Convolution: Product of Transforms

For two functions f and g , it must be noted that
 $L(f)L(g) \neq L(fg)$,

Convolution: Product of Transforms

For two functions f and g , it must be noted that $L(f)L(g) \neq L(fg)$, instead we have $L(f)L(g) = L(f \star g)$,

Convolution: Product of Transforms

For two functions f and g , it must be noted that $L(f)L(g) \neq L(fg)$, instead we have $L(f)L(g) = L(f \star g)$, where $f \star g$ is defined as

$$(f \star g)(t) = \int_0^t f(\tau) g(t - \tau) d\tau$$

Convolution: Product of Transforms

For two functions f and g , it must be noted that $L(f)L(g) \neq L(fg)$, instead we have $L(f)L(g) = L(f \star g)$, where $f \star g$ is defined as

$$(f \star g)(t) = \int_0^t f(\tau) g(t - \tau) d\tau$$

In words one says: product of transforms is not the transform of product, **instead**

Convolution: Product of Transforms

For two functions f and g , it must be noted that $L(f)L(g) \neq L(fg)$, instead we have $L(f)L(g) = L(f \star g)$, where $f \star g$ is defined as

$$(f \star g)(t) = \int_0^t f(\tau) g(t - \tau) d\tau$$

In words one says: product of transforms is not the transform of product, **instead** product of transforms is the transform of convolution.

Question: Find $L^{-1} \left(\frac{1}{(s^2 + \omega^2)^2} \right)$?

Question: Find $L^{-1} \left(\frac{1}{(s^2 + \omega^2)^2} \right)$?

Solution: We note that given entity seems a product of two familiar LT

Question: Find $L^{-1} \left(\frac{1}{(s^2 + \omega^2)^2} \right)$?

Solution: We note that given entity seems a product of two familiar LT i.e. $\frac{1}{(s^2 + \omega^2)^2} = \frac{1}{(s^2 + \omega^2)} \times \frac{1}{(s^2 + \omega^2)}$.

Question: Find $L^{-1} \left(\frac{1}{(s^2 + \omega^2)^2} \right)$?

Solution: We note that given entity seems a product of two familiar LT i.e. $\frac{1}{(s^2 + \omega^2)^2} = \frac{1}{(s^2 + \omega^2)} \times \frac{1}{(s^2 + \omega^2)}$. We also recall

$$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \frac{\sin(\omega t)}{\omega}.$$

Question: Find $L^{-1} \left(\frac{1}{(s^2 + \omega^2)^2} \right)$?

Solution: We note that given entity seems a product of two familiar LT i.e. $\frac{1}{(s^2 + \omega^2)^2} = \frac{1}{(s^2 + \omega^2)} \times \frac{1}{(s^2 + \omega^2)}$. We also recall

$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \frac{\sin(\omega t)}{\omega}$. Hence we write

$$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \frac{\sin(\omega t)}{\omega} \star \frac{\sin(\omega t)}{\omega}$$

Question: Find $L^{-1} \left(\frac{1}{(s^2 + \omega^2)^2} \right)$?

Solution: We note that given entity seems a product of two familiar LT i.e. $\frac{1}{(s^2 + \omega^2)^2} = \frac{1}{(s^2 + \omega^2)} \times \frac{1}{(s^2 + \omega^2)}$. We also recall

$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \frac{\sin(\omega t)}{\omega}$. Hence we write

$$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \frac{\sin(\omega t)}{\omega} \star \frac{\sin(\omega t)}{\omega}$$

$$\text{since } (f \star g)(t) = \int_0^t f(\tau) g(t - \tau) d\tau$$

$$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \int_0^t \frac{\sin(\omega \tau)}{\omega} \frac{\sin(\omega(t - \tau))}{\omega} d\tau$$

Question: Find $L^{-1} \left(\frac{1}{(s^2 + \omega^2)^2} \right)$?

Solution: We note that given entity seems a product of two familiar LT i.e. $\frac{1}{(s^2 + \omega^2)^2} = \frac{1}{(s^2 + \omega^2)} \times \frac{1}{(s^2 + \omega^2)}$. We also recall

$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \frac{\sin(\omega t)}{\omega}$. Hence we write

$$L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) = \frac{\sin(\omega t)}{\omega} \star \frac{\sin(\omega t)}{\omega}$$

$$\text{since } (f \star g)(t) = \int_0^t f(\tau) g(t - \tau) d\tau$$

$$\begin{aligned} L^{-1} \left(\frac{1}{s^2 + \omega^2} \right) &= \int_0^t \frac{\sin(\omega \tau)}{\omega} \frac{\sin(\omega(t - \tau))}{\omega} d\tau \\ &= \frac{1}{\omega^2} \int_0^t \frac{1}{2} [-\cos(\omega t) + \cos(2\omega \tau - \omega t)] d\tau \\ &= \frac{1}{2\omega^2} \left[-\tau \cos \omega t + \frac{\sin(\omega \tau)}{\omega} \right]_{\tau=0}^t \\ &= \frac{1}{2\omega^2} \left[-t \cos \omega t + \frac{\sin(\omega t)}{\omega} \right] \end{aligned}$$

Properties of Convolution:

$$f \star g = g \star f$$

$$f \star (g + h) = f \star g + f \star h$$

$$f \star (g \star h) = (f \star g) \star h$$

$$f \star 0 = 0 \star f = 0$$

$$f \star 1 \neq f \quad (\text{why?})$$

Question: (10ed-6.5-6) Find convolution $e^{at} \star e^{bt}$, ($a \neq b$)?

Question: (10ed-6.5-6) Find convolution $e^{at} \star e^{bt}$, ($a \neq b$)?

Solution: Since we have $(f \star g)(t) = \int_0^t f(\tau) g(t - \tau) d\tau$, we take $f(t) = e^{at}$ and $g(t) = e^{bt}$:

$$\int_0^t e^{a\tau} e^{b(t-\tau)} d\tau = \int_0^t e^{bt-b\tau} e^{a\tau} d\tau$$

Question: (10ed-6.5-6) Find convolution $e^{at} \star e^{bt}$, ($a \neq b$)?

Solution: Since we have $(f \star g)(t) = \int_0^t f(\tau) g(t - \tau) d\tau$, we take $f(t) = e^{at}$ and $g(t) = e^{bt}$:

$$\int_0^t e^{a\tau} e^{b(t-\tau)} d\tau = \int_0^t e^{bt-b\tau} e^{a\tau} d\tau$$

by rewriting $e^{a\tau} e^{b(t-\tau)} = e^{a\tau \ln(e)} e^{b(t-\tau) \ln(e)}$

Question: (10ed-6.5-6) Find convolution $e^{at} \star e^{bt}$, ($a \neq b$)?

Solution: Since we have $(f \star g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau$, we take $f(t) = e^{at}$ and $g(t) = e^{bt}$:

$$\int_0^t e^{a\tau} e^{b(t-\tau)} d\tau = \int_0^t e^{bt-b\tau} e^{a\tau} d\tau$$

by rewriting $e^{a\tau} e^{b(t-\tau)} = e^{a\tau \ln(e)} e^{b(t-\tau) \ln(e)}$

$$= \int_0^t e^{a\tau \ln(e)} e^{b(t-\tau) \ln(e)} d\tau = \int_0^t e^{a\tau \ln(e) - b(\tau-t) \ln(e)} d\tau$$

Question: (10ed-6.5-6) Find convolution $e^{at} \star e^{bt}$, ($a \neq b$)?

Solution: Since we have $(f \star g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau$, we take $f(t) = e^{at}$ and $g(t) = e^{bt}$:

$$\int_0^t e^{a\tau} e^{b(t-\tau)} d\tau = \int_0^t e^{bt-b\tau} e^{a\tau} d\tau$$

by rewriting $e^{a\tau} e^{b(t-\tau)} = e^{a\tau \ln(e)} e^{b(t-\tau) \ln(e)}$

$$= \int_0^t e^{a\tau \ln(e)} e^{b(t-\tau) \ln(e)} d\tau = \int_0^t e^{a\tau \ln(e) - b(\tau-t) \ln(e)} d\tau$$

Substituting $u = a\tau \ln(e) - b(\tau-t) \ln(e)$ implies at

$\tau = 0$, $u = bt$ and $\tau = t$, $u = at$ and

$$du = \frac{d}{d\tau} (a\tau \ln(e) - b(\tau-t) \ln(e)) = a - b \Rightarrow \frac{du}{a-b} = d\tau$$

Question: (10ed-6.5-6) Find convolution $e^{at} \star e^{bt}$, ($a \neq b$)?

Solution: Since we have $(f \star g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau$, we take $f(t) = e^{at}$ and $g(t) = e^{bt}$:

$$\int_0^t e^{a\tau} e^{b(t-\tau)} d\tau = \int_0^t e^{bt-b\tau} e^{a\tau} d\tau$$

by rewriting $e^{a\tau} e^{b(t-\tau)} = e^{a\tau \ln(e)} e^{b(t-\tau) \ln(e)}$

$$= \int_0^t e^{a\tau \ln(e)} e^{b(t-\tau) \ln(e)} d\tau = \int_0^t e^{a\tau \ln(e) - b(\tau-t) \ln(e)} d\tau$$

Substituting $u = a\tau \ln(e) - b(\tau-t) \ln(e)$ implies at

$\tau = 0$, $u = bt$ and $\tau = t$, $u = at$ and

$$du = \frac{d}{d\tau} (a\tau \ln(e) - b(\tau-t) \ln(e)) = a - b \Rightarrow \frac{du}{a-b} = d\tau$$

$$= \int_{\ln(e)bt}^{at \ln(e)} \frac{e^u}{(a-b) \ln(e)} du = \frac{e^{at \ln(e)} - e^{\ln(e)bt}}{(a-b) \ln(e)} = \frac{e^{at} - e^{bt}}{(a-b)} \quad \blacksquare$$

Question: (10ed-6.5-13)

$$y(t) + 2e^t \int_0^t y(\tau) e^{-\tau} d\tau = te^t$$

Question: (10ed-6.5-13)

$$y(t) + 2e^t \int_0^t y(\tau) e^{-\tau} d\tau = te^t$$

$$y(t) + 2 \int_0^t y(\tau) e^{t-\tau} d\tau = te^t$$

Question: (10ed-6.5-13)

$$y(t) + 2e^t \int_0^t y(\tau) e^{-\tau} d\tau = te^t$$

$$y(t) + 2 \int_0^t y(\tau) e^{t-\tau} d\tau = te^t$$

$$y(t) + 2(y(t) \star e^t) = te^t$$

Question: (10ed-6.5-13)

$$y(t) + 2e^t \int_0^t y(\tau) e^{-\tau} d\tau = te^t$$

$$y(t) + 2 \int_0^t y(\tau) e^{t-\tau} d\tau = te^t$$

$$y(t) + 2(y(t) \star e^t) = te^t$$

$$Y + 2\left(Y \frac{1}{s-1}\right) = \frac{1}{(s-1)^2}$$

Question: (10ed-6.5-13)

$$y(t) + 2e^t \int_0^t y(\tau) e^{-\tau} d\tau = te^t$$

$$y(t) + 2 \int_0^t y(\tau) e^{t-\tau} d\tau = te^t$$

$$y(t) + 2(y(t) \star e^t) = te^t$$

$$Y + 2 \left(Y \frac{1}{s-1} \right) = \frac{1}{(s-1)^2}$$

$$Y = \frac{1}{s^2 - 1}$$

$$y(t) = L^{-1} \left(\frac{1}{s^2 - 1} \right)$$

$$y(t) = \sinh t$$



Question: (10ed-6.5-14)

$$y(t) - \int_0^t y(\tau)(t-\tau) d\tau = 2 - \frac{1}{2}t^2$$

Question: (10ed-6.5-14)

$$y(t) - \int_0^t y(\tau)(t-\tau) d\tau = 2 - \frac{1}{2}t^2$$

$$y(t) - (y(t) \star t) = 2 - \frac{1}{2}t^2$$

$$Y - \left(Y \frac{1}{s^2} \right) = \frac{2}{s} - \frac{1}{s^3}$$

Question: (10ed-6.5-14)

$$y(t) - \int_0^t y(\tau)(t-\tau) d\tau = 2 - \frac{1}{2}t^2$$

$$y(t) - (y(t) \star t) = 2 - \frac{1}{2}t^2$$

$$Y - \left(Y \frac{1}{s^2} \right) = \frac{2}{s} - \frac{1}{s^3}$$

$$Y = -\frac{2s^2 - 1}{s - s^3}$$

$$Y = \frac{1}{2(s-1)} + \frac{1}{2(s+1)} + \frac{1}{s}$$

$$y(t) = \frac{1}{2}e^t + \frac{1}{2}e^{-t} + 1$$

$$y(t) = 1 + \cosh t \quad \blacksquare$$

Question: (10ed-6.5-22)

$$\frac{e^{-as}}{s(s-2)} =$$

Question: (10ed-6.5-22)

$$\frac{e^{-as}}{s(s-2)} = \frac{1}{s-2} \frac{e^{-as}}{s} = L(e^{2t}) L(u(t-a)) = (e^{2t}) \star u(t-a)$$

Question: (10ed-6.5-22)

$$\frac{e^{-as}}{s(s-2)} = \frac{1}{s-2} \frac{e^{-as}}{s} = L(e^{2t}) L(u(t-a)) = (e^{2t}) \star u(t-a)$$

choose $f(t) = e^{2t}$ and $g(t) = u(t-a)$ in convolution integral

$$(f \star g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau$$

Question: (10ed-6.5-22)

$$\frac{e^{-as}}{s(s-2)} = \frac{1}{s-2} \frac{e^{-as}}{s} = L(e^{2t}) L(u(t-a)) = (e^{2t}) \star u(t-a)$$

choose $f(t) = e^{2t}$ and $g(t) = u(t-a)$ in convolution integral

$$(f \star g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau$$

$$(e^{2t}) \star u(t-a) = \int_0^t e^{2\tau} u((t-\tau)-a) d\tau$$

Question: (10ed-6.5-22)

$$\frac{e^{-as}}{s(s-2)} = \frac{1}{s-2} \frac{e^{-as}}{s} = L(e^{2t}) L(u(t-a)) = (e^{2t}) \star u(t-a)$$

choose $f(t) = e^{2t}$ and $g(t) = u(t-a)$ in convolution integral

$$(f \star g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau$$

$$(e^{2t}) \star u(t-a) = \int_0^t e^{2\tau} u((t-\tau)-a) d\tau$$

$$\begin{aligned} &= \left[\frac{1}{2} u(t-a-\tau) (e^{2\tau} + e^{2t-2a}) \right]_{\tau=0}^t \\ &= \left[\frac{1}{2} u(t-a-\tau) (e^{2\tau} + e^{2t-2a}) \right]_{\tau=t} - \\ &\quad \left[\frac{1}{2} u(t-a-\tau) (e^{2\tau} + e^{2t-2a}) \right]_{\tau=0} \end{aligned}$$

$$= \frac{1}{2} u(-a) (e^{2t-2a} + e^{2t}) - \frac{1}{2} u(t-a) (e^{2t-2a} + 1) \quad \blacksquare$$

Question: (9ed-6.5-20) Using convolution theorem solve the following IVP

$$y'' + 5y' + 4y = 2e^{-2t} \quad ; \quad y(0) = 0, y'(0) = 0$$

Question: (9ed-6.5-20) Using convolution theorem solve the following IVP

$$y'' + 5y' + 4y = 2e^{-2t} \quad ; \quad y(0) = 0, \quad y'(0) = 0$$

$$s^2 Y - y'(0) - sy(0) + 5sY - 5y(0) + 4Y = \frac{2}{s+2}$$

$$Y = \frac{1}{s^2 + 5s + 4} \left(\frac{2}{s+2} + y'(0) + sy(0) + 5y(0) \right) =$$
$$2 \frac{1}{(s+2)} \left(\frac{1}{(s+4)(s+1)} \right)$$

Question: (9ed-6.5-20) Using convolution theorem solve the following IVP

$$y'' + 5y' + 4y = 2e^{-2t} \quad ; \quad y(0) = 0, \quad y'(0) = 0$$

$$s^2 Y - y'(0) - sy(0) + 5sY - 5y(0) + 4Y = \frac{2}{s+2}$$

$$Y = \frac{1}{s^2 + 5s + 4} \left(\frac{2}{s+2} + y'(0) + sy(0) + 5y(0) \right) =$$
$$2 \frac{1}{(s+2)} \left(\frac{1}{(s+4)(s+1)} \right)$$

$$Y = \frac{2}{(s+2)} [L(e^{-4t}) L(e^{-t})]$$

Question: (9ed-6.5-20) Using convolution theorem solve the following IVP

$$y'' + 5y' + 4y = 2e^{-2t} \quad ; \quad y(0) = 0, \quad y'(0) = 0$$

$$s^2 Y - y'(0) - sy(0) + 5sY - 5y(0) + 4Y = \frac{2}{s+2}$$

$$Y = \frac{1}{s^2 + 5s + 4} \left(\frac{2}{s+2} + y'(0) + sy(0) + 5y(0) \right) =$$
$$2 \frac{1}{(s+2)} \left(\frac{1}{(s+4)(s+1)} \right)$$

$$Y = \frac{2}{(s+2)} [L(e^{-4t}) L(e^{-t})] = 2L(e^{-2t}) [L(e^{-4t}) L(e^{-t})]$$

Question: (9ed-6.5-20) Using convolution theorem solve the following IVP

$$y'' + 5y' + 4y = 2e^{-2t} \quad ; \quad y(0) = 0, \quad y'(0) = 0$$

$$s^2 Y - y'(0) - sy(0) + 5sY - 5y(0) + 4Y = \frac{2}{s+2}$$

$$Y = \frac{1}{s^2 + 5s + 4} \left(\frac{2}{s+2} + y'(0) + sy(0) + 5y(0) \right) =$$
$$2 \frac{1}{(s+2)} \left(\frac{1}{(s+4)(s+1)} \right)$$

$$Y = \frac{2}{(s+2)} [L(e^{-4t}) L(e^{-t})] = 2L(e^{-2t}) [L(e^{-4t}) L(e^{-t})]$$

$$Y = 2e^{-2t} \star [e^{-4t} \star e^{-t}] = 2e^{-2t} \star \int_0^t e^{-4\tau} e^{-(t-\tau)} d\tau$$

Question: (9ed-6.5-20) Using convolution theorem solve the following IVP

$$y'' + 5y' + 4y = 2e^{-2t} \quad ; \quad y(0) = 0, \quad y'(0) = 0$$

$$s^2 Y - y'(0) - sy(0) + 5sY - 5y(0) + 4Y = \frac{2}{s+2}$$

$$Y = \frac{1}{s^2 + 5s + 4} \left(\frac{2}{s+2} + y'(0) + sy(0) + 5y(0) \right) =$$
$$2 \frac{1}{(s+2)} \left(\frac{1}{(s+4)(s+1)} \right)$$

$$Y = \frac{2}{(s+2)} [L(e^{-4t}) L(e^{-t})] = 2L(e^{-2t}) [L(e^{-4t}) L(e^{-t})]$$

$$Y = 2e^{-2t} \star [e^{-4t} \star e^{-t}] = 2e^{-2t} \star \int_0^t e^{-4\tau} e^{-(t-\tau)} d\tau$$

$$Y = 2e^{-2t} \star \left[-\frac{1}{3} e^{-t-3\tau} \right]_{\tau=0}^{\tau=t}$$

$$= 2e^{-2t} \star \left(\frac{1}{3} e^{-t} - \frac{1}{3} e^{-4t} \right) = \frac{2}{3} e^{-2t} \star (e^{-t} - e^{-4t})$$

$$Y = \frac{2}{3} \int_0^t e^{-2\tau} \left(e^{-(t-\tau)} - e^{-4(t-\tau)} \right) d\tau$$

$$Y = \frac{2}{3} \int_0^t e^{-2\tau} \left(e^{-(t-\tau)} - e^{-4(t-\tau)} \right) d\tau$$

$$y(t) = \frac{2}{3} \left(\frac{1}{2} e^{-t} (e^{-t} - 1)^2 (e^{-t} + 2) \right)$$

$$y(t) = \frac{1}{3} e^{-t} (e^{-t} - 1)^2 (e^{-t} + 2) \quad \blacksquare$$