

Advanced Engineering Mathematics

Mathematical Techniques for Engineering

Dr Athar Kharal

Humanities and Science Department
College of Aeronautical Engineering
National University of Sciences and Tecnology (NUST), Pakistan

Differentiation and Integration of LT

Theorem

If $F(s)$ is the Laplace transform of $f(t)$, then

$$F'(s) = - \int_0^{\infty} e^{-st} t f(t) dt$$

from which, also note that

$$L(t f(t)) = -F'(s) \quad \text{and} \quad L^{-1}(F'(s)) = -t f(t)$$

Corollary

By mathematical induction we also have

$$L(t^n f(t)) = (-1)^n F^{(n)}(s)$$

Example

$$L^{-1} \left(\ln \left(1 + \frac{\omega^2}{s^2} \right) \right) = ?$$

We observe that $\ln \left(1 + \frac{\omega^2}{s^2} \right) = \ln \left(\frac{s^2 + \omega^2}{s^2} \right) = \ln (s^2 + \omega^2) - \ln s^2$.

This indicates that a derivative of \ln may bring terms like $\frac{s}{s^2 + \omega^2}$ or $\frac{1}{s}$. Hence we set $F(s) = \ln \left(\frac{s^2 + \omega^2}{s^2} \right)$ and proceed for an application of differentiation of LT theorem i.e.

$$F'(s) = \frac{d}{ds} \ln (s^2 + \omega^2) - \frac{d}{ds} \ln s^2 = 2 \frac{s}{s^2 + \omega^2} - \frac{2}{s}$$

$$L^{-1} (F'(s)) = L^{-1} \left(2 \frac{s}{s^2 + \omega^2} - \frac{2}{s} \right) = 2 \cos (\omega t) - 2 = 2 (\cos (\omega t) - 1)$$

Since $L^{-1} (F'(s)) = -tf(t)$

$$\Rightarrow -tf(t) = 2 (\cos (\omega t) - 1)$$

$$f(t) = \frac{2}{t} (1 - \cos (\omega t))$$



Example

(10e-6.6-2) $L(3t \sinh(4t)) = ?$

We know $L(tf(t)) = -F'(s)$

$$L(3t \sinh(4t)) = 3L(t \sinh(4t))$$

$$= 3 \left(-\frac{d}{ds} L(\sinh(4t)) \right)$$

$$= -3 \left(\frac{d}{ds} \left(\frac{4}{s^2 - 16} \right) \right)$$

$$= 24 \frac{s}{(s^2 - 16)^2}$$



Example

For finding $L(t^n e^{kt})$, we note

$$L(e^{kt}) = \frac{1}{s-k}, \quad (= F(s))$$

by corollary $L(t^n f(t)) = (-1)^n F^{(n)}(s)$, so we take n differentiation

$$\frac{d}{ds} \left(\frac{1}{s-k} \right) = \frac{-1}{(s-k)^2}, \quad \frac{d^2}{ds^2} \left(\frac{1}{s-k} \right) = \frac{2}{(s-k)^3}$$

$$\frac{d^3}{ds^3} \left(\frac{1}{s-k} \right) = \frac{-6}{(s-k)^4}, \quad \text{hence}$$

$$\frac{d^n}{ds^n} \left(\frac{1}{s-k} \right) = \frac{(-1)^n n!}{(s-k)^{n+1}}$$

$$L(t^n e^{kt}) = \frac{(-1)^n n!}{(s-k)^{n+1}}$$



Example

(10e-6.7-6)

$$\frac{d}{dt}y_1(t) = 5y_1(t) + y_2(t)$$

$$\frac{d}{dt}y_2(t) = y_1(t) + 5y_2(t)$$

$$y_1(0) = 1, y_2(0) = -3$$

Taking Laplace Transforms

$$sY_1 - y_1(0) = 5Y_1 + Y_2$$

$$sY_2 - y_2(0) = Y_1 + 5Y_2$$

Using ICs

$$sY_1 - 1 = 5Y_1 + Y_2$$

$$sY_2 + 3 = Y_1 + 5Y_2$$

Example

(10e-6.7-12)

$$\frac{d^2}{dt^2} y_1(t) = -2 y_1(t) + 2 y_2(t)$$

$$\frac{d^2}{dt^2} y_2(t) = 2 y_1(t) - 5 y_2(t)$$

$$y_1(0) = 1, y_2(0) = 3, y_1'(0) = 0, y_2'(0) = 0$$

Taking Laplace Transforms

$$s^2 Y_1 - y_1'(0) - s y_1(0) = -2 Y_1 + 2 Y_2$$

$$s^2 Y_2 - y_2'(0) - s y_2(0) = 2 Y_1 - 5 Y_2$$

Using ICs

$$s^2 Y_1 - s = -2 Y_1 + 2 Y_2$$

$$s^2 Y_2 - 3s = 2 Y_1 - 5 Y_2$$