

Marked circuits and fixed-five rooted interfaces in cubic graphs

Structural reductions and finite censuses

Atharva Vaidya

Research draft, 27 July 2026

Abstract

Let H be a finite simple cubic graph and let S be a four-edge matching. As algebraic setup, we first identify five Eulerian edge sets covering every edge twice with an everywhere-anisotropic flow in a four-dimensional quadratic space over $\mathbb{F}_2$, including the exact convention for loops. We then record six narrowly scoped results. First, if H has a Tait colouring in which the four marks have one common colour, and

$$|\delta_H(X)| + |S \cap E(H[X])| \geq 4$$

whenever $H[X]$ contains a cycle, then S is contained in an even edge set whose every circuit component contains an even number of marks. The proof combines elementary shore lemmas with three published prescribed-cycle and decomposition results. It does not require universal separation after the common-colour Tait colouring has been obtained.

Second, suppose H is cyclically 4-edge-connected, the marks are separated in a fixed common-colour Tait colouring, and a differently coloured edge $f \notin S$ is forbidden. If no cycle of $H - f$ contains all four marks, then f belongs to an independent cyclic four-edge cut. Hence a cycle through all four marks and avoiding f always exists when H is cyclically 5-edge-connected.

Third, for the surviving exceptional rooted three-pole interface in a separate $D_5 = \binom{[5]}{2}$ flow model, we prove a necessary cycle-support obstruction. If a nonbridge root has only one possible label after the connector triangle is fixed, every cycle through that root has a label support which spans all five coordinates and is either non-bipartite or the star $K_{1,4}$. An exact gluing factorization shows how this condition arises when a fixed-five boundary signature equals one of the exceptional type patterns in the literature. Missing root base pairs also force explicit odd proper-core cuts. These are necessary restrictions, not eliminations.

Fourth, two exact solvers show that every nonempty nonbridge rooted three-pole signature through order 17 contains one of three base pairs: 15 645 623 roots in 654 676 canonical cores are checked. Fifth, on the 31 cyclically 4-edge-connected bridgeless non-Tait cubic graphs of order 22, all 14 322 independent-edge deletion poles have the full ten-type fixed-five boundary signature.

Sixth, an exhaustive canonical computation finds no *Tait-colourable* connected simple cubic graph on 22 vertices having a universally separated four-edge matching. Two separately written enumeration methods agree on 7 174 735 colourable graphs and 95 360 112 Tait colourings modulo global colour permutation. This manuscript neither proves nor disproves the Five-Cycle Double Cover Conjecture.

AI-use disclosure. OpenAI Codex agents, under the direction of Atharva Vaidya, generated the initial arguments, wrote the census programs, executed and audited the computations, and drafted this manuscript. Other Codex agents independently rechecked selected proofs and enumerations; those are AI-agent checks, not independent human verification or peer review. All mathematical arguments used here are displayed for line-by-line human checking, and the computational claims are separated from them. The human author should independently verify the paper and establish priority before submission.

1 Introduction and exact scope

A binary cycle of a graph is an edge set with even degree at every vertex. In a cubic graph its nonempty connected components are cycles. The first part of this paper asks when four marked edges lie in a binary cycle whose components each contain an even number of marks. This marked parity question arose in a larger investigation of the Five-Cycle Double Cover Conjecture, but the results below do not resolve that conjecture.

For clarity, the standard conjecture asks for at most five Eulerian edge subsets whose incidence multiplicities cover each edge exactly twice [8, 6]. The orientable variant imposes additional directional conditions and is a different problem. Neither version is proved here. In particular, our unrooted theorem has a marked-cut hypothesis, our rooted theorem leaves a cyclic four-cut interface, and the census is restricted to one order, simple graphs, and Tait-colourable inputs. All D_5 boundary signatures below use one fixed five-coordinate universe. They are not the unbounded-colour relation of Máčajová, Mazzuoccolo, and Tabarelli; our finite computations do not classify colourings needing more than five coordinates.

The six contributions

Result	Content	Nature of evidence
Theorem 4.1	Four common-colour marks satisfying the marked cyclic-cut inequality admit an even-marked binary cycle.	Human proof using explicitly identified imported theorems.
Theorem 5.1	A failed forbidden-edge reduction is localized to an independent cyclic four-cut; cyclic connectivity five closes the rooted case.	Human proof; only the four-independent-edge cycle theorem is imported.
Theorem 6.4	A nonbridge singleton root in the normalized D_5 three-pole interface forces every root cycle to have a spanning translation blocker.	Human algebraic and graph-cut proof; a necessary obstruction only.
Theorem 6.10	Failure of three rooted base pairs forces an odd proper-core cut carrying one common coordinate.	Elementary human proof; a necessary certificate only.
Theorem 7.3	Every nonempty nonbridge rooted three-pole signature through order 17 contains a base pair.	Two exact solvers with matching per-shard transcript digests.
Theorem 7.4	Every order-22 independent-edge deletion in the cyclically four-connected non-Tait cap corpus has the full fixed-five signature.	Two full boundary classifiers with byte-identical transcripts.
Theorem 8.2	No Tait-colourable connected simple cubic graph of order 22 has a universally separated four-edge matching.	Exhaustive computation with two algorithmically different enumerators.

The human arguments appear to be useful deductions from known cycle and boundary-signature machinery. The rooted factorizations and finite frontiers appear new to the authors, but the priority search performed for this draft was targeted rather than exhaustive.

A scope correction discovered during audit

An earlier project note stated the computational result over all connected simple cubic graphs of order 22. The programs, correctly for the motivating application, search only graphs having at least one Tait colouring. This qualifier is logically essential. If a graph has no Tait colouring, the assertion “separated in every Tait colouring” is vacuously true under the literal universal quantifier. Moreover, every cubic graph on 22 vertices has a matching of size at least four: a maximal matching of size m has endpoints dominating at most $6m$ vertices, so $m \geq \lceil 22/6 \rceil = 4$. Theorem 8.2 therefore states exactly the nonvacuous result established by the programs.

2 Definitions and imported results

All graphs are finite. Unless stated otherwise, they are simple. For $X \subseteq V(G)$, let $\delta_G(X)$ be the set of edges with exactly one end in X , and let $G[X]$ be the induced subgraph. A *cycle* is a connected 2-regular subgraph. A *binary cycle* is an edge subset $Q \subseteq E(G)$ such that every vertex has even degree in Q .

A graph is *cyclically k -edge-connected* if it has no edge cut of size less than k with a cycle on both shores. A *Tait colouring* of a cubic graph is a proper edge-colouring with three colours a, b, c . Every two colour classes induce a spanning 2-regular subgraph, called a *bichromatic factor*; its components are even cycles.

An exact elliptic-flow reformulation

For later use, put

$$\Gamma = \left\{ x \in \mathbb{F}_2^5 : \sum_{i=0}^4 x_i = 0 \right\}, \quad q(x) = \sum_{0 \leq i < j \leq 4} x_i x_j. \quad (2.1)$$

If $x \in \Gamma$, then

$$q(x) = \binom{\text{wt}(x)}{2} \pmod{2}.$$

Thus $q(x) = 1$ exactly for the ten vectors of weight two, which we identify with

$$D_5 = \binom{\{0, 1, 2, 3, 4\}}{2}. \quad (2.2)$$

The polar form is

$$B(x, y) = q(x + y) + q(x) + q(y) = \sum_{i=0}^4 x_i y_i \quad (x, y \in \Gamma). \quad (2.3)$$

It is nondegenerate: a vector orthogonal to every $e_i + e_j$ has all coordinates equal, and the all-one vector is not in Γ . The five nonzero singular vectors are $\mathbf{1} + e_i$, $0 \leq i < 5$. They span Γ , with their sum as their only dependence. Hence every isometry permutes them faithfully. Coordinate permutations already realize all five permutations, so

$$O(\Gamma, q) \cong S_5. \quad (2.4)$$

Proposition 2.1 (Elliptic-flow equivalence). *Let G be a finite multigraph. Five Eulerian edge subsets $C_0, \dots, C_4$ cover each edge exactly twice if and only if there is a map $\phi : E(G) \rightarrow \Gamma$ such that*

$$q(\phi(e)) = 1 \quad (e \in E(G)), \quad \sum_{\substack{\text{edge-ends} \\ \text{incident with } v}} \phi(e) = 0 \quad (v \in V(G)). \quad (2.5)$$

A loop is counted by its two edge-ends in the second sum.

Proof. Given the five edge subsets, let coordinate i of $\phi(e)$ be one exactly when $e \in C_i$. Exact double coverage makes $\phi(e)$ a weight-two vector, hence an element of $q^{-1}(1)$. Coordinate i of the vertex equation is precisely the degree in C_i modulo two. Conversely, $q(\phi(e)) = 1$ forces $\phi(e)$ to have exactly two nonzero coordinates. Defining $C_i = \{e : \phi(e)_i = 1\}$ therefore gives exact double coverage, and the five coordinate equations give the five even-degree conditions. The two constructions are inverse. $\square$

Parallel edges are distinct edge variables. A loop contributes twice to each selected degree, so it cancels in the characteristic-two vertex equation; counting a loop only once in a SAT/XOR incidence row would be an incorrect encoding. Empty C_i are allowed, so this five-coordinate formulation means “at most five” in the standard sense [8, 6]. It imposes no orientation condition. The underlying two-subset encoding also appears as a *CDC-colouring* in Máčajová, Mazzuoccolo, and Tabarelli [3, Definition 3.1]. Proposition 2.1 is an exact elementary reformulation, not a resolution or a priority claim; priority for the quadratic packaging has not been established.

Definition 2.2 (Separation). Let S be a matching in a Tait-coloured cubic graph. The marks are *separated in that colouring* if every bichromatic factor cycle contains at most one member of S . If the graph is Tait-colourable, then S is *universally separated* if it is separated in every Tait colouring.

We use the following three published results. They are the only non-elementary imported ingredients in the proof of Theorem 4.1.

- (I1) Knappe and Pitz [2, Fact 5.4] prove $g(3) = 3$. In the form used here: if G is 3-edge-connected and $F \subseteq E(G)$ has at most three edges, then a circuit contains F if and only if F contains no odd edge cut of G .
- (I2) Aldred, Ellingham, Hemminger, and Holton [1, Theorem 3.3] prove that a free edge system of size at most four in a quasi 4-connected graph lies on a cycle. In particular, four independent edges lie on a cycle. They also note that cyclically 4-edge-connected cubic graphs are quasi 4-connected.
- (I3) Every 3-connected cubic graph is obtained by repeated vertex 3-sums of cyclically 4-edge-connected cubic graphs. This is Theorem 6.1 of Nedela, Seifrtova, and Skoviera [5]; see also Pastor’s component-tree theorem [7].

Here a circuit in (I1) is a connected closed trail. In a cubic graph its edge-induced degrees are positive and even, hence equal to two, so its edge set is a cycle.

Lemma 2.3 (Tait cut parity). *In a Tait-coloured cubic graph, the numbers of cut edges of colours a, b, c have the same parity. In particular, the three edges of a three-edge cut have three distinct colours, and a Tait-colourable cubic graph has no bridge.*

Proof. For a colour p , counting incidences of the p -coloured perfect matching with X gives

$$|X| \equiv |\delta_G(X) \cap E_p| \pmod{2}.$$

This holds for all three colours. If the cut has size three, all three counts are odd and sum to three, so each is one. If a single edge were a cut, its colour count would have parity different from the other two. $\square$

Lemma 2.4 (Precolouring separated marks). *Let H be Tait-colourable and let S be universally separated. Every map $S \rightarrow \{a, b, c\}$ extends to a Tait colouring. In particular, there is a Tait colouring in which all marks have one common colour.*

Proof. Start with any Tait colouring. If a mark e currently has colour p and is requested to have colour q , interchange p and q on the unique pq -factor cycle through e . Universal separation says that this cycle contains no other mark, so all previously fixed marks keep their colours. Process the marks one at a time. $\square$

3 Two elementary shore lemmas

The marked hypothesis used throughout the first theorem is

$$|\delta_H(X)| + |S \cap E(H[X])| \geq 4 \quad \text{whenever } H[X] \text{ contains a cycle.} \quad (\text{MC})$$

Lemma 3.1 (Two-mark shore). *Let $A \subsetneq V(H)$, put $P = H[A]$, and suppose that P is connected and subcubic with minimum degree two. Suppose $|\delta_H(A)| \in \{2, 3\}$, the boundary edges have distinct ends in A , and exactly two marks lie in $E(P)$. If (MC) holds for every cyclic $X \subseteq A$, then one cycle of P contains both marks.*

Proof. Delete the bridges of P and form the tree of bridge-components. If P has no bridge, it is 2-edge-connected. A 2-edge-connected subcubic graph of minimum degree two has no cutvertex: two components behind a cutvertex would each require at least two incident edges at that vertex, giving degree at least four. Thus P is 2-connected. Subdivide the two marked edges. Any two vertices of a 2-connected graph lie on a common cycle, and suppressing the subdivisions gives a cycle through both marks.

Now suppose the bridge-component tree is nontrivial. Every leaf component K is cyclic; a single isolated vertex would have only one incident bridge but degree at least two in P . Let $p(K)$ count the pole-boundary edges incident with K , and let $s(K)$ count its internal marks. Exactly one bridge leaves K , so (MC) gives

$$1 + p(K) + s(K) \geq 4.$$

If K does not contain both marks, then $s(K) \leq 1$, hence $p(K) \geq 2$. Two such leaves would consume at least four distinct pole-boundary edges, impossible because there are at most three. Therefore a leaf contains both marks. That bridge-component is 2-edge-connected, subcubic, and of minimum degree two, so the preceding argument supplies the desired cycle. $\square$

Lemma 3.2 (One-mark shore). *Under the setup of Lemma 3.1, suppose instead that $|\delta_H(A)| = 3$, its ends in A are u_1, u_2, u_3 , and exactly one mark e lies in $E(P)$. If (MC) holds on every cyclic $X \subseteq A$, then for every pair $i \neq j$ there is a u_i - u_j path in P containing e .*

Proof. If P had a bridge, a leaf K of its bridge-component tree would be cyclic. With $p(K)$ and $s(K) \in \{0, 1\}$ as above,

$$1 + p(K) + s(K) \geq 4.$$

An unmarked leaf consumes all three pole-boundary edges, while a marked leaf consumes at least two. Two leaves cannot do so using only three boundary edges. Hence P is 2-edge-connected and, by the subcubic cutvertex argument, 2-connected.

Fix $i \neq j$, add an auxiliary edge $u_i u_j$, and use the standard fact that any two edges of a 2-connected graph lie on a common cycle. The assertion remains valid if the auxiliary edge is parallel to an existing edge. A cycle containing the auxiliary edge and e , with the auxiliary edge deleted, is the required path. $\square$

Corollary 3.3 (Low-cut reductions). *If a graph satisfying (MC) has no binary cycle containing all four marks with an even number of marks in each component, then it has no cyclic two-edge cut. Moreover, an unmarked cyclic three-edge cut cannot split the marks $2 + 2$.*

Proof. For a cyclic two-cut, (MC) puts at least two internal marks on each shore. Since there are four marks in total, the cut is unmarked and each shore has exactly two. Lemma 3.1 gives one two-mark cycle on each shore, whose union is a valid binary cycle. The same argument applies to the two shores of an unmarked $2 + 2$ cyclic three-cut. $\square$

4 The four-mark core theorem

Theorem 4.1 (Four-mark core closure). *Let H be a connected simple cubic graph and let S be a four-edge matching. Assume:*

- (i) *H has a Tait colouring in which every member of S has one common colour c ; and*
- (ii) *the marked cyclic-cut inequality (MC) holds.*

Then there is a binary cycle $Q \subseteq E(H)$ containing S such that every nonempty component of Q contains an even number of marks.

Remark 4.2. Universal separation is not a hypothesis of Theorem 4.1. By Lemma 2.4, it is one sufficient way to obtain hypothesis (i). The proof below uses only the one fixed common-colour Tait colouring.

Proof. Suppose that no such Q exists. By Corollary 3.3, H has no cyclic two-edge cut.

Step 1: H is 3-connected. Lemma 2.3 excludes bridges. A cutvertex in a bridgeless cubic graph is impossible, because one component behind the cutvertex would receive only one of its three incident edges, making that edge a bridge.

Suppose $\{u, v\}$ is a two-vertex cut. Every component D of $H - \{u, v\}$ has at least three boundary edges; otherwise the cubic degree sum makes both shores of $\delta_H(D)$ cyclic, contrary to the absence of a cyclic cut of size at most two. There are at least two components and at most six boundary incidences at u, v . Hence $uv \notin E(H)$, there are exactly two components D_1, D_2 , and each has three boundary edges. Neither component sends all three edges to only one of u, v , since the other would then be a cutvertex. Relabel so that D_1 sends two edges to u and one to v . Then $\delta_H(D_1 \cup \{u\})$ has size two. Both shores are cyclic by the cubic degree count and simplicity, a contradiction. Thus H is 3-connected.

Step 2: eliminate marked cyclic three-cuts. Let $B = \delta_H(A)$ be a cyclic three-edge cut and write

$$r = |B \cap S|, \quad a = |S \cap E(H[A])|, \quad d = |S \cap E(H[V(H) \setminus A])|.$$

The inequality (MC) on the two shores gives $a, d \geq 1$, and

$$a + d + r = 4. \tag{4.1}$$

In particular, $r \leq 2$. Since a shared boundary end would expose a cyclic cut of size at most two, the three cut edges have distinct ends on each shore.

If $r = 2$, then $a = d = 1$. On each shore, Lemma 3.2 gives a path through its unique internal mark between the ends of the two marked cut edges. The two paths and those two cut edges form one cycle containing all four marks.

Suppose $r = 1$. Up to exchanging shores, $a = 1$ and $d = 2$. Cap the two-mark shore by adjoining a new cubic vertex z at its three boundary ends; call the resulting cubic graph $\tilde{D}$. Lemma 2.3 shows that the fixed Tait colouring caps properly. The cap is 3-edge-connected. Indeed, it has no bridge, and a two-edge cut in a cubic graph has two cyclic shores; the shore avoiding z would induce a cyclic two-cut of H .

Let F consist of the cap edge corresponding to the marked member of B and the two internal marks. These three edges are independent. They contain no odd cut. A one-edge cut is absent. If the three edges formed a three-edge cut, that cut could not be a vertex star because F is independent. For a nontrivial three-cut, take the cyclic shore avoiding z . Both internal marks would be boundary edges and there would be no internal mark, contradicting $3 + 0 \geq 4$ in (MC). Imported result (I1) therefore gives a cycle through F .

At z , that cycle uses the marked cap edge and one other cap edge. Deleting z leaves a path through both internal marks between the corresponding boundary positions. On the one-mark shore, Lemma 3.2 supplies a path through its mark for the same boundary pair. Gluing with the two selected cut edges gives one four-mark cycle. Both cases contradict the assumed failure. Hence every cyclic three-cut that remains is unmarked.

By Corollary 3.3, an unmarked cyclic three-cut cannot split the marks $2 + 2$. Equation (4.1) shows that every such cut splits them $1 + 3$.

Step 3: organize all remaining cuts at once. Apply imported result (I3). Split along cyclic three-cuts, cap each shore by a root vertex, and continue until the factors are cyclically 4-edge-connected. Reverse reconstruction gives a tree $\mathcal{T}$: its nodes are the final factors and its edges record the vertex 3-sums. Each tree edge corresponds to a principal cyclic three-cut of H .

All principal cuts are unmarked, so every marked edge belongs to one factor. Give a node weight equal to its number of marks. Deleting any tree edge splits the total weight four as $1 + 3$. There is a node K for which every component of $\mathcal{T} - K$ has weight one. To prove this, begin at any node and move into a component of weight three whenever one exists. The walk never backtracks and must stop. At the stopping node no component has weight three; weights two and zero are excluded respectively by the $1 + 3$ rule and by positivity on both sides of every tree edge.

Let K contain r marked edges. It has $q = 4 - r$ incident one-mark branches, represented in the capped factor by root vertices $z_1, \dots, z_q$. Transfer the fixed Tait colouring to the cap. The r central marks have colour c ; at each z_j , let f_j be the unique incident edge of colour c . Let

$$F = (S \cap E(K)) \cup \{f_1, \dots, f_q\},$$

interpreted as a set, so a shared edge between adjacent cap roots is included once. The set F is a matching, $1 \leq |F| \leq 4$, it contains every central mark, and an edge of F is incident with each z_j .

If $|F| = 4$, imported result (I2) supplies a cycle through F . If $|F| \leq 3$, then F contains no odd cut: a one-cut is excluded by 3-edge-connectivity, a trivial three-cut is not a matching, and a nontrivial three-cut is excluded by cyclic 4-edge-connectivity. Imported result (I1) again supplies a cycle through F . Denote the resulting central cycle by Z .

For every j , the cycle Z passes through z_j and selects two of its three boundary positions. The corresponding entire branch of $\mathcal{T} - K$ is a shore with exactly one mark. Lemma 3.2 gives a path through that mark for the selected boundary pair. Substitute these paths while reversing the 3-sums. Each substitution preserves a single cycle. If two adjacent cap roots share the selected edge $z_i z_j$, reversing both sums turns that single edge into the single boundary edge between the two expanded shores; it is not duplicated. The final cycle contains the r central marks and one mark from each of the q branches, hence all four marks. This final contradiction proves the theorem. $\square$

Remark 4.3 (Why the conclusion allows two components). An unmarked cyclic three-cut with a $2 + 2$ mark split can be closed by one two-mark cycle on each shore. Nothing in the hypotheses forces those two cycles to merge. Thus the binary-cycle conclusion is the proved statement; an unconditional one-cycle strengthening is not claimed.

5 A forbidden root edge and the four-cut gate

We now use a stronger fixed-colouring hypothesis and obtain a rooted conclusion in a more highly connected class.

Theorem 5.1 (Independent four-cut gate). *Let H be a finite simple cyclically 4-edge-connected cubic graph. Let $S = \{s_1, s_2, s_3, s_4\}$ be a matching and let $f \notin S$. Assume that H has a Tait colouring in which*

- (i) *all members of S have one common colour c ;*
- (ii) *f has a different colour; and*
- (iii) *S is separated in this fixed colouring.*

If no cycle of $H - f$ contains all four members of S , then f belongs to an independent cyclic four-edge cut of H .

Proof. Write $f = uv$,

$$N_H(u) - \{v\} = \{u_1, u_2\}, \quad N_H(v) - \{u\} = \{v_1, v_2\},$$

and form the root-edge reduction

$$R = H \ominus f := H - \{u, v\} + \{\bar{u} = u_1 u_2, \bar{v} = v_1 v_2\}. \quad (5.1)$$

The graph H has at least eight vertices because S is a four-edge matching. It has no triangle: the triangle boundary has size three and its complementary shore is cyclic by the cubic degree count, contrary to cyclic 4-edge-connectivity. Consequently the four neighbours appearing in (5.1) do not create loops, parallel edges, or coincident new edge objects. Also H is 3-connected (for cubic graphs, cyclic connectivity and ordinary connectivity agree through order three), so deleting the adjacent pair u, v leaves a connected graph. Thus R is connected, simple, and cubic.

Map every mark $s \in S$ to an edge $\bar{s} \in E(R)$ by

$$\bar{s} = \begin{cases} \bar{u}, & \text{if } s \text{ is incident with } u, \\ \bar{v}, & \text{if } s \text{ is incident with } v, \\ s, & \text{otherwise.} \end{cases} \quad (5.2)$$

The four images are distinct. We claim that they form a matching. Only adjacency between a new image and an unchanged image needs proof. Suppose, without loss of generality, that $s = uu_1$ is a mark of colour c , while f has colour a . Then uu_2 has the third colour b . An unchanged mark incident with u_2 would lie with s on the bc -factor cycle, contradicting fixed-colouring separation. A mark incident with u_1 would already be adjacent to s . The other cases are symmetric.

If R were cyclically 4-edge-connected, imported result (I2) would give a cycle through the four independent images. Replace $\bar{u}$, when used, by u_1uu_2 , and similarly replace $\bar{v}$. The lifted cycle lies in $H - f$ and contains every original mark, contrary to hypothesis. Hence R has a cyclic cut $\delta_R(X)$ of size $k \leq 3$.

We pull this cut back to H . If u_1, u_2 lie on opposite shores, then $\bar{u}$ is a cut edge; placing u on either shore replaces $\bar{u}$ by exactly one crossing edge. If u_1, u_2 lie on one shore, place u there and create no crossing edge. The same choices apply to v . Unless both neighbour pairs are unsplit and lie on opposite shores, choose u, v on the same shore. Then f is internal, the cut size remains $k \leq 3$, and both shore cycles lift: an internal new edge is replaced by its two-edge path, while a crossing new edge was not part of either shore cycle. This contradicts cyclic 4-edge-connectivity of H .

Therefore u_1, u_2 lie together on one shore and v_1, v_2 lie together on the other. Restoring u, v on those forced shores gives

$$\delta_H(X^\uparrow) = \delta_R(X) \cup \{f\}. \quad (5.3)$$

Both shores remain cyclic, so $k + 1 \geq 4$. Since $k \leq 3$, we have $k = 3$.

It remains to prove independence. If two edges of the cyclic three-cut $\delta_R(X)$ shared an endpoint w on one shore, then only one edge at w would remain internal. No cycle of that shore uses w . Moving w to the other shore replaces two crossing edges by one and preserves cycles on both shores, producing a cyclic two-cut of R . The pullback argument just given excludes such a cut, since it would produce a cyclic cut of size at most three in H . Thus the three edges of $\delta_R(X)$ are independent. In the forced configuration they avoid the new edges and hence avoid u, v . Together with f , they form an independent cyclic four-edge cut of H . $\square$

Corollary 5.2 (Rooted closure at cyclic connectivity five). *Under the hypotheses of Theorem 5.1, if H is cyclically 5-edge-connected, then $H - f$ has a cycle containing all four marks.*

Proof. Otherwise Theorem 5.1 produces a cyclic four-edge cut. $\square$

Remark 5.3 (What remains at cyclic connectivity four). Theorem 5.1 is a one-way obstruction localization, not a characterization: a root-containing independent four-cut need not prevent an avoiding cycle. The unresolved case is the corresponding four-pole interface. No unrestricted rooted theorem follows from the gate alone.

6 Cycle translations in an exceptional rooted interface

This section records a second, algebraic restriction on a rooted interface. Its “root” is a distinguished cap edge whose possible D_5 -labels are studied; it is not the forbidden edge of Theorem 5.1.

The conclusion is a necessary obstruction, not a proof that the exceptional interface is impossible. Throughout this section, $\mathcal{R}_5$ and $\mathcal{C}_5$ denote exact signatures inside one fixed five-coordinate D_5 universe. This is narrower than the published unbounded-colour relation.

6.1 The D_5 model

Recall Γ , q , and $D_5 = q^{-1}(1)$ from (2.1)–(2.2). We identify a member of D_5 with its weight-two incidence vector and write addition in Γ as symmetric difference. A D_5 -labelling of a cubic multipole assigns a member of D_5 to every proper edge and semiedge so that the three incident labels sum to zero at each internal vertex. Such a local triple is exactly the three-edge set of a triangle in K_5 .

Let Q be a cubic three-pole whose ordered connector word has been normalized to

$$\tau = (01, 02, 12), \quad (6.1)$$

and let r be a distinguished proper edge. Its exact root signature is

$$\mathcal{R}_5(Q, r) = \{A \in D_5 : Q \text{ has a } D_5\text{-labelling with word } \tau \text{ and root label } A\}. \quad (6.2)$$

The stabilizer of the ordered word τ fixes 0, 1, 2 and may swap 3, 4. Therefore $\mathcal{R}_5(Q, r)$ is a union of the seven orbits

$$\{01\}, \{02\}, \{12\}, \{03, 04\}, \{13, 14\}, \{23, 24\}, \{34\}. \quad (6.3)$$

Lemma 6.1 (Cycle translation). *Let λ be a normalized root labelling of Q , let C be a cycle of proper edges, and let $h \in \Gamma$. If*

$$\lambda(e) + h \in D_5 \quad (e \in E(C)), \quad (6.4)$$

then adding h to every label on C , and changing no other label, gives another normalized root labelling.

Proof. Every vertex of C is incident with exactly two translated edges, so the change in its incident-label sum is $h + h = 0$. All other vertices are unchanged. A proper cycle contains no connector semiedge, so the word τ in (6.1) remains fixed. Condition (6.4) says that all translated values remain in the allowed label set. $\square$

Lemma 6.2 (Translation blockers). *Let $L \subseteq E(K_5) = D_5$ be nonempty, with its edges connected under edge adjacency. The only $h \in \Gamma$ for which*

$$A + h \in D_5 \quad (A \in L) \quad (6.5)$$

is $h = 0$ if and only if the graph $(\{0, 1, 2, 3, 4\}, L)$

(i) spans all five coordinate vertices; and

(ii) is either non-bipartite or is the four-edge star $K_{1,4}$.

Proof. Every nonzero element of Γ has weight two or four. First let h have weight four, say $h = \{0, 1, 2, 3, 4\} \setminus \{v\}$. For $A \in D_5$, the symmetric difference $A + h$ has weight two exactly when A avoids v . Thus all weight-four translations are blocked exactly when L spans all five vertices.

Now let $h = B$ have weight two. For $A \in D_5$,

$$|A \triangle B| = 2 \iff |A \cap B| = 1.$$

Consequently B translates every member of L within D_5 exactly when every edge of L crosses the cut

$$B \mid (\{0, 1, 2, 3, 4\} \setminus B). \quad (6.6)$$

If L spans and is non-bipartite, it is contained in no such cut. If L is spanning and bipartite, edge-adjacency connectedness makes it connected, so its bipartition is unique. A $2 + 3$ bipartition gives a weight-two translation. A $1 + 4$ bipartition has no side of size two; connectedness and spanning force all four incident edges, namely $K_{1,4}$. The weight-two and weight-four cases prove the equivalence. $\square$

Remark 6.3. Up to coordinate permutation, the inclusion-minimal connected blockers are $K_{1,4}$, the bull graph, and C_5 . This follows directly: a minimal spanning non-bipartite graph contains a triangle or a five-cycle. Minimality gives C_5 in the latter case. Starting from a triangle, the unused vertices must attach at distinct triangle vertices: attaching both to one vertex contains a star blocker, while a pendant path leaves a triangle plus a disjoint edge after its joining edge is deleted. The remaining case is the bull.

Theorem 6.4 (Cycle-support obstruction). *Suppose*

$$\mathcal{R}_5(Q, r) = \{A\}, \quad A \in \{01, 02, 12, 34\},$$

and r is a nonbridge of the proper graph of Q . In every normalized root labelling and on every proper cycle C containing r , the set of distinct labels

$$L_C = \{\lambda(e) : e \in E(C)\}$$

spans all five coordinates and is either non-bipartite or $K_{1,4}$.

Proof. Consecutive cycle labels share one coordinate because they are two edges of the local label triangle. Thus L_C is nonempty and connected under edge adjacency. If it were not a blocker, Lemma 6.2 would give a nonzero h satisfying (6.4). By Lemma 6.1, translation on C gives another normalized labelling whose root label is $A + h \in D_5$. Since $h \neq 0$, this differs from A , contradicting the singleton signature. Lemma 6.2 gives the stated classification. $\square$

6.2 Exact factorization across the cyclic three-cut

We now state the precise setting in which the singleton arises. Let $G = P + \{e, f\}$ be a bridgeless cubic cap with no cyclic two-edge cut. Assume that a cyclic three-edge cut, disjoint from e, f , has connected shores, splits the four terminals of P as $2 + 2$, and contains one whole cap edge on each shore. Cutting the three cut edges gives rooted three-poles (Q_1, e) and (Q_2, f) . Order corresponding connectors, normalize them to τ , and write

$$R = \mathcal{R}_5(Q_1, e), \quad S = \mathcal{R}_5(Q_2, f).$$

For nonempty $R, S \subseteq D_5$, let

$$\text{rel}(R, S) \subseteq \{E, I, D\}$$

record which of equality, one-point intersection, and disjointness occur among cross-pairs $(A, B) \in R \times S$. Let π denote the pairing of the four boundary semiedges induced by the deleted cap edges. In the standard four-pole notation, a doubled boundary word (A, A, B, B) has type

$$AA, \quad AT_\pi, \quad \text{or} \quad T_\pi T_\pi$$

according as $A = B$, $|A \cap B| = 1$, or $A \cap B = \emptyset$.

Proposition 6.5 (Exact rooted factorization). *The doubled part of the fixed-five boundary signature $\mathcal{C}_5(P)$ is*

$$\begin{aligned} \mathcal{C}_5(P) \cap \{AA, AT_\pi, T_\pi T_\pi\} = & \{AA : E \in \text{rel}(R, S)\} \\ & \cup \{AT_\pi : I \in \text{rel}(R, S)\} \\ & \cup \{T_\pi T_\pi : D \in \text{rel}(R, S)\}. \end{aligned} \quad (6.7)$$

Proof. Given a labelling of P with doubled word (A, A, B, B) , restore e with label A and f with label B . Cutting the three-edge cut gives two rooted labellings with a common ordered connector triangle. One global coordinate permutation normalizes it to τ , placing the transformed root labels in R, S ; equality, intersection size, and disjointness are invariant under that permutation.

Conversely, take defining labellings for any $A \in R$ and $B \in S$. Their connector words are identically τ , so corresponding semiedges glue. Delete the two roots while retaining their labels on the four boundary semiedges. The resulting word is (A, A, B, B) , with the type determined by the cross-relation. Both inclusions follow. $\square$

Lemma 6.6 (Sparse cross-relations). *Let R, S be nonempty and invariant under the stabilizer of τ .*

(a) $\text{rel}(R, S) = \{E\}$ if and only if

$$R = S = \{A\} \quad \text{for some } A \in \{01, 02, 12, 34\}. \quad (6.8)$$

(b) $\text{rel}(R, S) = \{D\}$ if and only if, after possibly swapping R, S ,

$$\emptyset \neq R \subseteq \{01, 02, 12\}, \quad \emptyset \neq S \subseteq \{d \in D_5 : d \cap a = \emptyset \text{ for every } a \in R\}, \quad (6.9)$$

where S is stabilizer-invariant. There are thirteen unordered patterns in (6.9).

Proof. If every cross-pair is equal, choosing one member on either side forces both sets to be the same singleton. The singleton is stabilizer-invariant exactly for the four fixed labels in (6.8), proving (a).

For (b), every pair described in (6.9) is disjoint. Conversely, suppose every cross-pair is disjoint. If one side contains 34, the other side consists of triangle edges. If one side contains a mixed edge $u3$, invariance also puts $u4$ there, and the only edge disjoint from both is the triangle edge on $\{0, 1, 2\} \setminus \{u\}$. Thus, after swapping, one side is a nonempty subset of $\{01, 02, 12\}$, and the other is exactly restricted as in (6.9). For a one-edge triangle subset the common disjoint neighbourhood has two stabilizer orbits, giving three nonempty invariant choices; this contributes $3 \cdot 3 = 9$. A two-edge subset has neighbourhood $\{34\}$, contributing three, and the three-edge subset contributes one. Hence $9 + 3 + 1 = 13$. $\square$

Lemma 6.7 (Cap roots are nonbridges). *In the cap geometry preceding Proposition 6.5, neither e nor f is a bridge of its rooted shore.*

Proof. Suppose e is a bridge of Q_1 , splitting its proper graph into connected vertex sets X, Y . Let k of the three cut connectors be incident with X . If $k = 0$ or 3 , then e is a bridge of G , contrary to bridgelessness. Thus $k \in \{1, 2\}$. Choose $Z = X$ for $k = 1$, and $Z = Y$ for $k = 2$. Exactly two edges of G leave Z : e and one connector.

If Z has n vertices and m internal edges, cubic degree counting gives $3n = 2m + 2$, so

$$m - n + 1 = n/2 > 0.$$

Hence Z contains a cycle. Its complement is connected: the other bridge component of Q_1 connects through the two remaining connectors to the connected opposite shore Q_2 . Applying the same degree count to the complement gives a cycle there as well. The two leaving edges form a cyclic two-edge cut of G , a contradiction. The argument for f is symmetric. $\square$

Máčajová, Mazzuoccolo, and Tabarelli [3] isolate the following two exceptional type sets in their unbounded-colour signature relation:

$$\begin{aligned} \mathcal{E}_4(i; j, k) &= \{AA, AT_j, AT_k, T_j T_k\}, \\ \mathcal{E}_5(i; j, k) &= \{T_i T_i, T_j T_j, T_k T_k, T_i T_j, T_i T_k\}, \quad \{i, j, k\} = \{2, 3, 4\}. \end{aligned} \tag{6.10}$$

Below, the statements “ $\mathcal{C}_5(P) = \mathcal{E}_4$ ” and “ $\mathcal{C}_5(P) = \mathcal{E}_5$ ” use the same type masks as conditional targets in the fixed-five model. They do not identify $\mathcal{C}_5(P)$ with the full published relation. Intersecting these displayed sets with the three types in Proposition 6.5 gives the following literal consequence.

Corollary 6.8 (Exceptional rooted obligations). *(i) If $\mathcal{C}_5(P) = \mathcal{E}_4(i; j, k)$ and $\pi = i$, then $\text{rel}(R, S) = \{\mathbf{E}\}$. Thus $R = S = \{A\}$ for one fixed label $A \in \{01, 02, 12, 34\}$, both roots are nonbridges, and every cycle through either root has a support classified by Theorem 6.4.*

(ii) If $\mathcal{C}_5(P) = \mathcal{E}_4(i; j, k)$ and $\pi \in \{j, k\}$, then $\text{rel}(R, S) = \{\mathbf{E}, \mathbf{I}\}$.

(iii) If $\mathcal{C}_5(P) = \mathcal{E}_5(i; j, k)$, then the cross-relations are disjointness only:

$$\text{rel}(R, S) = \{\mathbf{D}\}.$$

Hence one of the thirteen patterns in Lemma 6.6(b) occurs.

Proof. For $\mathcal{E}_4$, the intersection with $\{AA, AT_\pi, T_\pi T_\pi\}$ is $\{AA\}$ when $\pi = i$, and $\{AA, AT_\pi\}$ when $\pi = j$ or k . For $\mathcal{E}_5$, it is $\{T_\pi T_\pi\}$ for every π . Proposition 6.5, Lemmas 6.6 and 6.7, and Theorem 6.4 give the assertions. $\square$

6.3 Missing base pairs force proper-core cuts

For a normalized labelling ϕ and a coordinate a , let

$$E_{\bar{a}}(\phi) = \{e \text{ a proper edge of } Q : a \notin \phi(e)\}, \quad h_a = \{0, 1, 2, 3, 4\} \setminus \{a\}.$$

Lemma 6.9 (Complement-cycle switch). *Suppose $a \notin \phi(r)$. If r lies on a proper cycle $C \subseteq E_{\bar{a}}(\phi)$, then replacing every label $\phi(e)$, $e \in C$, by $\phi(e) \triangle h_a$ gives another normalized labelling. Consequently, if $\phi(r) \triangle h_a \notin \mathcal{R}_5(Q, r)$, then r is a bridge of $E_{\bar{a}}(\phi)$.*

Proof. Every label on C is a two-subset of the four-set h_a ; symmetric difference with h_a is its complementary two-subset and therefore remains in D_5 . At a cycle vertex the same vector h_a is added twice, so the vertex equation is unchanged. No boundary semiedge is changed. The last assertion uses the elementary fact that an edge of a finite graph is a nonbridge exactly when it lies on a cycle. $\square$

Define the three base pairs

$$\mathcal{P}_0 = \{12, 03, 04\}, \quad \mathcal{P}_1 = \{02, 13, 14\}, \quad \mathcal{P}_2 = \{01, 23, 24\}.$$

Theorem 6.10 (Base-pair cut certificate). *Suppose $\mathcal{R}_5(Q, r)$ is nonempty, contains a label other than 34, and contains none of $\mathcal{P}_0, \mathcal{P}_1, \mathcal{P}_2$ as a subset. Then some realizing labelling ϕ and $a \in \{3, 4\}$ make r a bridge of $E_{\bar{a}}(\phi)$.*

More precisely, an inner root label 12, 02, or 01 forces this bridge for both $a = 3$ and $a = 4$; an outer orbit without its inner mate supplies one bridge for each coordinate after applying the symmetry $3 \leftrightarrow 4$.

Proof. The fixed boundary word $(01, 02, 12)$ is invariant under $3 \leftrightarrow 4$, so the exact root signature is invariant as well. Complementation in the four-set avoiding 4 pairs

$$12 \leftrightarrow 03, \quad 02 \leftrightarrow 13, \quad 01 \leftrightarrow 23,$$

and complementation in the four-set avoiding 3 gives the same three pairs with 3, 4 interchanged. If an inner label occurs, failure of its base pair and signature invariance exclude both outer mates. If an outer orbit occurs, failure of its base pair excludes the inner mate. Lemma 6.9 gives the claimed bridges. $\square$

Here is the promised ordinary cut certificate. Let $\delta_Q^p(X)$ denote the edge cut in the proper core of Q , explicitly excluding its three boundary semiedges. If X is one component of $E_{\bar{a}}(\phi) - r$, then

$$\delta_Q^p(X) = \{r\} \dot{\cup} F_a, \quad F_a \subseteq \{e : a \in \phi(e)\}.$$

For $a = 3, 4$, no boundary label uses a . The coordinate- a support on proper edges is therefore Eulerian, so $|F_a|$ is even. If r is a nonbridge of the unlabelled proper core, then $F_a \neq \emptyset$. Thus Theorem 6.10 supplies such a nonbridge root with an odd proper-core cut of size at least three, containing r , whose other edges carry one common outer coordinate.

For completeness, if $\mathcal{R}_5(Q, r) = \{34\}$, complementation in the four-sets avoiding 0, 1, 2 shows that r is simultaneously a bridge of $E_{\bar{0}}(\phi)$, $E_{\bar{1}}(\phi)$, and $E_{\bar{2}}(\phi)$. Boundary semiedges use these coordinates, so the even- $|F_a|$ conclusion is not asserted in this case.

Remark 6.11 (Exact limitation). Corollary 6.8 neither realizes nor excludes any of the remaining root signatures. In particular, Theorem 6.4 does not prove that a nonbridge root has two possible labels: it only classifies what every root cycle must look like if the label remains unique. The cut certificate above is likewise necessary, not contradictory. The $\{\mathbf{E}, \mathbf{I}\}$ and thirteen disjoint cases receive no further elimination here.

7 Fixed-five rooted finite frontiers

7.1 A human-checkable cap reduction

Let P be a connected simple proper core with four specified terminals $T = \{t_1, t_2, t_3, t_4\}$, each of proper degree two, and all other vertices of proper degree three. Attaching one semiedge at

each terminal gives a terminal-distinct cubic four-pole. A perfect matching of T is *simple for P* if neither of its pairs is already an edge of P .

Lemma 7.1 (Simple cap matching). *At least one of the three perfect matchings of T is simple for P .*

Proof. Suppose each perfect matching contains an edge of $P[T]$, and choose one such edge from each of the three matching classes of K_4 . The three chosen edges form either a three-edge star or a triangle. A star would give its terminal centre three proper neighbours, although that terminal has degree two. In the triangle case, all three triangle vertices have both proper incident edges inside the triangle, making it a component of P disjoint from the fourth terminal. Both possibilities contradict the setup. $\square$

The non-Tait consequence used next retains the fixed-five scope. Máčajová, Mazzuoccolo, and Tabarelli [3, Lemma 3.8] prove that a Tait-colourable pole has no isolated vertex in its signature graph. The T_i vertex is isolated in $\mathcal{E}_4$, and A is isolated in $\mathcal{E}_5$. Moreover, the cited construction uses at most four colours, so the same exclusion holds when the exact signature is restricted to one fixed five-set.

Theorem 7.2 (Simple-cap enumeration reduction). *Suppose no proper edge of P is a bridge and $\mathcal{C}_5(P)$ is one of the fixed-five target masks $\mathcal{E}_4, \mathcal{E}_5$ in (6.10). Some simple matching on T adds two independent edges e, f to produce a connected simple bridgeless cubic graph $G = P + \{e, f\}$ which is not Tait-colourable. Deleting e, f recovers P .*

Proof. Choose the matching by Lemma 7.1. Its pairs are nonedges with four distinct endpoints, so adding them preserves simplicity and makes every terminal cubic. The graph remains connected. Every old edge lies on a cycle because the proper core has no bridge. Each new edge lies on a cycle because its endpoints are joined by a path in connected P . Thus G is bridgeless.

If G had a Tait colouring, deleting e, f and retaining their colours on the four resulting semiedges would Tait-colour P , contrary to the fixed-five consequence of the cited lemma. $\square$

7.2 Root base pairs through order 17

A rooted three-pole core of order n here is a connected simple graph with exactly three degree-two terminals, all other vertices cubic, and one distinguished proper edge. Its three boundary semiedges are ordered and normalized to (01, 02, 12).

Theorem 7.3 (Finite rooted base-pair theorem). *For every odd $n \leq 17$, every nonempty fixed-five root signature of a nonbridge root in such an order- n core contains at least one of $\mathcal{P}_0, \mathcal{P}_1, \mathcal{P}_2$.*

This is a computer-assisted finite theorem. Canonical input was generated by

$$\text{geng -cq -d2 -D3 } n \text{ } m:m, \quad m = (3n - 3)/2,$$

which forces exactly three degree-two vertices. At order 17, eight canonical shards contain 654 676 cores and 15 645 623 nonbridge roots. Two separately written engines were used: a direct finite-domain propagator and an incremental SAT encoding with a one-hot ten-label domain and the complete local support table. Each reports

empty signatures	337 059,
signatures containing a base pair	15 308 564,
violations	0,
solver calls	52 216 251.

Their deterministic transcript digests agree separately on all eight shards. The retained verifier checks generator status, row accounting, both implementation summaries, and every digest equality. Lower odd orders are included in the archived report.

Theorem 7.3 is not an induction and supplies no universal root-signature theorem. Combined with Theorem 6.10, it says that the ordinary cut certificate does not occur among nonempty nonbridge roots in this finite simple-core range because a base pair is always present.

7.3 The cyclically four-connected order-22 cap corpus

Theorem 7.4 (Finite full-signature cap census). *Let G be a cyclically 4-edge-connected, bridgeless, non-Tait, connected simple cubic graph on 22 vertices, and let e, f be independent edges. Then the deletion pole $G - \{e, f\}$ has all ten parity-permitted boundary types in its fixed-five signature:*

$$\mathcal{C}_5(G - \{e, f\}) \text{ has mask } 0x3ff.$$

The quantifier is finite and exhaustive. Of 7 319 447 canonical connected simple cubic graphs on 22 vertices, 7 187 627 are bridgeless. Independent direct edge-colouring and perfect-matching filters agree on exactly 12 892 non-Tait graphs and on the hard corpus digest

230f2cd88e72011d73a40c5f3d7d5f9fb0fca6110de2ec2541581bc50982037c.

Exact cut classification leaves 31 cyclically 4-edge-connected graphs. Deleting every independent edge pair gives 14 322 ordered pole records.

Two classifiers ask ten questions for each record. One uses CaDiCaL; the other is independently written finite-domain code. Both return

$$\begin{aligned} \text{rows} &= 14\,322, \\ \text{queries} &= \text{SAT answers} = 143\,220, \\ \text{nonfull rows} &= 0. \end{aligned}$$

Their uncompressed output streams are byte-identical, with SHA-256

52edd8a189012acefd3565a79d5c58b6f209bbe40156560eb57e2c569fd7c091.

These checks prove Theorem 7.4 conditional on the generator and two implementations.

Corollary 7.5 (Exact finite cap consequence). *No bridge-free connected simple terminal-distinct order-22 four-pole with fixed-five signature $\mathcal{E}_4$ or $\mathcal{E}_5$ can have a cyclically 4-edge-connected simple two-edge cap.*

Proof. Theorem 7.2 places such a cap in the bridgeless non-Tait order-22 corpus. Theorem 7.4 says the recovered deletion pole has the full ten-type signature, not either exceptional proper subset. $\square$

This corollary does not cover a bridge-bearing or repeated-terminal pole, a nonsimple core, a cap with a cyclic two- or three-cut, higher orders, or the published arbitrary-colour signature relation. A larger all-cap order-22 run was still in progress when this draft was frozen and is not claimed here.

8 The order-22 computational census

8.1 The exact property

For a Tait-colourable cubic graph H , define a conflict graph $\Omega(H)$ whose vertices are the edge objects of H . Join two edge objects when they are adjacent in H , or when some Tait colouring places them on the same bichromatic factor cycle.

Lemma 8.1 (Conflict-graph equivalence). *A four-edge set S is a universally separated matching if and only if it is an independent set of size four in $\Omega(H)$.*

Proof. Adjacency conflicts are exactly the negation of the matching condition. Every other conflict records one colouring and one bichromatic cycle containing two selected edges. Absence of all such conflicts is exactly separation in every Tait colouring. $\square$

Theorem 8.2 (Order-22 census). *No Tait-colourable connected simple cubic graph on 22 vertices has a universally separated four-edge matching.*

8.2 Canonical generation

The program **geng** from nauty and Traces 2.9.3 [4] was run in four modular shards:

```
geng -cq -d3 -D3 22 0/4
geng -cq -d3 -D3 22 1/4
geng -cq -d3 -D3 22 2/4
geng -cq -d3 -D3 22 3/4
```

Here **-c** requires connectedness, **-d3 -D3** requires degree three, and **geng** canonically generates simple graphs. The four disjoint shards contain respectively

$$1\,376\,411, \quad 2\,078\,782, \quad 1\,716\,645, \quad 2\,147\,609$$

graphs, for a total of 7 319 447.

8.3 Two enumeration methods

The primary program fixes the three colours at one vertex, thereby quotienting the six global colour permutations. It recursively enumerates every remaining Tait colouring, records each bichromatic factor cycle, builds $\Omega(H)$, and performs exact backtracking for an independent four-set.

The replay program does not use the primary colouring recursion. It enumerates every perfect matching M , retains M exactly when its complement is a union of even cycles, and chooses independently one of the two alternating colourings on every complementary cycle. It then records the three bichromatic factor partitions and builds the same semantic conflict graph. Each Tait colouring modulo global colour permutation is visited six times: three choices of distinguished colour matching and two orders for the other two colours.

Both methods skip a graph only after establishing that it has no Tait colouring. On every shard they agree on the graph count, the number of Tait-colourable graphs, the number of normalized Tait colourings, and the zero witness count:

Shard	Graphs	Tait-colourable	Normalized colourings	Witnesses
0	1,376,411	1,338,944	17,481,016	0
1	2,078,782	2,029,518	28,040,733	0
2	1,716,645	1,686,315	22,708,101	0
3	2,147,609	2,119,958	27,130,262	0
Total	7,319,447	7,174,735	95,360,112	0

The replay additionally visits 312 583 931 perfect matchings. By Lemma 8.1, the exhaustive zero-witness output on all 7 174 735 eligible graphs proves Theorem 8.2, conditional on the correctness of the generator and implementations.

8.4 Controls and limitations

The two implementations agree on complete lower-order corpora:

Order	Graphs	Tait-colourable	Colourings	Perfect matchings	Witnesses
12	85	80	307	902	0
16	4,060	3,848	23,420	73,929	0
18	41,301	39,687	307,606	995,256	0

These three controls were rerun from the present source during the manuscript audit.

A positive target-three control at order 24 has graph6 encoding

W'??A???A_@_A_c0?S_Gc'_?@0@@?@00??_????_??H_A?E

Both programs enumerate 36 normalized Tait colourings, return the same first separated triple (edge identifiers 0, 1, 34), and reject a separated four-set. This exercises the positive witness path, although it is not a positive four-mark control.

The computation does not address multigraphs, orders 24 and above, or the rooted avoiding-cycle property. The two programs share the mathematical conflict-graph specification and the graph6 input format; their enumeration engines are different, but they are not formal proof-producing verifiers. Reproduction of the full census is the available check of the finite claim.

9 Reproducibility

The source distribution containing this manuscript stores the two programs at

```
scratch/tait_all_coloring_mark_separation.cpp
scratch/verify_order22_separation_via_matchings.cpp
```

and the machine-readable run record at

```
scratch/order22-universal-four-separation-result.json
```

Paths are relative to the project root, one directory above this preprint directory.

The finite truth-table checks supporting the rooted algebra, though not needed by its displayed proofs, are stored at

```

scratch/rooted_cycle_translation_blockers.py
scratch/verify_rooted_cycle_translation_blockers.mjs
scratch/rooted-cycle-translation-blockers-result.json
scratch/rooted_three_pole_signature_factorization.py
scratch/verify_rooted_three_pole_signature_factorization.mjs
scratch/rooted-three-pole-signature-factorization-result.json

```

The first replay checks all $2^{10} - 1$ nonempty K_5 edge supports. The second enumerates all nonempty pairs of stabilizer-orbit unions and confirms the four equality-only, 26 ordered disjoint-only, and thirteen unordered disjoint-only cases.

The rooted order-17 package and the completed cyclically-four order-22 cap slice are stored at

```

search/rooted-three-pole-frontier-20260727/
  order17-base-pair-report.json
  artifacts/order17-base-pair/
  verify_order17_base_pair_run.py
search/four-pole-order22-cap-20260727/
  artifacts/order22-cyclic4.g6
  artifacts/order22-cyclic4-poles.g6.gz
  artifacts/cyclic4-full-cadical.tsv.gz
  artifacts/cyclic4-full-csp.tsv.gz
  verify_completed_run.py

```

For the latter package, the verifier was run only through its cyclically-four extraction and full-signature checks. Its subsequent all-cap phase requires artifacts from the still-running larger census and is not part of this paper.

The semantic build and run commands are:

```

clang++ -O3 -std=c++20 \
  scratch/tait_all_coloring_mark_separation.cpp \
  -o /tmp/tait_all_coloring_mark_separation

geng -cq -d3 -D3 22 SHARD/4 |
  /tmp/tait_all_coloring_mark_separation \
  --target 4 --stop-after-first --progress 250000

clang++ -O3 -std=c++20 \
  scratch/verify_order22_separation_via_matchings.cpp \
  -o /tmp/verify_order22_separation_via_matchings

geng -cq -d3 -D3 22 SHARD/4 |
  /tmp/verify_order22_separation_via_matchings \
  --target 4 --stop-after-first --progress 250000

```

The frozen SHA-256 identifiers are:

Artifact	SHA-256
Primary source	a00c2f53a3b95a06a7e8767aae2eb2bfa794fc95eeeba18bcac34aea6a12072a
Replay source	104dca4ac143581a9b306172354428a96913f3fe1701c60b2a018510990eceed
geng 2.9.3 binary	ad2f68adf733dbed7cad543841cfa329740596ee5cd136f17a0a17f6e744f5ad
Primary run binary	a63d8048528e3dfa751049fbad6aec3b4909d1f7508f9e56859edbb5af9a6aa0
Replay run binary	3de46d3c343fb3b23679334f1058104488077386f49b4a2541273aa1e5d7d703
Corrected census report	53faa21acfaf75e6bf1d54d6c7d73150a57207067be4f44e20c55be220c322a2
Corrected run JSON	59d8e99cffb573de6ffa8a4df122099db3fd856d8e87803920e9cf69d6946ca8

The recorded environment was macOS 26.5.2 on arm64 with Apple clang 21.0.0. Mach-O linker UUIDs can make recompilations byte-distinct, so the source hashes and semantic counts are the portable anchors; the binary hashes identify the executables used in the archived runs. The JSON record includes all shard counts, commands, controls, and scope warnings. Its arithmetic totals were independently recomputed during this manuscript audit.

10 Relationship among the results and open boundary

Theorem 4.1 is unrooted: its binary cycle may contain a prescribed edge f . Theorem 5.1 is rooted but assumes the marks are separated in one fixed common-colour Tait colouring and leaves a cyclic four-cut interface. Neither theorem implies the other. Proposition 2.1 supplies the exact algebraic language for the later rooted interface. Theorem 6.4 then restricts a different, exceptional three-pole interface, but only conditionally on a singleton root signature; Theorem 6.10 turns a wider failure into an ordinary cut certificate. Neither is an elimination. Theorems 7.3 and 7.4 establish exact finite frontiers in the fixed-five simple-core model, not universal signature theorems. Theorem 8.2 says only that a particularly strong four-mark hypothesis cannot occur at order 22 in a Tait-colourable simple cubic graph. It is diagnostic evidence, not a substitute for the missing four-cut argument.

The marked-cycle human-proof frontier left by this paper is:

Analyze the rooted four-pole obtained when the forbidden edge belongs to an independent cyclic four-edge cut, while preserving the inherited mark-parity and fixed-colouring constraints.

The distinct fixed-five signature frontier is to prove universal base-pair closure or reduce the coordinate-cut certificates, and then to eliminate the surviving cyclically-two/three-cut cap interfaces beyond the completed finite slice. Nothing here classifies the published arbitrary-colour relation. Even a resolution of that local statement would still require the remaining global reductions in a Five-Cycle Double Cover proof. No claim about the orientable conjecture is made.

Acknowledgements

The author thanks the authors of the cited structural and prescribed-cycle theorems. The substantial OpenAI Codex contribution is described in the disclosure on the first page and is repeated here to avoid ambiguity: the present text and the underlying project are AI-assisted research artifacts awaiting independent human review.

References

- [1] R. E. L. Aldred, M. N. Ellingham, R. L. Hemminger, and D. A. Holton. Cycles in quasi 4-connected graphs. *Australasian Journal of Combinatorics*, 15:37–46, 1997. <https://ajc.maths.uq.edu.au/pdf/15/ocr-ajc-v15-p37.pdf>.
- [2] Paul Knappe and Max Pitz. Circuits through prescribed edges. *Journal of Graph Theory*, 93(4):470–482, 2020. <https://doi.org/10.1002/jgt.22497>.
- [3] Edita Máčajová, Giuseppe Mazzuoccolo, and Gloria Tabarelli. Cycle separating cuts in possible counterexamples to the cycle double cover and the Berge–Fulkerson conjectures. *Ars Mathematica Contemporanea*, 26:P2.03, 2026. <https://doi.org/10.26493/1855-3974.3409.c13>.
- [4] Brendan D. McKay and Adolfo Piperno. Practical graph isomorphism, II. *Journal of Symbolic Computation*, 60:94–112, 2014. <https://doi.org/10.1016/j.jsc.2013.09.003>.
- [5] Roman Nedela, Michaela Seifrtová, and Martin Škoviera. Decycling cubic graphs. *Discrete Mathematics*, 347(8):114039, 2024. <https://doi.org/10.1016/j.disc.2024.114039>.
- [6] Sang-il Oum. A proof of the cycle double cover conjecture by OpenAI: An exposition, 2026. <https://arxiv.org/abs/2607.16356>.
- [7] A. V. Pastor. On the decomposition of a 3-connected graph into cyclically 4-edge-connected components. *Journal of Mathematical Sciences*, 232(1):61–83, 2018. <https://doi.org/10.1007/s10958-018-3859-0>.
- [8] Cun-Quan Zhang. *Circuit Double Cover of Graphs*. Cambridge University Press, 2012. <https://doi.org/10.1017/CB09781139016391>.