Extra Review Problems for Exam 2

Problems

- 1. Find a vector function for the line tangent to the curve $\mathbf{r}(t) = \langle \cos t, \sin t, \cos 4t \rangle$ at $t = \pi/3$.
- 2. An object moves with velocity vector $\mathbf{v} = \langle t, t^2, \cos t \rangle$, starting at $\langle 0, 0, 0 \rangle$ at t = 0. Find the position function $\mathbf{r}(t)$.
- 3. The position function of a particle is given by $\mathbf{r}(t) = \langle t^2, 5t, t^2 16 \rangle$, for $t \ge 0$. When (at what value of t) is the speed of the particle a minimum?
- 4. A curve is given by the position function $\mathbf{r}(t) = \langle t^2, 2, t^3 \rangle$. Find the length of the curve over the interval $0 \le t \le 1$.
- 5. Find the curvature of the curve $\mathbf{r}(t) = \langle t, t^2, t \rangle$ at time t = 1.

Answers

- 1. The tangent line to the curve $\mathbf{r}(t)$ at $t = \pi/3$ is given by $\mathbf{L}(t) = \left\langle \frac{1}{2} \frac{t\sqrt{3}}{2}, \frac{\sqrt{3}}{2} + \frac{t}{2}, -\frac{1}{2} + 2t\sqrt{2} \right\rangle$
- 2. $\mathbf{r}(t) = \left\langle \frac{1}{2}t^2, \frac{1}{3}t^3, \sin t \right\rangle$
- 3. The speed function is $\sqrt{8t^2 + 25}$. Clearly, by inspection, its minimum value occurs at t = 0. The minimum speed is 5.
- 4. The length of the curve $\mathbf{r}(t)$ over the interval [0, 1] is $s = \int_0^1 t \sqrt{4 + 9t^2} dt = \frac{13\sqrt{13} 8}{27} \approx 1.44$
- 5. The curvature of $\mathbf{r}(t)$ at time t = 1 is $\kappa(1) = \frac{\sqrt{3}}{9} \approx 0.192$