

My Understanding of Survival Strategies in the Malaria Parasite *Plasmodium falciparum*

Atin Thakur,
IISER Pune.

atin.thakur@students.iiserpune.ac.in

The survival of *Plasmodium falciparum* within human erythrocytes depends on its ability to exploit host resources while adapting to environmental and drug pressures. One such example is heme metabolism. Investigation of the heme biosynthetic pathway suggests that the parasite detoxifies heme released during hemoglobin digestion by converting it into hemozoin within the digestive vacuole while still retaining genes required for de novo heme biosynthesis. Hence, investigating the relative contribution of parasite-derived and host-derived metabolic pathways to parasite survival would be interesting.

For instance, the parasite genome encodes a functional PfALAD enzyme that forms an octameric complex and is transported to the plastid-like organelle known as the apicoplast. However, quantitative enzyme assays indicate that parasite-derived PfALAD contributes only a minor fraction of the total ALAD activity observed in infected erythrocytes. Instead, host-derived ALAD appears to provide the predominant contribution.

Furthermore, a study on *Plasmodium berghei*, where host erythrocyte ALAD bound specifically to parasite membrane proteins & disruption of this interaction reduced ALAD activity, decreased heme synthesis, and inhibited parasite growth. These observations suggest that parasite survival depends not only on its own metabolic activities but also on the exploitation of host-derived enzymes and resources.

Survival under drug pressure represents another important adaptive strategy. Mutations in *PfCRT* are strongly associated with chloroquine resistance, with the K76T mutation and SVMNT haplotype being highly prevalent among resistant parasite populations. Hence, genetic variation can be selected under antimalarial pressure and contribute to parasite persistence.

The consequences of such mutations can be explored using structural approaches. Molecular docking can be used to examine the effects of mutations on ligand binding, stability, and interaction patterns, while molecular dynamics simulations may further explain how these mutations alter transport mechanisms and contribute to drug resistance.

Taken together, these studies suggest that *P. falciparum* survival relies on two complementary strategies: metabolic exploitation of host resources and evolutionary adaptation through resistance-associated mutations. Understanding these mechanisms may reveal vulnerabilities that can be targeted for future antimalarial drugs.

References:

1. **Bonday ZQ, Dhanasekaran S, Rangarajan PN, Padmanaban G.** Import of host δ -aminolevulinate dehydratase into the malarial parasite: identification of a new drug target. *Nat Med.* 2000;6(8):898–903. doi:10.1038/78659.

2. **Dhanasekaran S, Chandra NR, Chandrasekhar Sagar BK, Rangarajan PN, Padmanaban G.** δ -Aminolevulinic acid dehydratase from *Plasmodium falciparum*: molecular characterization and identification of a potent inhibitor. *J Biol Chem.* 2004;279(8):6934–6942.
3. **Varadharajan S, Dhanasekaran S, Bonday ZQ, Rangarajan PN, Padmanaban G.** Involvement of δ -aminolaevulinate synthase encoded by the parasite gene in de novo haem synthesis by *Plasmodium falciparum*. *Biochem J.* 2002;367(2):321–327. doi:10.1042/BJ20020834.
4. **Vathsala PG, Pramanik A, Dhanasekaran S, Ushadevi C, Pillai CR, Subbarao SK, et al.** Widespread occurrence of the *Plasmodium falciparum* chloroquine resistance transporter (*pfcr*t) gene haplotype SVMNT in *P. falciparum* malaria in India. *Am J Trop Med Hyg.* 2004;70(3):256–259.