

EXPLORING PERFECT BINARY TREES WITH RELATION TO THE HK-PROPERTY

ATISHAYA MAHARJAN AND MAHSA N. SHIRAZI

ABSTRACT.

1. INTRODUCTION AND PRELIMINARIES

For a given graph $G = (V, E)$, $V(G)$ and $E(G)$ denotes the vertex sets and edge sets of the graph G . For an arbitrary vertex, $v \in V(G)$, all vertices adjacent to v by an edge are called the neighbours of v and is denoted by $N_G(v)$. The degree of a vertex $v \in V(G)$ is the cardinality of the set of neighbours of v , and is denoted by $\deg_G(v)$.

For $n \in \mathbb{Z}^+$ such that $0 \leq n \leq |V(G)|$, a path of length n in G is a sequence of distinct vertices v_1, v_2, \dots, v_n such that v_i is adjacent to v_{i+1} for $1 \leq i \leq n-1$. A cycle is an extension of a path such that the last vertex is connected to the first vertex, i.e for a path of length n , $v_n v_1 \in E(G)$. As such, the length of the cycle is $n+1$.

A connected graph is a graph if for all $u, v \in V(G)$, there exists a uv -path. An independent set is a set of vertices such that no two vertices in the set are adjacent to each other. It is denoted by I . We denote a family of independent sets of a graph G as \mathcal{I}_G . A family of independent set of a graph G of cardinality n is denoted by \mathcal{I}_G^n . For $v \in V(G)$, the family of independent sets, $\mathcal{I}_G^n(v) := \{A \in \mathcal{I}_G^n : v \in A\}$ is called a star of \mathcal{I}_G^n and v is called its center.

A tree is a connected graph with no cycles, it is denoted by T . For a vertex $v \in V(T)$, if $\deg_T(v) = 1$, it is called a leaf. A vertex that is not a leaf is called an interior vertex. The depth of a vertex is defined as length of the path from the root vertex to it. We study a more particular class of trees called binary trees, denoted by T_B , where each interior vertex v has exactly 2 children and all leaves have the same depth. Further, a perfect binary tree is a binary tree in which every vertex $v \in V(T)$ has either 0 or 2 children. A perfect binary tree is denoted by T_{PB} , however in this paper we will simply drop the subscript and denote it as T for clarity. A level n of a perfect binary tree is a set of vertices such that all vertices in the set have a depth of n .

The star centers of a graph are interesting because they relate to the EKR theorem. From the EKR theorems, Holroyd and Talbot, [6], introduced the HK-property:

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, WINNIPEG, R3T 2N2, CANADA

E-mail addresses: maharjaa@myumanitoba.ca, mahsa.nasrollahi@gmail.com.

Date: March 1, 2024.

Key words and phrases. EKR, HK-property, Perfect binary trees, Escape paths.

This was in order to hopefully aid in the study of the EKR property which Holroyd and Talbot [5] conjectured as follows:

The HK-property holds true for $k \leq 4$, but was proven false independently by [1, 2, 3]. The counterexample that they arrived at is a type of graph that is defined as a class of trees called “lobsters” [4]. This is interesting for this paper as the counterexample graph resembles a binary tree.

The rest of this paper is organized in the following manner: In section 2, we introduce the flip function and escape paths. In section 3, we present our results and examples. In section 4, we present our open problems and future works.

In [4], Estrugo and Pastine came up with a generalized flip function which is defined as follows:


This definition admits the following properties [4]:

They [4] also defined the escape path as follows:

Definition 2.3. Let G be a graph and $P = v_1 v_2, \dots v_n$ a path of length n such that $P \subset G$. We say that P is an escape path from v_1 to v_n in G if $\deg(v_n) = 1$ and $\deg(v_i) = 2$ for $2 \leq i \leq n - 2$. If this is the case we say that v_1 has an escape path to v_n .

Our objective is to expand the flip function to accomodate the structure of the perfect binary tree. We want this function to map independent sets from an arbitrary vertex into independent sets that contain a leaf. Doing so, we will show that the flip function is injective. Furthermore, we aim to conserve the property of the flip function being an involution.

We first show that for any vertex set of a fixed depth, k , the vertex choice does not matter. To do so, we first define an indexing for the vertices of depth k .



Definition 2.4 (Depth Vertex). Let $\mathcal{V}_k \in V(T)$ be the set of vertices of depth k . We call \mathcal{V}_k as the depth vertex set of depth k . Index all vertices in \mathcal{V}_k from left to right as $v_{k,i}$, where k is the depth of the vertex and i is the index of the vertex in \mathcal{V}_k such that $1 \leq i \leq 2^{k-1}$.

We will now define the flip function on the vertices of depth k .

Definition 2.5 (Flip Function on Depth Vertex Set). Let T be a perfect binary tree and \mathcal{V}_k be the depth vertex set of depth k . Then, the flip of \mathcal{V}_k in T , denoted by $flip_{\mathcal{K}} : \mathcal{V}_k(G) \rightarrow \mathcal{V}_k(G)$, is the function defined as follows:

$$flip_{\mathcal{K}}(v_{k,i}) = v_{(k, 2^{k-1}+1-i)}$$

We now have the following lemma:

Lemma 2.6 ($flip_{\mathcal{K}}$ is a bijective involution). Let T be a perfect binary tree and \mathcal{V}_k be the depth vertex set of depth k . Then, the flip function $flip_{\mathcal{K}}$ is an involution.

Proof. Let \mathcal{V}_k be the depth vertex set of depth k . Then, applying the flip function twice, we have:

$$\begin{aligned} flip_{\mathcal{K}}(flip_{\mathcal{K}}(v_{k,i})) &= flip_{\mathcal{K}}(v_{(k, 2^{k-1}+1-i)}) \\ &= v_{(k, 2^{k-1}+1-(2^{k-1}+1-i))} \\ &= v_{(k,i)} \end{aligned}$$

Hence, $flip_{\mathcal{K}}$ is an involution. \square

We then proceed to show that the flip function on a depth vertex set maps independent sets into independent sets.

Lemma 2.7. Let T be a perfect binary tree and \mathcal{V}_k be the depth vertex set of depth k . Let $\mathcal{I}_{\mathcal{V}_k}^m$ be the family of independent sets of cardinality m . Then, the flip function $flip_{\mathcal{K}}$ maps independent sets into independent sets and induces a bijection from $\mathcal{I}_{\mathcal{V}_k}^m$ onto itself.

In addition, $flip_{\mathcal{K}}(\mathcal{I}_{\mathcal{V}_k}(v_{k,1})) = \mathcal{I}_{\mathcal{V}_k}(v_{k,2^{k-1}})$.

Proof. Let \mathcal{I} be an independent set and assume two unique vertices $v_{k,i}, v_{k,j} \in flip_{\mathcal{K}}(\mathcal{I})$, i.e $i \neq j$. Since both $v_{k,i}$ and $v_{k,j}$ belong to the depth vertex set, and since this is a perfect binary tree, we see that $v_{k,i}$ and $v_{k,j}$ are not adjacent to each

other. Hence, $flip_{\mathcal{K}}(\mathcal{I})$ is an independent set. Thus, $flip_{\mathcal{K}}$ maps independent sets into independent sets.

Now, let $\mathcal{I}_{\mathcal{V}_k}^m$ be the family of independent sets of cardinality m of the depth vertex set \mathcal{V}_k . Then, $flip_{\mathcal{K}}$ is a bijection from $\mathcal{I}_{\mathcal{V}_k}^m$ onto itself.

Finally, we have $flip_{\mathcal{K}}(\mathcal{I}_{\mathcal{V}_k}(v_{k,1})) = \mathcal{I}_{\mathcal{V}_k}(v_{k,2^{k-1}+1-1}) = \mathcal{I}_{\mathcal{V}_k}(v_{k,2^{k-1}})$, as required.

Hence, $flip_{\mathcal{K}}$ maps independent sets into independent sets and induces a bijection from $\mathcal{I}_{\mathcal{V}_k}^m$ onto itself. \square

From [Lemma 2.7](#) and [2.6](#), we can easily see that we can pick any arbitrary vertex in any depth vertex set to show that our next flip function holds.

Now, our objective is to procure a similar result to [\[4\]](#) to show that our flip function induces a one to one mapping from any family of independent set $\mathcal{I}_T(v)$ for any vertex $v \in V(T)$ into any family of independent set $\mathcal{I}_T(l)$ for a leaf $l \in V(T)$.

3. OPEN PROBLEMS AND FUTURE WORKS

REFERENCES

- [1] R. Baber. *Some results in extremal combinatorics*. ProQuest LLC, Ann Arbor, MI, 2011. Thesis (Ph.D.)—University of London, University College London (United Kingdom).
- [2] Peter Borg. Stars on trees. *Discrete Math.*, 340(5):1046–1049, 2017.
- [3] Peter Borg and Fred Holroyd. The Erdős-Ko-Rado properties of various graphs containing singletons. *Discrete Math.*, 309(9):2877–2885, 2009.
- [4] Emiliano J.J. Estrugo and Adrián Pastine. On stars in caterpillars and lobsters. *Discrete Appl. Math.*, 298:50–55, 2021.
- [5] Fred Holroyd and John Talbot. Graphs with the Erdős-Ko-Rado property. *Discrete Math.*, 293(1-3):165–176, 2005.
- [6] Glenn Hurlbert and Vikram Kamat. Erdős-Ko-Rado theorems for chordal graphs and trees. *J. Combin. Theory Ser. A*, 118(3):829–841, 2011.