

model-calibration

Alexander Keth

2017-05-04

Contents

0.1	User Input	3
1	Whole system plots!	4
1.1	Overall biomass	4
1.2	Biomass timeseries	5
1.3	Biomass@age timeseries	6
1.4	Number timeseries	7
1.5	Number@age timeseries	8
1.6	SSB & Recruitment	9
1.7	Biomass benchmark	10
1.8	Biomass benchmark 2	11
1.9	Physics	12
2	Physics	13
2.1	Chl_a	13
2.2	Denitrification	14
2.3	NH3	15
2.4	NO3	16
2.5	Temp	17
2.6	salt	18
2.7	Fluxes 1	19
2.8	Fluxes 2	20
2.9	Relative change of water column height compared to nominal_dz	21
3	Calibration plots	22
3.1	Structural nitrogen	22
3.2	Reserve nitrogen	23
3.3	Biomass per ageclass	24
3.4	Eat per ageclass	25
3.5	Growth per ageclass	26
3.6	Growth in relation to initial conditions	27
3.7	Numbers	28
3.8	Biomass	29
4	Distribution plots	30
4.1	Numbers @ age	30
4.2	Biomass @ age	31
5	Diet Plots	32
5.1	Diet plot 1: Cephalopod	33

5.2	Diet plot 2: Diatom	34
5.3	Diet plot 3: Labile detritus	35
5.4	Diet plot 4: Megazoobenthos	36
5.5	Diet plot 5: Refractory detritus	37
5.6	Diet plot 6: Shallow piscivorous fish	38
5.7	Diet plot 7: Small planktivorous fish	39
6	Spatial Plots 1	40
6.1	Spatial Plot 1: Carrion3 1	40
6.2	Spatial Plot 2: Cephalopod 1	41
6.3	Spatial Plot 3: Diatom 1	42
6.4	Spatial Plot 4: Labile detritus 1	43
6.5	Spatial Plot 5: Megazoobenthos 1	44
6.6	Spatial Plot 6: Refractory detritus 1	45
6.7	Spatial Plot 7: Shallow piscivorous fish 1	46
6.8	Spatial Plot 8: Shallow piscivorous fish 2	47
6.9	Spatial Plot 9: Small planktivorous fish 1	48
6.10	Spatial Plot 10: Small planktivorous fish 2	49
7	Spatial Plots 2	50
7.1	Spatial Plot 1: Carrion3	51
7.2	Spatial Plot 2: Cephalopod	52
7.3	Spatial Plot 3: Diatom	53
7.4	Spatial Plot 4: Labile detritus	54
7.5	Spatial Plot 5: Megazoobenthos	55
7.6	Spatial Plot 6: Refractory detritus	56
7.7	Spatial Plot 7: Shallow piscivorous fish	57
7.8	Spatial Plot 8: Small planktivorous fish	58

NOTE: This vignette is optimised for longer simulation runs. Therefore the output is not as pleasant due to the fact that the dummy setas file have a running time of 5 years.

In order to use this vignette make sure to render `model-preprocess.Rmd` first. Either save the resulting list of dataframes as shown in `data-raw/data-vignette-model-preprocess.R` or render both vignettes `model-preprocess.Rmd` and `model-calibration.Rmd` in the same R-instance. Of course, you can also use a personalised version of `model-preprocess.Rmd`. Please make sure to add all resulting dataframes to the list of dataframes at the end of the preprocess vignette and change `model-calibration.Rmd` accordingly.

```
library("atlantistools")
library("ggplot2")
library("gridExtra")

fig_height2 <- 11
gen_labels <- list(x = "Time [years]", y = "Biomass [t]")

# You should be able to build the vignette either by clicking on "Knit PDF" in RStudio or with
# rmarkdown::render("model-calibration.Rmd")
```

0.1 User Input

This section is used to read in the SETAS dummy files. Please change this accordingly.

```
result <- preprocess

d <- system.file("extdata", "setas-model-new-trunk", package = "atlantistools")

# External recruitment data
ex_rec_ssb <- read.csv(file.path(d, "setas-ssb-rec.csv"), stringsAsFactors = FALSE)

# External biomass data
ex_bio <- read.csv(file.path(d, "setas-bench.csv"), stringsAsFactors = FALSE)

# bgm file
bgm <- file.path(d, "VMPPA_setas.bgm")
```

1 Whole system plots!


1.1 Overall biomass

```
df_bio <- combine_groups(result$biomass, group_col = "species", combine_thresh = 10)
```

```
## Joining, by = "species"
```


```
plot <- plot_bar(df_bio)
```

```
update_labels(plot, labels = gen_labels)
```




1.2 Biomass timeseries

```
plot <- plot_line(result$biomass)
update_labels(plot, labels = gen_labels)
```




1.3 Biomass@age timeseries

```
plot <- plot_line(result$biomass_age, col = "agecl")
update_labels(p = plot, labels = c(gen_labels, list(colour = "Ageclass")))
```




1.4 Number timeseries

```
plot <- plot_line(result$nums)
update_labels(p = plot, labels = list(x = "Time [years]", y = "Numbers"))
```




1.5 Number@age timeseries

```
plot <- plot_line(result$num_age, col = "agecl")
update_labels(p = plot, labels = list(x = "Time [years]", y = "Numbers", colour = "Ageclass"))
```



1.6 SSB & Recruitment

```
plot_rec(result$ssb_rec, ex_data = ex_rec_ssb)
```



1.7 Biomass benchmark

```
names(ex_bio)[names(ex_bio) == "biomass"] <- "atoutput"
```

```
data <- result$biomass
```

```
data$model <- "atlantis"
```

```
comp <- rbind(ex_bio, data, stringsAsFactors = FALSE)
```


```
# Show atlantis as first factor!
```

```
comp$model <- factor(comp$model, levels = c("atlantis", sort(unique(comp$model))[sort(unique(comp$model)) != "atlantis"])))
```

```
# Create plot
```


```
plot <- plot_line(comp, col = "model")
```

```
update_labels(plot, gen_labels)
```




1.8 Biomass benchmark 2

```
plot <- plot_line(result$biomass) %>% update_labels(labels = gen_labels)
plot_add_range(plot, ex_bio)
```



1.9 Physics


```
plot <- plot_line(result$physics, wrap = NULL)
custom_grid(plot, grid_x = "polygon", grid_y = "variable")
```




2 Physics

```
physics <- result$physics %>%  
  flip_layers() %>%  
  split(., .$variable)  
  
plots <- lapply(physics, plot_line, wrap = NULL) %>%  
  lapply(., custom_grid, grid_x = "polygon", grid_y = "layer")  
  
for (i in seq_along(plots)) {  
  cat(paste0("## ", names(plots)[i]), sep = "\n")  
  plot <- update_labels(plots[[i]], labels = list(y = names(plots)[i]))  
  print(plot)  
  cat("\n\n")  
}
```


2.1 Chl_a




2.2 Denitrification




2.3 NH3




2.4 NO3



2.5 Temp




2.6 salt




2.7 Fluxes 1

```
plot <- flip_layers(result$flux) %>%  
  plot_line(wrap = NULL, col = "variable")  
custom_grid(plot, grid_x = "polygon", grid_y = "layer")
```



2.8 Fluxes 2


```
plot <- flip_layers(result$sink) %>%  
  plot_line(wrap = NULL, col = "variable")  
custom_grid(plot, grid_x = "polygon", grid_y = "layer")
```



2.9 Relative change of water column height compared to nominal_dz

```
check_dz <- result$dz %>%  
  dplyr::left_join(result$nominal_dz, by = c("polygon", "layer")) %>%  
  dplyr::mutate(check_dz = atoutput.x / atoutput.y) %>%  
  dplyr::filter(!is.na(check_dz)) # remove sediment layer
```


```
plot <- plot_line(check_dz, x = "time", y = "check_dz", wrap = "polygon", col = "layer")  
update_labels(plot, list(x = "Time [years]", y = expression(dz/nominal_dz)))
```



3 Calibration plots


3.1 Structural nitrogen

```
df_rel <- convert_relative_initial(result$structn_age)
plot <- plot_line(df_rel, col = "agecl")
plot <- update_labels(plot, list(x = "Time [years]", y = expression(SN/SN[init])))
plot_add_box(plot)
```




3.2 Reserve nitrogen

```
df_rel <- convert_relative_initial(result$resn_age)
plot <- plot_line(df_rel, col = "agec1")
plot <- update_labels(plot, list(x = "Time [years]", y = expression(RN/RN[init])))
plot_add_box(plot)
```




3.3 Biomass per ageclass

```
df_rel <- convert_relative_initial(result$biomass_age)
plot <- plot_line(df_rel, col = "agecl")
plot <- update_labels(plot, list(x = "Time [years]", y = expression(Biomass/Biomass[init])))
plot_add_box(plot)
```




3.4 Eat per ageclass

```
df_rel <- convert_relative_initial(result$eat_age)
plot <- plot_line(df_rel, col = "agec1")
plot <- update_labels(plot, list(x = "Time [years]", y = expression(Cons./Cons.[init])))
plot_add_box(plot)
```




3.5 Growth per ageclass

```
df_rel <- convert_relative_initial(result$growth_age)
plot <- plot_line(df_rel, col = "agec1")
plot <- update_labels(plot, list(x = "Time [years]", y = expression(Growth/Growth[init])))
plot_add_box(plot)
```




3.6 Growth in relation to initial conditions

```
plot <- plot_line(result$growth_rel_init, y = "gr_rel", col = "agecl")
update_labels(plot, list(y = expression((Growth - Growth[req])/Growth[req])))
```




3.7 Numbers

```
df_rel <- convert_relative_initial(result$nums_age)
plot <- plot_line(df_rel, col = "agec1")
plot <- update_labels(plot, list(x = "Time [years]", y = expression(Numbers/Numbers[init])))
plot_add_box(plot)
```



3.8 Biomass


```
df_rel <- convert_relative_initial(result$biomass)
plot <- plot_line(df_rel)
plot <- update_labels(plot, list(x = "Time [years]", y = expression(Biomass/Biomass[init])))
plot_add_box(plot)
```



4 Distribution plots


4.1 Numbers @ age

```
df <- agg_perc(result$nums_age, groups = c("time", "species"))
plot <- plot_bar(df, fill = "agecl", wrap = "species")
update_labels(plot, labels = list(x = "Time [years]", y = "Numbers [%]"))
```



4.2 Biomass @ age


```
df <- agg_perc(result$biomass_age, groups = c("time", "species"))
plot <- plot_bar(df, fill = "agecl", wrap = "species")
update_labels(plot, labels = list(x = "Time [years]", y = "Biomass [%]"))
```



5 Diet Plots


```
## Joining, by = c("time", "pred", "agecl", "prey")  
## Joining, by = c("time", "pred", "agecl", "prey")
```


5.1 Diet plot 1: Cephalopod




5.2 Diet plot 2: Diatom

Diet proportions for species: Diatom




5.3 Diet plot 3: Labile detritus




5.4 Diet plot 4: Megazoobenthos




5.5 Diet plot 5: Refractory detritus



5.6 Diet plot 6: Shallow piscivorous fish




5.7 Diet plot 7: Small planktivorous fish





6 Spatial Plots 1


Joining, by = "polygon"


6.1 Spatial Plot 1: Carrion3 1



























7 Spatial Plots 2




```
## Joining, by = c("time", "polygon")

## geom_path: Each group consists of only one observation. Do you need to
## adjust the group aesthetic?



## geom_path: Each group consists of only one observation. Do you need to
## adjust the group aesthetic?
```





Species: Cephalopod with stanza: 1






Species: Diatom with stanza: 1








Species: Megazoobenthos with stanza: 1





Species: Shallow piscivorous fish with stanza: 1



Species: Shallow piscivorous fish with stanza: 2

