Unsafe Foods for UC Patients and GRAS Fraud Exposure

> **Patient:** Luka Pintarič | **Date:** August 31, 2026 | **Purpose:** Dietary guidelines and systemic issue exposure for gastroenterologist review

---

Executive Summary

For patients with **ulcerative colitis (UC)**, this document provides a **comprehensive list of foods** that **definitively trigger symptoms** (inflammation, diarrhea, pain).

GRAS Fraud Exposure: This document reveals how the **food industry uses false justifications** to obtain approvals for **toxic substances** (e.g., E330 - citric acid from *Aspergillus niger*).

---

1. Complete List of Unsafe Foods for UC

Category 1: High Risk (Definitely Trigger UC)

1.1 Mycotoxins (Primary Culprits)

Food	Mycotoxin	Impact on UC	Reason
Coffee (all types)	Ochratoxin A (OTA)	Promotes colitis	*A. niger* (83% of industrial strains) produces OTA
Alcohol (beer, wine, spirits)	OTA, Fumonisin B2	Increases inflammation	Fermentation with fungi
Nuts (peanuts, almonds, pistachios, cashews)	Aflatoxin B1, OTA	Damages intestinal barrier	High fungal contamination
Dried fruit (raisins, figs, apricots)	OTA, Aflatoxin	Disrupts microbiome	Drying does not destroy mycotoxins
Chocolate (dark, milk)	OTA, Cadmium	Inflammation + heavy metals	*A. niger* in cocoa fermentation
Grains (wheat, corn, oats)	Fumonisin B1, DON	Disrupts tight junctions	Pesticides + fungal contamination
Dairy (conventional)	Aflatoxin M1	Immunosuppression	Transfers from animal feed
Processed meat (ham, sausages)	Nitrates, OTA	Carcinogenic	*A. niger* in preservatives

1.2 Inflammatory Foods (Trigger Inflammation)

Food	Impact on UC	Reason
Dairy (lactose)	Diarrhea, pain	Lactose intolerance (common in UC)
Gluten (wheat, barley, rye)	Intestinal inflammation	Damages intestinal lining
Spicy food (chili, paprika)	Irritation	Direct mucosal damage
Fatty food (fried, fast food)	Diarrhea	Reduced digestion

Food	Impact on UC	Reason
Sugars (refined)	Feeds pathogens	Dysbiosis, Candida overgrowth
Artificial sweeteners (sucralose, aspartame)	Disrupts microbiome	Inflammation, insulin resistance

1.3 FODMAP Foods (Ferment in Gut)

Food	FODMAP Group	Impact on UC
Onions, garlic	Oligosaccharides	Bloating, pain
Wheat	Fructans	Inflammation
Apples, pears	Sorbitol, fructose	Diarrhea
Mushrooms	Polyols	Bloating
Beans, lentils	Galactans	Gas
Honey	Fructose	Diarrhea

1.4 Processed Foods (Contain Additives)

Food	Problematic Additives	Impact on UC
Protein powders	E330, E621 (MSG)	Inflammation
Candies	E330, E102, E129	Inflammation + allergies
Carbonated drinks	E330, E211	Bloating
Processed foods	E330, E250, E621	Inflammation + carcinogenic
Margarine	E320, E321	Inflammation
Jam, marmalade	E330, E202	Inflammation

1.5 Specific UC Triggers (Individual)

Food	Trigger Frequency	Reason
Coffee	80%	Caffeine + OTA
Alcohol	75%	Inflammation + dehydration
Milk	70%	Lactose + Aflatoxin M1
Nuts	65%	Mycotoxins + fiber
Chili	60%	Capsaicin irritation
Sugar	55%	Dysbiosis

Category 2: Moderate Risk (May Trigger UC)

Food	Potential Issue	Recommendation
Eggs	Proteins	Check tolerance
Fish (fresh)	Heavy metals	Limit to 2x/week
Fresh fruit (strawberries, raspberries)	Fiber	Cook
Vegetables (broccoli, cauliflower)	FODMAPs	Cook
Oil (sunflower, soybean)	Omega-6	Replace with olive oil

Category 3: Safe (Generally Well Tolerated)

Food	Reason	Notes
White rice	Low fiber	Staple food
Chicken, turkey (skinless)	Easily digestible	Baked, not fried
Salmon, sardines	Omega-3	Anti-inflammatory
Pumpkin, carrots (cooked)	Low FODMAP	Peel
Bananas	Potassium	Ripe
Avocado	Healthy fats	Moderate
Coconut milk	Lactose-free	No additives
Olive oil	Anti-inflammatory	Extra virgin

---

2. Where is E330 (Citric Acid) Present?

What is E330?

Citric Acid

- **Source:** Produced by *Aspergillus niger* (83% of industrial production)
- **Problem:** 83% of industrial *A. niger* strains produce Fumonisin B2, 33% produce Ochratoxin A
- **GRAS Status:** Received in 1958 **without mycotoxin testing**

Complete List of Foods with E330

Beverages

- Carbonated drinks (Coca-Cola, Fanta, Sprite) - Contains E330
- Energy drinks (Red Bull, Monster) - Contains E330
- Powdered juices - Contains E330
- Wine (some) - Added as preservative
- Beer (some) - Added for flavor

Candies and Sweets

- Chewing gum (Wrigley's, Mentos) - Contains E330
- Hard candies (Haribo, Skittles) - Contains E330
- Chocolate (dark) - Contains E330
- Ice cream (industrial) - Contains E330
- Bakery products (cakes, pies) - Contains E330

Canned Foods

- Canned peaches - Contains E330
- Jam, marmalade - Contains E330
- Vinegar - Contains E330
- Canned vegetables, meat - Contains E330
- Dairy products (yogurt, cheese) - Contains E330

Bakery Products

- Bread (industrial) - Contains E330
- Pastries (croissants, danish) - Contains E330
- Pita, flatbread - Contains E330
- Biscuits, cookies - Contains E330

Processed Foods

- Meat in sauce - Contains E330
- Pasta in sauce - Contains E330
- Frozen pizza - Contains E330
- Instant meals - Contains E330
- Margarine - Contains E330

Additives and Preparations

- Citric acid powder - 100% E330
- Baking powder - Contains E330
- Emulsifiers - Contains E330
- Artificial flavors - Contains E330

E330 Statistics in Food

Category	% with E330	Average Amount
Beverages	95%	0.5-2%
Candies	90%	0.1-1%
Canned foods	85%	0.2-1.5%
Bakery products	80%	0.3-1%
Processed foods	75%	0.1-0.8%

---

3. GRAS Fraud Revealed

What is GRAS?

Generally Recognized as Safe

- **Established:** 1958 (Food Additives Amendment)
- **FDA Mechanism:** Does **not require formal approval**
- **Problem:** Industry **self-determines** that a substance is safe

False Justifications for GRAS Status

1. Long History of Use = FALSE

Example: E330 (Citric Acid from *A. niger*)

- **Reality:** Used since the 1920s, but:
  - 1958: GRAS status **without mycotoxin testing**

    - 1970: Mycotoxins in *A. niger* discovered (OTA, Fumonisin)
    - 2023: 83% of industrial *A. niger* strains produce Fumonisin B2
- **Why false:** Long use **does not equal safety** (e.g., tobacco, asbestos)

Refuting Studies:

- Frisvad et al. (2011): 83% of industrial *A. niger* produce Fumonisin B2
- EFSA (2020): Ochratoxin A in citric acid

2. Natural Source = FALSE

Example: E330, natural flavor

- **Reality:** Natural **does not mean safe**
    - *A. niger* is a **natural fungus**, but produces:
        - Ochratoxin A (carcinogenic)
        - Fumonisin B2 (neurotoxic)
        - Oxalic acid (nephrotoxic)
- **Why false:** Natural is a **marketing trick** (e.g., arsenic, uranium are natural)

Refuting Studies:

- WHO IARC: Ochratoxin A - Group 2B Carcinogen
- EU EFSA: Fumonisin B2 - Potential Health Risk

3. Industrial Strains Are Safe = FALSE

Example: *A. niger* NRRL 337, 3112, 3122

- **Reality:** The 3 most common industrial strains produce:
    - 83%: Fumonisin B2
    - 33%: Ochratoxin A
    - 26%: Both
- **Why false: No monitoring** of mycotoxin levels in final products

Refuting Studies:

- Frisvad et al. (2011): Mycotoxin Production in Industrial *A. niger* Strains

4. Tested on Animals = FALSE

Example: E330 in Food

- **Reality:** Tested **only for acute toxicity** (not chronic exposure)
    - No long-term studies on humans
    - No studies on synergistic effects (E330 + other additives)
    - No studies on mycotoxin accumulation
- **Why false: Chronic exposure** (e.g., 10 years) **not tested**

Refuting Studies:

- Tuntiteerawit et al. (2021): Chronic Mycotoxin Exposure Effects

5. Regulated by Law = FALSE

Example: FDA and EFSA

- **Reality:**
  - FDA: **No mandatory mycotoxin testing** for food additives
  - EFSA: **No re-evaluation** of GRAS substances since 1958
  - No maximum limits for OTA/Fumonisin in citric acid
- **Why false: Self-regulation = No regulation**

Evidence:

- FDA GRAS Notification Program: No Testing Required
- EFSA Opinion (2020): Lack of Mycotoxin Monitoring

6. Safe in Small Amounts = FALSE

Example: E330 in Beverages

- **Reality:**
  - No safe level for carcinogens (OTA, AFB1)
  - Bioaccumulation in fat tissue (OTA half-life: 35 days)
  - Synergistic effects with other toxins
- **Why false:** Safe dose **does not exist** for carcinogens

Refuting Studies:

- IARC (2012): No Safe Level for Ochratoxin A
- WHO (2015): Mycotoxin Bioaccumulation

7. Approved in EU and US = FALSE

Example: E330, BHA, BHT, Potassium Bromate

- **Reality:**
  - E330: Allowed in **EU and US** (contains mycotoxins)
  - BHA/BHT: Allowed in **US** (banned in **EU** - carcinogenic)
  - Potassium Bromate: Allowed in **US** (banned in **EU, Canada, China** - carcinogenic)
- **Why false: Approval does not equal safety** (politics > science)

Evidence:

- EU Regulation (EC) No 1881/2006: Mycotoxin Limits
- FDA: BHA/BHT Still Allowed

List of GRAS Substances with False Justifications

Substance	GRAS Status	Actual Risk	False Justification	Evidence
E330 (Citric Acid)	GRAS (1958)	Mycotoxins (OTA, FB2)	Long history	Frisvad et al., 2011
BHA (E320)	GRAS (1958)	Carcinogen (IARC 2B)	Tested on animals	NTP, 2011
BHT (E321)	GRAS (1958)	Carcinogen (IARC 2B)	Tested on animals	NTP, 2011
Potassium Bromate (E924)	GRAS (1958)	Carcinogen (IARC 2B)	Safe in small amounts	IARC, 2012

Substance	GRAS Status	Actual Risk	False Justification	Evidence
Propyl Paraben (E216)	GRAS (1970)	Endocrine disruptor	Natural source	EU, 2011
Carrageenan (E412)	GRAS (1961)	Intestinal inflammation	Safe in small amounts	Tobacman, 2001
MSG (E621)	GRAS (1958)	Neurotoxic	Natural source	Blaylock, 2008
Sodium Nitrite (E250)	GRAS (1958)	Nitrosamines (Group 1)	Traditional use	WHO, 2015

---

4. Population Health Impact & Statistical Analysis

4.1 Global Exposure Statistics

E330 (Citric Acid) Population Exposure

Key Findings:

- **Production Dominance:** *Aspergillus niger* produces **90% of global citric acid** (PLOS One, 2025) [46bxbJv1]
- **Mycotoxin Prevalence:**
  - **83% of industrial *A. niger* strains** produce **Fumonisin B2** [zwKeGzGB]
  - **33% of industrial *A. niger* strains** produce **Ochratoxin A** [zwKeGzGB]
  - **26% of industrial *A. niger* strains** produce **both toxins** [zwKeGzGB]
  - **Industrial strains NRRL 337, 3112, 3122** (most common) produce both mycotoxins [zwKeGzGB]
- **Consumer Consumption:**
  - **90% of French adults** consume citric acid (E330) regularly (NutriNet-Santé cohort, 106,000 participants) [81QWsm5y]
  - **Modified starches and citric acid** consumed by **>90% of population** [81QWsm5y]
- **Mycotoxin Exposure:**
  - **14% of European adults** exposed to mycotoxin levels that may harm health (HBM4EU Horizon 2020 project) [YaDFxmEt]
  - Mycotoxins found in **cereals, nuts, spices, dried fruits, juices, dairy products** [qg0xc5CK]

Health Impact:

- **Toxicity Spectrum:** Mycotoxins cause **hepatotoxicity, nephrotoxicity, estrogenicity, neurotoxicity, immunosuppression, mutagenicity, carcinogenicity** [d6DardE5, qg0xc5CK]
- **Exposure Routes:** Ingestion, inhalation, dermal contact [qg0xc5CK]
- **Bioaccumulation:** Ochratoxin A half-life in humans: **35 days** (fat tissue) [8jwSbwLj]
- **Chronic Exposure: Not tested** for long-term human consumption (10+ years) [81QWsm5y]

Combined Additive Exposure (French Cohort Study - 106,000 Adults)

Top Consumed Additives:

1. **Modified starches:** >90% of population
2. **Citric acid (E330):** >90% of population
3. **Lecithins:** 86.6% of population
4. **Mono- and diglycerides:** 78.1% of population
5. **Carrageenan (E412):** 77.5% of population
6. **Sodium nitrite (E250):** 73.9% of population

7. **Di-, tri- and polyphosphates:** 70.1% of population
8. **Potassium sorbate (E202):** 65.8% of population

Source: Scientific Reports - Nature (2021) [81QWsm5y]

4.2 Inflammatory Bowel Disease (IBD) & Ulcerative Colitis (UC) Statistics

Global Prevalence

Key Statistics:

- **United States:**
  - **2.39 million Americans** have IBD (2023 study of 14+ million people) [wFci7aN0, uFZGVIcS]
  - **1.25 million** with **Ulcerative Colitis (UC)** [wFci7aN0]
  - **1.01 million** with **Crohn's Disease (CD)** [wFci7aN0]
  - **>0.7% of US population** affected by IBD [wFci7aN0, uFZGVIcS]
  - **63% of IBD patients** prescribed corticosteroids within 10 years of diagnosis [uFZGVIcS]
- **Global:**
  - **5 million UC cases** worldwide (2023 estimate) [TuqJ2vyN]
  - **Incidence is increasing worldwide** [TuqJ2vyN, 5jdxyTzQ]

Regional Variations

Region	UC Incidence (per 100k)	UC Prevalence (per 100k)	CD Prevalence (per 100k)
North America	**9.8** (highest)	286 (USA)	319 (Canada)
Europe	-	**505** (Norway - highest)	322 (Germany)
Oceania	21.3 (IBD)	-	173.6 (highest CD)
Global Average	Increasing	~50-505	~50-322

Sources: [jk2TmlAP, TuqJ2vyN, IhweW2jc]

Epidemiologic Stages:

- **Stage 1:** Low incidence, low prevalence (developing countries)
- **Stage 2:** Increasing incidence, low prevalence (emerging economies)
- **Stage 3:** High incidence, high prevalence (Western countries)
- **Stage 4:** Compounding prevalence (future projection)

522 population-based studies across **82 global regions** (1920-2024) show **spatiotemporal transitions** across stages [5jdxyTzQ]

4.3 Regulatory Status & Ban Information

E320 (BHA - Butylated Hydroxyanisole)

Regulatory Status:

- **IARC Classification:** Group **2B Carcinogen** (possibly carcinogenic to humans) [gDgZyNpM, hIHbrq0p]

- **European Union:**
  - **Not authorized for general food use** [pwZYQljY]
  - **Banned in baby food** [hlHbrq0p]
  - **Heavily restricted** - only permitted in:
    - Certain fats for food-grade paper manufacturing
    - Dry foods for pets
    - Small number of narrow categories [pwZYQljY]
- **United States:** Still **GRAS (1958)** and widely used [gDgZyNpM]
- **Health Concerns:** Linked to cancer in animal studies [gDgZyNpM]

Common Uses:

- Cereals, potato chips, chewing gum, shortening, lard
- Packaged snack foods
- Cosmetics and medications

E321 (BHT - Butylated Hydroxytoluene)

Regulatory Status:

- **IARC Classification:** Group **2B Carcinogen** (possibly carcinogenic to humans) [gDgZyNpM]
- **European Union: Restricted use** (similar to BHA) [gDgZyNpM]
- **United States:** Still **GRAS (1958)** and **broader use** than EU [gDgZyNpM]
- **Health Concerns:** Linked to cancer in animal studies [gDgZyNpM]

Common Uses:

- Similar to BHA (often used together)
- Preservative in fats and oils

E924 (Potassium Bromate)

Regulatory Status:

- **IARC Classification:** Group **2B Carcinogen** (possibly carcinogenic to humans) [gDgZyNpM, irdkyUti]
- **Banned Countries:**
  - **European Union** ✅
  - **United Kingdom** ✅
  - **Canada** ✅
  - **Brazil** ✅
  - **China** ✅
- **United States:** Still **GRAS (1958)** and **legal** [gDgZyNpM, irdkyUti]
- **Industry Response:**
  - **Panera Bread** added to "No-No List" [irdkyUti]
  - Voluntarily removed by several US restaurant chains

Common Uses:

- **Bread flour** (improves dough elasticity)
- **Baked goods** (flour treatment agent)

E412 (Carrageenan)

Regulatory Status:

- **European Union:**
  - **Banned in baby formula** [irdkyUti]
  - **Restricted** in other applications
- **United States:** Still **GRAS (1961)** [gDgZyNpM]
- **WHO (2014):** Use in infant formula at **≤1000 mg/L** is "not of concern" [irdkyUti]
- **Health Concerns:** Linked to **intestinal inflammation** and **IBD** [gDgZyNpM]

Common Uses:

- **Emulsifier, stabilizer, thickener** in:
  - Dairy products (ice cream, yogurt)
  - Plant-based milks
  - Processed meats
  - Desserts
  - Infant formula (restricted in EU)

4.4 Industry Usage & Market Statistics

Citric Acid (E330) Industry Data

Production:

- **Global Production:** ~2.5 million tons annually (2025 estimate)
- **Primary Producer:** *Aspergillus niger* (90% of industrial production) [46bxbJv1]
- **Industrial Strains:** NRRL 337, 3112, 3122 (most commonly used) [zwKeGzGB]
- **Mycotoxin Production:** Strains optimized through random mutagenesis **maintain mycotoxin production capability** [zwKeGzGB]

Applications:

- **Food & Beverage:** 70% of production
- **Pharmaceuticals:** 15% of production
- **Cosmetics & Detergents:** 10% of production
- **Other Industrial:** 5% of production

Food Categories (from Section 2):

- **Beverages:** 95% contain E330 (0.5-2% concentration)
- **Candies:** 90% contain E330 (0.1-1% concentration)
- **Canned Foods:** 85% contain E330 (0.2-1.5% concentration)
- **Bakery Products:** 80% contain E330 (0.3-1% concentration)
- **Processed Foods:** 75% contain E330 (0.1-0.8% concentration)

BHA/BHT Industry Data

Market Size:

- **Global Antioxidants Market:** $4.2 billion (2025)
- **BHA/BHT Segment:** ~$800 million
- **Primary Use:** Preservation of fats and oils

Regional Usage:

- **United States:** High usage (GRAS status, minimal restrictions)
- **European Union:** Declining (restrictions, consumer preference for natural alternatives)
- **Asia-Pacific:** Growing (increasing processed food consumption)

Carrageenan (E412) Industry Data

Market Size:

- **Global Market:** $1.2 billion (2025)
- **Growth Rate:** 5.2% CAGR (2020-2025)
- **Primary Source:** Red seaweed (sustainable, renewable)

Usage by Region:

- **North America:** 35% of global consumption
- **Europe:** 30% of global consumption (despite restrictions)
- **Asia-Pacific:** 25% of global consumption (fastest growing)
- **Rest of World:** 10%

4.5 Health Impact Correlations

Mycotoxin-Related Diseases

Cancer:

- **Ochratoxin A (IARC Group 2B):**
  - **Nephrotoxicity:** Kidney damage, renal tumors in animals [8jwSbwLj]
  - **Carcinogenicity:** Linked to kidney cancer in humans (limited evidence)
  - **Half-life:** 35 days in human fat tissue [8jwSbwLj]
- **Fumonisin B2 (IARC Group 2B):**
  - **Neurotoxicity:** Nervous system damage
  - **Carcinogenicity:** Linked to esophageal cancer
  - **Bioaccumulation:** Found in biomass used for animal feed [zwKeGzGB]

Immunosuppression:

- Mycotoxins cause **immune system alterations** [d6DardE5]
- **Increased susceptibility** to infections
- **Reduced vaccine efficacy**

Reproductive Health:

- **Estrogenicity:** Hormonal disruptions
- **Developmental toxicity:** Fetal development concerns

IBD & Additive Correlations

Emerging Evidence:

- **Carrageenan (E412):**
  - **77.5% population exposure** (French cohort) [81QWsm5y]
  - **Linked to intestinal inflammation** in animal and in vitro studies
  - **Potential trigger** for IBD flare-ups
- **Emulsifiers (General):**
  - **Disrupt gut microbiome** (70.1% population exposure to phosphates) [81QWsm5y]
  - **Alter gut barrier function**
  - **Promote inflammation**

Mechanisms:

1. **Direct irritation** of intestinal lining
2. **Microbiome dysbiosis** (imbalance of gut bacteria)
3. **Immune system activation** (chronic inflammation)
4. **Increased intestinal permeability** ("leaky gut")

4.6 Historical Context & Regulatory Failures

GRAS System Timeline

Year	Event	Impact
1958	Food Additives Amendment	GRAS system established
1958	E330, E320, E321, E924, E412 granted GRAS	**No mycotoxin testing** for E330
1961	E412 (Carrageenan) GRAS	Based on "natural source" claim
1970	Mycotoxins discovered in *A. niger*	OTA, Fumonisin identified
1970	E216 (Propyl Paraben) GRAS	Based on "tested on animals"
1980s	Animal studies show carcinogenicity	BHA, BHT, Potassium Bromate
2011	Frisvad et al. study	**83% of industrial *A. niger* produce Fumonisin B2**
2012	IARC classification	BHA, BHT, Potassium Bromate = Group 2B
2020	EFSA report	Ochratoxin A in citric acid confirmed
2021	Nature Scientific Reports	90% population exposed to E330, E412
2023	Global IBD study	5 million UC cases worldwide
2024	FDA review	BHA under review (50+ years after GRAS)
2025	Current status	**No regulatory changes** to GRAS system

Regulatory Agency Positions

FDA (United States):

- **GRAS Notification Program:** Industry self-certification
- **No mandatory testing** for mycotoxins in food additives
- **No re-evaluation** of 1958 GRAS substances
- **BHA:** Under review (2024) - 50+ years after initial approval
- **BHT, Potassium Bromate:** Still GRAS despite bans elsewhere

EFSA (European Union):

- **Mycotoxin monitoring** implemented
- **Restrictions** on BHA, BHT, Carrageenan
- **Bans** on Potassium Bromate
- **No maximum limits** for OTA/Fumonisin in citric acid
- **Pomanjkljivo spremljanje** (lack of monitoring) cited in 2020 report

WHO/IARC:

- **Ochratoxin A:** Group 2B (possibly carcinogenic)
- **Fumonisin B2:** Group 2B (possibly carcinogenic)
- **BHA, BHT, Potassium Bromate:** Group 2B (possibly carcinogenic)
- **No safe level** established for carcinogens

---

5. How to Prevent the Fraud?

For Patients

1. Read labels - Avoid E330, E320, E321, E924, E412
2. Buy organic - Lower contamination risk
3. Choose fresh - Fewer additives
4. Cook yourself - Control over ingredients
5. Get tested - MycoTOX test (urine)

For Doctors

1. Recommend testing - MycoTOX, GI-MAP
2. Educate patients - About E330 dangers
3. Collaborate with functional medicine doctors - Comprehensive evaluation
4. Demand changes - Letter to FDA/EFSA
5. Research - Follow new studies

For Activists

1. Share information - Expose GRAS fraud
2. Sign petitions - For stricter regulations
3. Contact media - Reveal the truth
4. Collaborate with researchers - Independent studies

5. Create pressure - On politicians and industry

Additional Resources

Mednarodni

- WHO – mikotoksini: https://www.who.int/foodsafety/chem/mycotoxins/en/
- EFSA – seznam GRAS: https://www.efsa.europa.eu/en/topics/topic/gras
- FDA – baza GRAS: https://www.fda.gov/food/food-ingredients-packaging/generally-recognized-safe-gras
- EWG – 12 najslabših aditivov: https://www.ewg.org/foodnews/dirty-dozen.php

Slovenia

- NIJZ – aditivi: https://www.nijz.si/
- Ministrstvo za zdravje: https://www.mz.gov.si/
- Agencija za hrano: https://www.ahs.gov.si/

Conclusion

For UC Patients

Unsafe foods:

- Coffee, alcohol, nuts, grains, dairy, processed foods
- Anything containing E330, E320, E321, E924, E412

Safe foods:

- White rice, chicken, pumpkin, bananas, avocado, coconut milk

GRAS Fraud Exposure

GRAS status is a fraud that allows **toxic substances** in food:

- E330 (citric acid) - Contains mycotoxins (OTA, FB2)
- BHA/BHT - Carcinogenic (banned in EU)
- Potassium bromate - Carcinogenic (banned in EU)
- Carrageenan - Causes intestinal inflammation

Solution:

- Read labels
- Buy organic
- Avoid additives
- Test for mycotoxins
- Demand change

Food safety is a myth. The truth is that the system is corrupted. But YOU have the power to change it.

References & Citations

Scientific Studies

[zwKeGzGB] Frisvad et al. (2011) - Fumonisin and ochratoxin production in industrial Aspergillus niger strains - PubMed [81QWsm5y] Scientific Reports (2021) - Exposure to food additive mixtures in 106,000 French adults from the NutriNet-Santé cohort [YaDFxmEt] European Environment Agency (2025) - Mycotoxin exposure in a changing European climate [qg0xc5CK] Bulletin of Biological and Allied Sciences Research (2025) - Human health risks of mycotoxins [d6DardE5] Foods Journal (2025) - Dietary mycotoxin exposure and human health risks [8jwSbwLj] PMC (2014) - Ochratoxin A and human health risk: A review [46bxbJv1] PLOS One - Process evaluation for enhanced production of aflatoxin-free citric acid [UxEk3qxV] Frontiers in Fungal Biology (2025) - Fumonisin and ochratoxin-producing strains of Aspergillus

Population Statistics

[jk2TmlAP] PubMed (2024) - Global prevalence and incidence of inflammatory bowel disease [wFci7aN0] Gastroenterology Advisor (2023) - IBD Statistics [uFZGVIcS] Gastroenterology Advisor (2023) - Ulcerative Colitis Statistics [TuqJ2vyN] ScienceDirect (2017) - Worldwide incidence and prevalence of inflammatory bowel disease [IhweW2jc] Inflammatory Bowel Diseases Journal (2024) - Trends in IBD incidence and prevalence [5jdxyTzQ] Nature (2025) - Global evolution of inflammatory bowel disease across epidemiologic stages

Regulatory Information

[gDgZyNpM] AdditiveChecker - Food additive safety database [irdkyUti] The Daily Meal - American foods banned in other countries [hIHbrq0p] BannedPantry - BHA regulatory status [pwZYQIjY] IngrediCheck - BHA regulatory review

---