

Relaxation and Odd-Even Effect in Fermi Liquids

Kittipat Pongarunotai, Puripat Thumbanthu

Outline

1

Introduction

- k -space
- Fermi surface

2

Recovering equations

- Linearization of the collision integral

3

Odd-Even effect

1

Introduction

1.1

***k*-space**

k -space

- Analogous to momentum space
- Each point $\mathbf{k}$ in k -space corresponds to a wave $e^{i\mathbf{k}}$
- **Fourier transform** converts between k -space to real space




Figure 1: REPRESENTATIVE WAVES OF k -SPACE

- Distance from origin $\propto$ Wave frequency
- Direction on k -space $\implies$ Wave direction

1.2

Fermi surface

2

Recovering equations

2.1

Linearization of the collision integral

Ideas

- Consider **only perturbation near Fermi's equilibrium**

$$f(t, \mathbf{p}) = f_0(\mathbf{p}) + \delta f(t, \mathbf{p})$$

- Transform the integrals into a linear operator

Preparation

Fermi equilibrium

$$f_0(\mathbf{p}) = \frac{1}{\exp[\beta(\varepsilon(\mathbf{p}) - \mu)] + 1}$$

1

$$\frac{\partial f_0}{\partial \varepsilon} = -\frac{1}{T} f_0(1 - f_0)$$

2

$$\frac{\partial f_0}{\partial \mu} = \frac{1}{T} f_0(1 - f_0)$$

3

$$\frac{\partial f_0}{\partial \varepsilon} = -\frac{\partial f_0}{\partial \mu}$$

4

$$f(t, \mathbf{p}) = f_0(\mathbf{p}) + \delta f(t, \mathbf{p})$$

Perturbation

5

$$\approx f_0(\mathbf{p}) + \frac{\partial f_0}{\partial \mu} \delta \mu$$

Taylor series. Fix ε

6

$$\therefore \delta f(t, \mathbf{p}) = \frac{\partial f_0}{\partial \mu} \delta \mu$$

7

$$f(t, \mathbf{p}) = f_0(\mathbf{p}) + \delta f(t, \mathbf{p})$$

Perturbation

5

$$\approx f_0(\mathbf{p}) + \frac{\partial f_0}{\partial \mu} \delta \mu$$

Taylor series. Fix ε

6

$$\therefore \delta f(t, \mathbf{p}) = \frac{\partial f_0}{\partial \mu} \delta \mu$$

7

Perturbation of interest

$$\delta \mu = T \psi$$

8

$\delta \mu$ represents temperature independent change in chemical potential

$$\delta f(t, \mathbf{p}) = \frac{\partial f_0}{\partial \mu} T \psi \quad 9$$

$$= \left(-\frac{1}{T} f_0 (1 - f_0) \right) \times T \psi \quad 10$$

$$= f_0 (1 - f_0) \psi \quad 11$$

Collision integral

$$\hat{\mathcal{J}}[f] = - \int \dots \int \frac{1}{(2\pi)^6} d\mathbf{p}_2 d\mathbf{p}'_1 d\mathbf{p}'_2 W(\mathbf{p}'_1, \mathbf{p}'_2 | \mathbf{p}_1, \mathbf{p}_2) \\ [f(\mathbf{p}_1)f(\mathbf{p}_2)(1 - f(\mathbf{p}'_1))(1 - f(\mathbf{p}'_2)) - f(\mathbf{p}'_1)f(\mathbf{p}'_2)(1 - f(\mathbf{p}_1))(1 - f(\mathbf{p}_2))] \quad 12$$

3

Odd-Even effect