

The study of the dynamics of quantum bipartite entangled systems using Feynman path integrals

Puripat Thumbanthu

Advisor: Assoc. Prof. Ekapong Hirunsirisawat, Dr. Tanapat Deesuwan

Started: May 16 2024

This version is not to be published. It is my own notes during the research.

Contents

1	Introduction	4
1.1	Historical backgrounds	4
1.2	Academical backgrounds	5
1.3	List of symbols	5
2	The generalized Gaussian integrals	6
2.1	Preliminary form	6
2.2	Evaluation	6
2.3	Tables of Gaussian integrals with varying orders	8
3	Propagator and joint probability distribution	10
3.1	General theory for the propagator of two particles	10
3.1.1	Non-interacting systems	10
3.1.2	Interacting systems	11
3.1.3	Form of the two particle perturbation series	12
3.2	Initial states of choice	13

3.3	The spring system	13
3.3.1	First order perturbation term	13
3.3.2	Joint probability contribution from the first order perturbation term on a separable state	16
3.3.3	Joint probability contribution from the first order perturbation term on an entangled state	18
3.3.4	Second order perturbation term	18
3.4	The delta function collision problem	19
3.4.1	First order perturbation term	19
3.5	The coulomb problem	19
4	Correlation function	20
5	Implementation of simulations	22
6	The Schrödinger's equation for two particles	26
A	Integrals evaluation code	28
A.1	Polynomial extraction	28
A.2	Code for the spring problem	29
A.2.1	First order perturbation	29

Introduction

1.1 Historical backgrounds¹

I'm captured by the path integrals method of quantum mechanics, and its ability to intuitively produce the classical world, which is governed by the principle of least action. It appears so naturally in path integrals that I thought I was hallucinating. I wasn't. However, one question arises in my mind: what does the principle of least action look like for more than one particle? And what if they're entangled? I was literally mashing random words together at that point, but it has led me into the realm of physics that I don't think anyone really cares before: the dynamics of an entangled system.

Entanglement is usually defined as the inseparability of states, commonly used to describe the spin of a particle, which has been studied in great extent. It has found its use in quantum computer, giving its blazing speed. However, from my knowledge, no literature has mentioned the effect of entanglement on the movement of particles. Will entangled particles move together in the same direction? If I separate a pair of entangled particles, put one in a potential, and let the other be free, does the particle in the potential have any effect on the free particle? And thus, this research (*project*) was born.

This research started out by two people; me, and the other one that shall not be named. However, I have a lot of conflicts with that person. Long story short, he quitted; and thus, I'm on my own journey.

¹This introduction is rewritten after the school project has passed. I've declared this project to be independent of the school ever since. So, it's going to be much more authentic and straightforward.

1.2 Academical backgrounds

1.3 List of symbols

The generalized Gaussian integrals

2.1 Preliminary form

A Gaussian integral is the integral of the Gaussian function $\exp[-x^2]$:

$$\int_{-\infty}^{\infty} \exp[-x^2] dx = \sqrt{\pi}. \quad (2.1)$$

However, this is not that useful for path integrals, because it demands a more generalized form of this Gaussian function. Therefore, we focus on integrals of the form

$$I_1 = \int_{-\infty}^{\infty} x^n \exp[-(ax^2 + bx + c)] dx, \quad (2.2)$$

where a , b , and c are constants and n is a positive integer. This integral will be very useful later on when evaluating the Born expansion of the propagator.

Because we're just going to be dealing with integrals from negative to positive infinity, I shall omit the bounds of the integrals. Thus, every integral written with no bounds from now on are assumed to be a definite integral from negative to positive infinity.

2.2 Evaluation

To evaluate I_1 (eq. (2.2)), we shall complete the square first, then substitute the exponents to fit the form of the standard Gaussian integral.

$$\begin{aligned} \int x^n e^{-(ax^2+bx+c)} dx &= \int x^n \exp\left[-a\left(x + \frac{b}{2a}\right)^2 + \frac{b^2}{4a} - c\right] dx \\ &= e^{\frac{b^2}{4a}-c} \int x^n \exp\left[-a\left(x + \frac{b}{2a}\right)^2\right] dx. \end{aligned}$$

Let

$$-u^2 = -a\left(x + \frac{b}{2a}\right)^2 \quad (2.3)$$

$$x = \frac{u}{\sqrt{a}} - \frac{b^2}{2a} \quad (2.4)$$

$$\frac{dx}{du} = \frac{d}{du} \left(\frac{u}{\sqrt{a}} - \frac{b^2}{2a} \right) \quad (2.5)$$

$$dx = \frac{1}{\sqrt{a}} du. \quad (2.6)$$

Then,

$$I_1 = e^{\frac{b^2}{4a}-c} \frac{1}{\sqrt{a}} \int \left(\frac{u}{\sqrt{a}} - \frac{b^2}{2a} \right)^n e^{-u^2} du. \quad (2.7)$$

$$= e^{\frac{b^2}{4a}-c} \frac{1}{\sqrt{a}} \int \sum_{k=0}^n \binom{n}{k} \left(\frac{b^2}{2a} \right)^{n-k} \left(\frac{u}{\sqrt{a}} \right)^k e^{-u^2} du. \quad (2.8)$$

$$= e^{\frac{b^2}{4a}-c} \frac{1}{\sqrt{a}} \sum_{k=0}^n \left[\binom{n}{k} \left(\frac{b^2}{2a} \right)^{n-k} \left(\frac{1}{\sqrt{a}} \right)^k \int u^k e^{u^2} du \right], \quad (2.9)$$

i.e., I_1 can be written as a sum of

$$I_0 = \int u^k e^{-u^2} du. \quad (2.10)$$

Notice that when n is odd, the integrand of I_0 is an odd function. Therefore, I_0 is zero whenever n is odd. When n is even however, the integrand is even. Thus, it can be simplified to

$$2 \int_0^\infty u^{2m} e^{-u^2} du \quad (2.11)$$

where $k = 2m$. We then do another substitution by letting $-t = -u^2$. Thus, $u = \sqrt{t}$ and $du = dt/2\sqrt{t}$. Both infinity and zero aren't affected by a square root, therefore the bound doesn't change.

Our integral then becomes

$$2 \times \int_0^\infty t^m e^{-t} \frac{1}{2\sqrt{t}} dt. \quad (2.12)$$

$$= \int_0^\infty t^{m-\frac{1}{2}} e^{-t} dt \quad (2.13)$$

$$= \int_0^\infty t^{m+\frac{1}{2}-1} e^{-t} dt \quad (2.14)$$

$$= \Gamma \left(m + \frac{1}{2} \right) \quad (2.15)$$

$$= \Gamma \left(\frac{k+1}{2} \right). \quad (2.16)$$

Since the integral evaluates to the gamma function for only even numbers, we add a term that makes $I_0 = 0$ when n is odd:

$$I_0 = \frac{1}{2}((-1)^k + 1) \Gamma \left(\frac{k+1}{2} \right). \quad (2.17)$$

Thus, from eq. (2.9),

$$I_1 = e^{\frac{b^2}{4a}-c} \frac{1}{\sqrt{a}} \sum_{k=0}^n \left[\binom{n}{k} \left(\frac{b^2}{2a} \right)^{n-k} \left(\frac{1}{\sqrt{a}} \right)^k \frac{1}{2}((-1)^k + 1) \Gamma \left(\frac{k+1}{2} \right) \right] \quad (2.18)$$

To continue this, we then expand the combinatorics and use the formulas of arguments with half-integer real part to get

$$e^{\frac{b^2}{4a}-c} \frac{1}{\sqrt{a}} \sum_{k=0}^n \frac{n!}{k!(n-k)!} \left(\frac{b^2}{2a}\right)^n \left(\frac{2a}{b^2}\right)^k \left(\frac{1}{\sqrt{a}}\right)^k \left(\sqrt{\pi} \frac{k!}{4^{\frac{k}{2}} (k/2)!}\right) \times \frac{1}{2}((-1)^k + 1) \quad (2.19)$$

$$= e^{\frac{b^2}{4a}-c} n! \sqrt{\frac{\pi}{a}} \left(\frac{b^2}{2a}\right)^n \sum_{k=0}^n \frac{1}{(n-k)!(k/2)!} \left(\frac{\sqrt{a}}{b^2}\right)^k \frac{1}{2}((-1)^k + 1) \quad (2.20)$$

To simplify the summation, we replace k with $2k$, and the upper limit with m where $m = \lfloor n/2 \rfloor$.

Thus, we get

$$\left[\sum_{k=0}^m \frac{1}{(n-2k)!k!} \left(\frac{\sqrt{a}}{b^2}\right)^{2k} \right] e^{\frac{b^2}{4a}-c} n! \left(\frac{b^2}{2a}\right)^n \sqrt{\frac{\pi}{a}} \quad (2.21)$$

As far as I know, this equation cannot be simplified further.

Comment on the generalized Gaussian integral Note that the integral in eq. (2.2) can be casted in the form of Meijer's G function for $n \geq 0$:

$$\int x^n \exp[-ax^2 - bx - c] dx = \frac{e^{-c}}{2\sqrt{\pi}ab} \left[-\frac{2^{n+1}a}{b^n} G_{2,1}^{1,2} \left(\frac{1-n}{2}, -\frac{n}{2} \middle| \frac{4ae^{-2i\pi}}{b^2} \right) + a^{\frac{1-n}{2}} b G_{1,2}^{2,1} \left(\frac{1-n}{2} \middle| \frac{b^2}{4a} \right) \right] \quad (2.22)$$

However, this form doesn't seem to be very helpful when we try to evaluate the polynomial part of the Gaussian integral because of the nature of the Meijer's G function that's defined with integral of products.

Originally, I was going to evaluate everything using `SymPy`, and not really care about anything else. However, I came to realize that `SymPy` doesn't bother integrating any Gaussian integrals that have complex number components. My own implementation will be written in chapter 5

2.3 Tables of Gaussian integrals with varying orders

$$n = 0: +\frac{1}{1} \sqrt{\frac{\pi}{a}} e^{\frac{b^2}{4a}-c}$$

$$n = 1: -\frac{b}{2} \sqrt{\frac{\pi}{a^3}} e^{\frac{b^2}{4a}-c}$$

$$n = 2: +\frac{1}{4} \sqrt{\frac{\pi}{a^5}} (2a + b^2) e^{\frac{b^2}{4a}-c}$$

$$n = 3: -\frac{b}{8} \sqrt{\frac{\pi}{a^7}} (6a + b^2) e^{\frac{b^2}{4a}-c}$$

$$n = 4: +\frac{1}{16} \sqrt{\frac{\pi}{a^9}} (12a^2 + 12ab^2 + b^4) e^{\frac{b^2}{4a}-c}$$

$$n = 5: -\frac{b}{32} \sqrt{\frac{\pi}{a^{11}}} (60a^2 + 20ab^2 + b^2) e^{\frac{b^2}{4a} - c}$$

$$n = 6: +\frac{1}{64} \sqrt{\frac{\pi}{a^{13}}} (120a^3 + 180a^2b^2 + 30ab^4 + b^6) e^{\frac{b^2}{4a} - c}$$

$$n = 7: -\frac{b}{128} \sqrt{\frac{\pi}{a^{15}}} (840a^3 + 420a^2b^2 + 42ab^4 + b^6) e^{\frac{b^2}{4a} - c}$$

$$n = 8: +\frac{1}{256} \sqrt{\frac{\pi}{a^{17}}} (1680a^4 + 3360a^3b^2 + 840a^2b^4 + 56ab^6 + b^8) e^{\frac{b^2}{4a} - c}$$

$$n = 9: -\frac{b}{512} \sqrt{\frac{\pi}{a^{19}}} (15120a^4 + 10080a^3b^2 + 1512a^2b^4 + 72ab^6 + b^8) e^{\frac{b^2}{4a} - c}$$

Propagator and joint probability distribution

3.1 General theory for the propagator of two particles

3.1.1 Non-interacting systems

The propagator for a single particle system is used to find the probability distribution $\tilde{P}(x)$ of a state at anytime, i.e., for a state $|\psi(t_0)\rangle$,

$$\tilde{P}(q_F, t_F) = \langle q_F | \psi(t_F) \rangle = \int K(q_f, q_0; \Delta t) \psi(q_0, t_0) dq_0. \quad (3.1)$$

When taken the modulus squared, $|\tilde{P}(q_F, t_F)|^2$, or just $P(q_F, t_F)$, represents the probability of finding that particle at the point (q_F, t_F) in spacetime.

We shall then extend this idea to describe a two particle system. Let there be a state ket $|\eta(t)\rangle$, which simultaneously represents the state of both particle one and particle two. Assume that at some point in time, $|\eta\rangle$ is separable:

$$|\eta(t_0)\rangle = |\psi(t_0)\rangle \otimes |\phi(t_0)\rangle. \quad (3.2)$$

If the subsystem of $|\eta\rangle$ is non-interacting, the Hamiltonian must be completely separable, which also implies that the time evolution operator \hat{U} is also separable. Consider $\langle q_F, q'_F | \eta(t_F) \rangle$ where the unprimed q belongs to the position in the Hilbert space of particle one, and the primed q 's, in Hilbert space of particle two.

$$\begin{aligned} \langle q_F, q'_F | \eta(t_F) \rangle &= \langle q_F, q'_F | (|\psi(t_F)\rangle |\phi(t_F)\rangle) \\ &= \psi_i(q_F, t_F) \phi_i(q'_F, t_F) \\ &= \iint K_\psi(q_F, t_F; q_0, t_0) K_\phi(q'_F, t_F; q'_0, t'_0) [\psi_i(q_F, t_F) \phi_i(q'_F, t_F)] dq_0 dq'_0. \end{aligned}$$

For most systems, K_ψ and K_ϕ is identical because both ψ_i and ϕ_i is affected by the same potential. But for some, K_ψ and K_ϕ might not be identical. E.g., if ψ is in a potential well, but ϕ is a free particle that's infinitely far away. K_ψ must be of the potential well, and K_ϕ must be of the free particle.

It can be seen that $\langle q_F, q'_F | \eta(t_F) \rangle$ represents the probability distribution of both the two particles. E.g., for a state ket $|\psi\rangle \otimes |\phi\rangle$,

$$(\langle q_F | \langle q'_F |) (|\psi(t_F)\rangle |\phi(t'_F)\rangle) \quad (3.3)$$

$$= \langle q_F | \psi(t_F) \rangle \langle q'_F | \phi(t'_F) \rangle \quad (3.4)$$

$$= \iint dq_0 dq'_0 K(q_F, t_F; q_0, t_0) K(q'_F, t'_F; q'_0, t'_0) (\psi(q_0, t_0) \phi(q'_0, t'_0)) \quad (3.5)$$

$$= \int dq_0 K(q_F, t_F; q_0, t_0) \psi(q_0, t_0) \int dq'_0 K(q'_F, t'_F; q'_0, t'_0) \phi(q'_0, t'_0), \quad (3.6)$$

eq. (3.4) says that $\langle q_F, q'_F | \eta(t_F) \rangle$ is the product of the probability that ψ is at (q_F, t_F) , and (q'_F, t'_F) . Therefore, $\langle q_F, q'_F | \eta(t_F) \rangle$ represents the joint probability distribution of the two states, and it can be found via the product of propagators,

$$K(q_F, t_F; q_0, t_0) K(q'_F, t'_F; q'_0, t'_0). \quad (3.7)$$

3.1.2 Interacting systems

Let there be a state ket $|\eta\rangle$ which describes the quantum state of two particles. When the two particles are interacting, the time-evolution operator cannot be factored into tensor products of two operators. Therefore, there can only be one combined time-evolution operator for both of the systems:

$$\hat{U}(t_F, t_0) = \exp \left[-i\Delta t \left(\frac{\hat{p}^2}{2m} + \frac{\hat{p}'^2}{2m'} + V(\hat{q}, \hat{q}', t) \right) \right] \quad (3.8)$$

The consequence of the combined time-evolution operator is that, we cannot calculate the joint probability distribution between different time of the subsystem. For simplicity' sake, let the index $j = i + 1$, where $t_j = t_i + \Delta t$ in which $\Delta t \rightarrow 0$.

$$\langle q_j, q'_j | \eta(t_j) \rangle \quad (3.9)$$

$$= \langle q_j, q'_j | \hat{U}(t_j, t_i) | \eta(t_i) \rangle. \quad (3.10)$$

$$= \iint dq_i dq'_i \langle q_j, q'_j | \hat{U}(t_j, t_i) | q_i, q'_i \rangle \langle q_i, q'_i | \eta(t_i) \rangle. \quad (3.11)$$

As seen, the form of transition element is

$$\langle q_j, q'_j | \exp \left[-i\Delta t \left(\frac{\hat{p}^2}{2m} + \frac{\hat{p}'^2}{2m'} + V(\hat{q}, \hat{q}', t) \right) \right] | q_i, q'_i \rangle. \quad (3.12)$$

The terms in the exponents can be separated due to the vanishing commutator when $\Delta t \rightarrow 0$. By separating the terms in the exponents and inserting two complete sets of momentum basis, the

equation above turns into

$$\begin{aligned}
 & \frac{1}{(2\pi)^2} \iint dp dp' \langle q_j, q'_j | \exp \left[-i\Delta t \frac{\hat{p}^2}{2m} \right] | p, p' \rangle \langle p, p' | \exp \left[-i\Delta t \frac{\hat{p}'^2}{2m'} \right] \exp[-i\Delta t V(\hat{q}, \hat{q}', t)] | q_i, q'_i \rangle \\
 &= \frac{1}{(2\pi)^2} \iint dp dp' \exp \left[-iu\Delta t \left(\frac{p^2}{2m} + \frac{p'^2}{2m'} + V(q_i, q_i, \Delta t) \right) \right] \langle q_j | p \rangle \langle q'_j | p' \rangle \langle q_i | p \rangle^* \langle q'_i | p' \rangle^* \\
 &= \frac{1}{(2\pi)^4} e^{-i\Delta t V(q_i, q'_i, t)} \int dp \exp \left[-i\Delta t \frac{p^2}{2m} + ip(q_j - q_i) \right] \int dp' \exp \left[-i\Delta t \frac{p'^2}{2m'} + ip'(q'_j - q'_i) \right] \\
 &= \frac{(mm')^{\frac{1}{2}}}{8\pi^3 i \Delta t} \exp \left[-i\Delta t V(q_i, q'_i, t) + \frac{im'}{2\Delta t} (q'_j - q'_i)^2 + \frac{im}{2\Delta t} (q_j - q_i)^2 \right]. \tag{3.13}
 \end{aligned}$$

To find the propagator for an interacting system, we need to perform successive integrals on q and q' , i.e.,

$$K_\eta = \int \dots \int dq_N dq'_N \dots dq_1 dq'_1 \langle q_F, q'_F | \hat{U}(t_F, t_N) | q_N, q'_N \rangle \dots \langle q_1, q'_1 | \hat{U}(t_1, t_0) | q_0, q'_0 \rangle \tag{3.14}$$

Notice that when there is no interaction between the two systems ($V = 0$), the integrals become separable and reduces down to the form of the non-interacting system's propagator but off by a normalization factor.

There are two common forms of interaction, which is the spring interaction and the coulomb interaction. Both of which includes the term $(q_i - q'_i)^2$ in $V(q, q')$, which causes major problems in integration. When expanded, there is a $q_i q'_i$ term that makes the integral inseparable which causes the integral pattern to not repeat; therefore, we resort to perturbation.

3.1.3 Form of the two particle perturbation series

The perturbation series for one particle are already given by Feynman in his path integrals textbook:

$$K_n(F, 0) = (-i)^n \int \dots \int K_0(F, n) \prod_{i=1}^n V(i) K_0(i, i-1) d\tau_i \tag{3.15}$$

where K_n is the n 'th order perturbation, $K(j, k) = K(q_j, t_j; q_k, t_k)$, $K = \sum_n K_n$ and $d\tau_i = dq_i dq'_i dt_i$. To apply it with two particles, we extend it:

$$K_n(F, 0; F', 0') = (-i)^n \int \dots \int K_0(F, n) K_0(F', n') \left[\prod_{i=1}^n V(i) K_0(i, i-1) K_0(i', i'-1) dq_i dq'_i dt_i \right]. \tag{3.16}$$

We can also expand the joint probability distribution in the form of the perturbation series.

$$\eta(q_F, q'_F; t_F) = \int K(q_F, q'_F; q_0, q'_0; t_F - t_0) \eta(q_0, q'_0; t_0) dq_0 dq'_0 \tag{3.17}$$

$$= \int \sum_n K_n(q_F, q'_F; q_0, q'_0; t_F - t_0) \eta(q_0, q'_0; t_0) dq_0 dq'_0 \tag{3.18}$$

$$= \sum_n \int K_n(q_F, q'_F; q_0, q'_0; t_F - t_0) \eta(q_0, q'_0; t_0) dq_0 dq'_0 \tag{3.19}$$

3.2 Initial states of choice

We're interested in the time evolution of a Gaussian wave packet, mostly due to its mathematical simplicity. The normalized state of a Gaussian wave packet is given by

$$\sqrt{\frac{1}{2\pi\sigma}} \exp\left[ipq - \frac{(q-s)^2}{4\sigma^2}\right]. \quad (3.20)$$

For two separable particles, its joint probability is

$$\langle q, q' | \eta(t_0) \rangle = \sqrt{\frac{1}{2\pi\sigma}} \sqrt{\frac{1}{2\pi\sigma'}} \exp\left[ipq - \frac{(q-s)^2}{4\sigma^2}\right] \exp\left[ip'q' - \frac{(q'-s')^2}{4\sigma'^2}\right] \quad (3.21)$$

As we'll see, the higher order perturbation terms (at least of the spring system) are expressed as the integral of the product between the wave function and some polynomials, which we'll find later.

$$\langle q | \xi(t) \rangle \quad (3.22)$$

$$= \int K_0(q_F, q_0; t_F) \langle q_0 | \xi(t) \rangle dq_0 \quad (3.23)$$

$$= \sqrt{\frac{m}{2\pi i t_F}} \int \exp\left[\frac{im}{2t_F}(q_F - q_0)^2\right] \left(\sqrt{\frac{1}{2\pi\sigma}} \exp\left[ipq_0 - \frac{(q_0-s)^2}{4\sigma^2}\right] \right) dq_0. \quad (3.24)$$

$$= \frac{1}{2\pi\sqrt{\sigma\sigma'}} \exp\left[ipq - \frac{(q-s)^2}{4\sigma^2}\right] \exp\left[ip'q' - \frac{(q'-s')^2}{4\sigma'^2}\right] \quad (3.25)$$

$$= \frac{1}{2\pi} \sqrt{\frac{m}{it_F\sigma}} \int \exp\left[\frac{im}{2t_F}(q_F - q_0)^2 + ipq_0 - \frac{(q_0-s)^2}{4\sigma^2}\right] dq_0 \quad (3.26)$$

$$= \frac{1}{2\pi} \sqrt{\frac{m}{it_F\sigma}} \int \exp\left[\left(\frac{imq_F^2}{2t_F} - \frac{s^2}{4\sigma^2}\right) + q\left(-\frac{imq_F}{t_F} + ip + \frac{s}{2\sigma^2}\right) + q^2\left(\frac{im}{2t_F} - \frac{1}{4\sigma^2}\right)\right] dq_0 \quad (3.27)$$

$$= \frac{1}{2\pi} \sqrt{\frac{m}{it_F\sigma}} \cdot 2\sigma \sqrt{\frac{\pi t_F}{-2im\sigma^2 + t_F}} \exp\left[\frac{4impq_F\sigma^2 - mq_F^2 + 2mq_Fs - ms^2 - 2ip^2t_F\sigma^2 - 2pst_F}{4m\sigma^2 + 2it_F}\right] \quad (3.28)$$

$$= \sqrt{\frac{m\sigma}{2m\sigma^2 + it_F}} \exp\left[\frac{4impq_F\sigma^2 - mq_F^2 + 2mq_Fs - ms^2 - 2ip^2t_F\sigma^2 - 2pst_F}{4m\sigma^2 + 2it_F}\right]. \quad (3.28)$$

To plot its probability amplitude, we have to take its modulus squared, which equals to

$$\frac{m\sigma}{\sqrt{4m^2\sigma^4 + t_F^2}} \exp\left[\frac{-2m^2q_F^2\sigma^2 + 4m^2q_Fs\sigma^2 - 2m^2s^2\sigma^2 + 4mpq_Ft_F\sigma^2 - 4mpst_F\sigma^2 - 2p^2t_F^2\sigma^2}{4m^2\sigma^4 + t_F^2}\right] \quad (3.29)$$

3.3 The spring system

The spring potential is given by $V(q, q') = \frac{k}{2}(q - q')^2$, which I shall let $\alpha = \frac{k}{2}$ for simplicity's sake.

3.3.1 First order perturbation term

From eq. (3.16), set $t_1 = t, t_0 = 0$

$$K_1(F, 0; F', 0)$$

$$\begin{aligned}
 &= (-i)^1 \iiint K_0(F, 1)K_0(F', 1')K_0(1, 0)K_0(1, 0')\alpha(q_1 - q'_1)^2 dq_1 dq'_1 dt. \quad (3.30) \\
 &= -i\alpha \int_0^{t_F} \left[\iint K_0(q_F, q_1; t_F - t)K_0(q'_F, q'_1; t_F - t)K_0(q_1, q_0; t)K_0(q'_1, q'_0; t)(q_1 - q'_1)^2 dq_1 dq'_1 \right] dt.
 \end{aligned}$$

We then let the terms in the square bracket,

$$\iint K_0(q_F, q_1; t_F - t)K_0(q'_F, q'_1; t_F - t)K_0(q_1, q_0; t)K_0(q'_1, q'_0; t)(q_1 - q'_1)^2 dq_1 dq'_1 \quad (3.31)$$

equals I ; therefore, $K_1 = -i\alpha \int_0^{t_F} I dt$.

We then separate I into three integrals:

$$I_{P1} = \int q_1^2 K_0(q_F, q_1; t_F - t)K_0(q_1, q_0; t) dq_1 \int K_0(q'_F, q'_1; t_F - t)K_0(q'_1, q'_0; t) dq'_1, \quad (3.32)$$

$$I_{P2} = \int K_0(q_F, q_1; t_F - t)K_0(q_1, q_0; t) dq_1 \int q_1^2 K_0(q'_F, q'_1; t_F - t)K_0(q'_1, q'_0; t) dq'_1, \quad (3.33)$$

$$I_{P3} = \int q_1 K_0(q_F, q_1; t_F - t)K_0(q_1, q_0; t) dq_1 \int q'_1 K_0(q'_F, q'_1; t_F - t)K_0(q'_1, q'_0; t) dq'_1, \quad (3.34)$$

where $I = I_{P1} + I_{P2} + 2I_{P3}$. The integrals without the factor q_1 and q_1^2 can be reduced into the kernel for the free particle:

$$I_{P1} = K_0(q'_F, q'_0; t_F) \int q_1^2 K_0(q_F, q_1; t_F - t)K_0(q_1, q_0; t) dq_1 \quad (3.35)$$

$$I_{P2} = K_0(q_F, q_0; t_F) \int q_1^2 K_0(q'_F, q'_1; t_F - t)K_0(q'_1, q'_0; t) dq'_1 \quad (3.36)$$

Since I_{P2} can be obtained by switching all the primed variables with the corresponding unprimed in I_{P1} , we're left with two family of integrals:

$$\int q_1 K_0(q_F, q_1; t_F - t)K_0(q_1, q_0; t) dq_1 \quad \text{and} \quad \int q_1^2 K_0(q_F, q_1; t_F - t)K_0(q_1, q_0; t) dq_1. \quad (3.37)$$

To evaluate these, we first simplify the product of kernel under the assumption that $t_F > t$.

$$\begin{aligned}
 &K_0(q_F, q_1; t_F - t)K_0(q_1, q_0; t) \\
 &= \sqrt{\frac{m}{2\pi i(t_F - t)}} \sqrt{\frac{m}{2\pi i t}} \exp \left[\frac{im}{2(t_F - t)}(q_F - q_1)^2 + \frac{im}{2t}(q_1 - q_0)^2 \right] \quad (3.38) \\
 &= \frac{m}{2\pi} \sqrt{-\frac{1}{t(t_F - t)}} \exp \left[q_1^2 \left(\frac{im}{2t} + \frac{im}{2(t_F - t)} \right) - q_1 \left(\frac{imq_0}{t} + \frac{imq_F}{t_F - t} \right) + \left(\frac{imq_0^2}{2t} + \frac{imq_F^2}{2(t_F - t)} \right) \right] \\
 &= \frac{m}{2\pi} \sqrt{-\frac{1}{t(t_F - t)}} \exp \left[-q_1^2 \left(\frac{mt_F}{2it(t_F - t)} \right) - q_1(im) \left(\frac{q_0}{t} + \frac{q_F}{t_F - t} \right) - \left(\frac{m}{2i} \right) \left(\frac{q_0^2}{t} + \frac{q_F^2}{t_F - t} \right) \right]
 \end{aligned}$$

The normalization factor are pulled out. Both integrals in eq. (3.37) can be evaluated with

$$a = \frac{mt_F}{2it(t_F - t)}, \quad b = im \left(\frac{q_0}{t} + \frac{q_F}{t_F - t} \right) \quad \text{and} \quad c = \left(\frac{m}{2i} \right) \left(\frac{q_0^2}{t} + \frac{q_F^2}{t_F - t} \right) \quad (3.39)$$

in which,

$$\exp \left[\frac{b^2}{4a} - c \right] = \exp \left[\frac{im}{2t_F}(q_F - q_0)^2 \right] = \sqrt{\frac{2\pi i t_F}{m}} K_0(q_F, q_0; t_F). \quad (3.40)$$

To summarize,

$$\begin{aligned} \int q_1 K_0(q_F, q_1; t_F - t) K_0(q_1, q_0; t) dq_1 &= -\frac{b}{2} \sqrt{\frac{\pi}{a^3}} \times \frac{m}{2\pi} \sqrt{-\frac{1}{t(t_F - t)}} \times \sqrt{\frac{2\pi i t_F}{m}} K_0(q_F, q_0; t_F) \\ &= -\frac{b}{2} \sqrt{\frac{\pi}{a^3}} \times \sqrt{\frac{m t_F}{2\pi i t(t_F - t)}} K_0(q_F, q_0; t_F), \end{aligned} \quad (3.41)$$

and

$$\int q_1^2 K_0(q_F, q_1; t_F - t) K_0(q_1, q_0; t) dq_1 = \frac{1}{4} \sqrt{\frac{\pi}{a^5}} (2a + b^2) \times \sqrt{\frac{m t_F}{2\pi i t(t_F - t)}} K_0(q_F, q_0; t_F). \quad (3.42)$$

On the Gaussian integral with degree one,

$$-\frac{b}{2} \sqrt{\frac{\pi}{a^3}} = \sqrt{\frac{2\pi t}{m t_F^3}} \frac{\sqrt{-i(t_F - t)^3}}{i(t_F - t)} \times [q_0(t_F - t) + q_F t]; \quad (3.43)$$

thus from eq. (3.41),

$$\int q_1 K_0(q_F, q_1; t_F - t) K_0(q_1, q_0; t) = \sqrt{\frac{2\pi t}{m t_F^3}} \frac{\sqrt{-i(t_F - t)^3}}{i(t_F - t)} \times [q_0(t_F - t) + q_F t] \quad (3.44)$$

$$\begin{aligned} &\times \sqrt{\frac{m t_F}{2\pi i t(t_F - t)}} K_0(q_F, q_0; t_F) \\ &= -\frac{1}{t_F} [q_0(t_F - t) + q_F t] K_0(q_F, q_0; t_F). \end{aligned} \quad (3.45)$$

On the Gaussian integral with degree two,

$$\frac{1}{4} \sqrt{\frac{\pi}{a^5}} (2a + b^2) = -\sqrt{\frac{2\pi t}{m^3 t_F^5}} \frac{\sqrt{i(t_F - t)^5}}{(t_F - t)^2} [m(q_0(t_F - t) + q_F t)^2 + i t t_F (t_F - t)]; \quad (3.46)$$

and thus,

$$\int q_1^2 K_0(q_F, q_1; t_F - t) K_0(q_1, q_0; t) dq_1 \quad (3.47)$$

$$\begin{aligned} &= \sqrt{\frac{m t_F}{2\pi i t(t_F - t)}} K_0(q_F, q_0; t_F) \times -\sqrt{\frac{2\pi t}{m^3 t_F^5}} \frac{\sqrt{i(t_F - t)^5}}{(t_F - t)^2} [m(q_0(t_F - t) + q_F t)^2 + i t t_F (t_F - t)] \\ &= -\frac{1}{m t_F^2} [m(q_0(t_F - t) + q_F t)^2 + i t t_F (t_F - t)] K_0(q_F, q_0; t_F). \end{aligned} \quad (3.48)$$

We're now in the place to finally construct the first order propagator term for the spring system. Recall that

$$K_1 = -i\alpha \int_0^{t_F} I dt = -i\alpha \left[\int_0^{t_F} I_{P1} dt + \int_0^{t_F} I_{P2} dt + 2 \int_0^{t_F} I_{P3} dt \right] \quad (3.49)$$

From earlier,

$$\begin{aligned} I_{P1} &= K_0(q'_F, q'_0; t_F) \int q_1^2 K_0(q_F, q_1; t_F - t) K_0(q_1, q_0; t) dq_1 \\ &= -\frac{1}{m t_F^2} [m(q_0(t_F - t) + q_F t)^2 + i t t_F (t_F - t)] K_0(q_F, q_0; t_F) K_0(q'_F, q'_0; t_F), \end{aligned} \quad (3.50)$$

$$\begin{aligned}
 I_{P2} &= K_0(q_F, q_0; t_F) \int q_1'^2 K_0(q_F', q_1'; t_F - t) K_0(q_1', q_0'; t) dq_1' \\
 &= -\frac{1}{mt_F^2} \left[m(q_0'(t_F - t) + q_F't)^2 + itt_F(t_F - t) \right] K_0(q_F, q_0; t_F) K_0(q_F', q_0'; t_F),
 \end{aligned} \tag{3.51}$$

$$\begin{aligned}
 I_{P3} &= -\frac{1}{t_F} [q_0(t_F - t) + q_F t] K_0(q_F, q_0; t_F) \times -\frac{1}{t_F} [q_0'(t_F - t) + q_F' t] K_0(q_F', q_0'; t_F) \\
 &= \frac{1}{t_F^2} [q_0(t_F - t) + q_F t] [q_0'(t_F - t) + q_F' t] K_0(q_F, q_0; t_F) K_0(q_F', q_0'; t_F).
 \end{aligned} \tag{3.52}$$

Thus,

$$\begin{aligned}
 K_1 &= -i\alpha K_0(q_F, q_0; t_F) K_0(q_F', q_0'; t_F) \frac{1}{t_F^2} \left[-\frac{1}{m} \int_0^{t_F} m(q_0(t_F - t) + q_F t)^2 + itt_F(t_F - t) dt \right. \\
 &\quad \left. - \frac{1}{m} \int_0^{t_F} m(q_0'(t_F - t) + q_F' t)^2 + itt_F(t_F - t) dt \right. \\
 &\quad \left. + \int_0^{t_F} [q_0(t_F - t) + q_F t] [q_0'(t_F - t) + q_F' t] dt \right] \tag{3.53}
 \end{aligned}$$

The integrals are then taken out to be evaluated term by term. Since the integrand of these integrals are all polynomials, we can just plug it into `SymPy.jl`:

$$\int_0^{t_F} m(q_0(t_F - t) + q_F t)^2 dt = \frac{t_F^3}{6} (2m(q_0 + q_F)^2 + it_F) \tag{3.54}$$

$$\int_0^{t_F} m(q_0'(t_F - t) + q_F' t)^2 dt = \frac{t_F^3}{6} (2m(q_0' + q_F')^2 + it_F) \tag{3.55}$$

$$\int_0^{t_F} [q_0(t_F - t) + q_F t] [q_0'(t_F - t) + q_F' t] dt = \frac{t_F^3}{6} (2q_0 q_0' + q_0 q_F' + q_0' q_F + 2q_F q_F') \tag{3.56}$$

Therefore, the first order perturbation term of the spring system takes the form

$$\begin{aligned}
 K_1(q_F, q_0; q_F', q_0'; t_F) &= -i\frac{\alpha t_F}{6} K_0(q_F, q_0; t_F) K_0(q_F', q_0'; t_F) \\
 &\quad \times [-2(m(q_0 + q_F)^2 + m(q_0' + q_F')^2 + it_F) + 2q_0 q_0' + q_0 q_F' + q_0' q_F + 2q_F q_F']. \tag{3.57}
 \end{aligned}$$

3.3.2 Joint probability contribution from the first order perturbation term on a separable state

As said earlier, we're interested in the time evolution of a Gaussian wave packet earlier, with the initial state of the two particles being

$$\eta(q_0, q_0; t_0) = \frac{1}{2\pi\sqrt{\sigma\sigma'}} \exp \left[ipq_0 + ip'q_0' - \frac{(q_0 - s)^2}{4\sigma^2} - \frac{(q_0' - s')^2}{4\sigma'^2} \right]. \tag{3.58}$$

We shall set $t_0 = 0$ for simplicity's sake. The state at any following time t_F is then

$$\eta(q_F, q_F'; t_F) = \iint (K_0(q_F, q_0; q_F', q_0'; t_F - t_0) + K_1(q_F, q_0; q_F', q_0'; t_F - t_0)) \eta(q_0, q_0'; t_0) dq dq_0' \tag{3.59}$$

Thus, using the first order perturbation term from eq. (3.57),

$$\begin{aligned} \eta(q_F, q'_F; t_F) = \frac{1}{2\pi\sqrt{\sigma\sigma'}} \iint K_0(q_F, q_0; q'_F, q'_0; t_F - t_0) & \left\{ 1 - i\frac{\alpha t_F}{6} \left[-2(m(q_0 + q_F)^2 \right. \right. \\ & \left. \left. + m(q'_0 + q'_F)^2 + it_F) + 2q_0q'_0 + q_0q'_F + q'_0q_F + 2q_Fq'_F \right] \right\} \\ & \exp\left[ipq_0 + ip'q'_0 - \frac{(q_0 - s)^2}{4\sigma^2} - \frac{(q'_0 - s')^2}{4\sigma'^2} \right] dq_0 dq'_0 \end{aligned} \quad (3.60)$$

In the end, this integral can be evaluated term by term as Gaussian integrals with the same exponential argument.

$$ipq_0 - \frac{(q_0 - s)^2}{4\sigma^2} + \frac{im}{2t_F}(q_F - q_0)^2 \quad (3.61)$$

$$= -\left(\frac{1}{4\sigma^2} - \frac{im}{2t_F}\right)q_0^2 - \left(\frac{imq_F}{t_F} - ip - \frac{s}{2\sigma^2}\right)q_0 - \left(\frac{s^2}{4\sigma^2} - \frac{imq_F^2}{2t_F}\right), \quad (3.62)$$

with

$$a = \frac{1}{4\sigma^2} - \frac{im}{2t_F}, \quad b = \frac{imq_F}{t_F} - ip - \frac{s}{2\sigma^2} \quad \text{and} \quad \frac{s^2}{4\sigma^2} - \frac{imq_F^2}{2t_F}. \quad (3.63)$$

From section 3.2,

$$\exp\left[\frac{b^2}{4a} - c\right] = \exp\left[\frac{4impq_F\sigma^2 - mq_F^2 + 2mq_Fs - ms^2 - 2ip^2t_F\sigma^2 - 2pst_F}{4m\sigma^2 + 2it_F}\right] \equiv \Xi, \quad (3.64)$$

and

$$\sqrt{\frac{\pi}{a}} = 2\sigma\sqrt{\frac{\pi t_F}{t_F - 2im\sigma^2}}, \quad (3.65)$$

$$-\frac{b}{2}\sqrt{\frac{\pi}{a^3}} = 2\sigma\sqrt{\frac{\pi t_F}{(t_F - 2im\sigma^2)^3}}(-2iq_F + 2ipt_F\sigma^2 + st_F), \quad (3.66)$$

$$\frac{1}{4}\sqrt{\frac{\pi}{a^5}} = 2\sigma\sqrt{\frac{\pi t_F}{(t_F - 2im\sigma^2)^5}}(-4imt_F\sigma^4 + 2t_F^2\sigma^2 + (-2imq_F\sigma^2 + 2ipt_F\sigma^2 + st_F)^2). \quad (3.67)$$

Now let's consider the integral

$$I_1(p, s, \sigma, m; q_0, n) \quad (3.68)$$

$$= \int q_0^n K_0(q_F, q_0; t_F) \exp\left[ipq_0 - \frac{(q_0 - s)^2}{4\sigma^2}\right] dq_0 \quad (3.69)$$

$$= \sqrt{\frac{m}{2\pi it_F}} \int q_0^n \exp\left[ipq_0 - \frac{(q_0 - s)^2}{4\sigma^2} + \frac{im}{2t_F}(q_F - q_0)^2\right] dq_0 \quad (3.70)$$

$$= \sqrt{\frac{m}{2\pi it_F}} \int q_0^n \exp\left[-\left(\frac{1}{4\sigma^2} - \frac{im}{2t_F}\right)q_0^2 - \left(\frac{imq_F}{t_F} - ip - \frac{s}{2\sigma^2}\right)q_0 - \left(\frac{s^2}{4\sigma^2} - \frac{imq_F^2}{2t_F}\right)\right] dq_0$$

Let's abbreviate $I(q_F, p, s, \sigma, m; q_0, n)$ as $I(n)$ for now. For $n = 0$, $I(0)$ becomes

$$\int K_0(q_F, q_0; t_F) \exp\left[ipq_0 - \frac{(q_0 - s)^2}{4\sigma^2}\right] dq_0 \quad (3.71)$$

$$= \sqrt{\frac{m}{2\pi it_F}} \int \exp\left[-\left(\frac{1}{4\sigma^2} - \frac{im}{2t_F}\right)q_0^2 - \left(\frac{imq_F}{t_F} - ip - \frac{s}{2\sigma^2}\right)q_0 - \left(\frac{s^2}{4\sigma^2} - \frac{imq_F^2}{2t_F}\right)\right] dq_0 \quad (3.72)$$

$$= \sqrt{\frac{m}{2\pi i t_F}} \times 2\sigma \sqrt{\frac{\pi t_F}{t_F - 2im\sigma^2}} \times \Xi \quad (3.73)$$

$$= \sigma \Xi \sqrt{\frac{2m}{it_F(t_F - 2im\sigma^2)}}. \quad (3.74)$$

For $n = 1$,

$$I(1) = \int q_0 K_0(q_F, q_0; t_F) \exp\left[ipq_0 - \frac{(q_0 - s)^2}{4\sigma^2}\right] dq_0 \quad (3.75)$$

$$= \sqrt{\frac{m}{2\pi i t_F}} \times 2\sigma \sqrt{\frac{\pi t_F}{(t_F - 2im\sigma^2)^3}} (-2iq_F + 2ipt_F\sigma^2 + st_F) \times \Xi \quad (3.76)$$

$$= \sigma \Xi \sqrt{\frac{2m}{it_F(t_F - 2im\sigma^2)^3}} \times (-2iq_F + 2ipt_F\sigma^2 + st_F) \quad (3.77)$$

For $n = 2$,

$$I(2) = \int q_0^2 K_0(q_F, q_0; t_F) \exp\left[ipq_0 - \frac{(q_0 - s)^2}{4\sigma^2}\right] dq_0 \quad (3.78)$$

$$= \sqrt{\frac{m}{2\pi i t_F}} \times 2\sigma \sqrt{\frac{\pi t_F}{(t_F - 2im\sigma^2)^5}} \left(-4imt_F\sigma^4 + 2t_F^2\sigma^2 + (-2imq_F\sigma^2 + 2ipt_F\sigma^2 + st_F)^2\right) \times \Xi$$

$$= \sigma \Xi \sqrt{\frac{2m}{it_F(t_F - 2im\sigma^2)^5}} \left(-4imt_F\sigma^4 + 2t_F^2\sigma^2 + (-2imq_F\sigma^2 + 2ipt_F\sigma^2 + st_F)^2\right) \quad (3.79)$$

3.3.3 Joint probability contribution from the first order perturbation term on an entangled state

3.3.4 Second order perturbation term

The second order perturbation term, K_2 is

$$K_2 = \int \dots \int K_0(F, 2)V(2)K_0(2, 1)V(1)K_0(1, 0) dq_1 dq'_1 dt_1 dq_2 dq'_2 dt_2 \quad (3.80)$$

$$= \int \dots \int K_0(q_F, q_2; t_F - t_2)K_0(q'_F, q'_2; t_F - t_2)K_0(q_2, q_1; t_2 - t_1)K_0(q_2, q_1; t_2 - t_1) \quad (3.81)$$

$$\begin{aligned} & \times K_0(q_1, q_0; t_1 - t_0)K_0(q_1, q_0; t_1 - t_0)(q_2 - q'_2)^2(q_1 - q'_1)^2 dq_1 dq'_1 dt_1 dq_2 dq'_2 dt_2 \\ & = \int_{t_1}^{t_F} \int_{t_0}^{t_F} \left[\int \dots \int K_0(q_F, q_2; t_F - t_2)K_0(q'_F, q'_2; t_F - t_2)K_0(q_2, q_1; t_2 - t_1) \right. \\ & \quad \times K_0(q_2, q_1; t_2 - t_1)K_0(q_1, q_0; t_1 - t_0)K_0(q_1, q_0; t_1 - t_0) \left(q_1^2 q_2^2 - 2q_1^2 q_2 q'_2 \right. \\ & \quad \left. + q_1^2 (q'_2)^2 - 2q_1 q_2^2 q'_1 + 4q_1 q_2 q'_1 q'_2 - 2q_1 q'_1 (q'_2)^2 + q_2^2 (q'_1)^2 \right. \\ & \quad \left. \left. - 2q_2 (q'_1)^2 q'_2 + (q'_1)^2 (q'_2)^2 \right) dq_1 dq'_1 dq_2 dq'_2 \right] dt_1 dt_2 \end{aligned} \quad (3.82)$$

The integral once again can be broken into nine integrals that must be integrated w.r.t. time twice later on. All those nine integrals have a product of propagator as a multiplier. We shall evaluate those first, separating the primed and unprimed variables.

$$\begin{aligned} & K_0(q_F, q_2; t_F - t_2)K_0(q'_F, q'_2; t_F - t_2)K_0(q_2, q_1; t_2 - t_1)K_0(q_2, q_1; t_2 - t_1)K_0(q_1, q_0; t_1 - t_0) \\ & \quad \times K_0(q_1, q_0; t_1 - t_0) \end{aligned}$$

$$= \frac{im^3}{8\pi^3(t_F - t_2)(t_2 - t_1)(t_1 - t_0)} \exp \left[\frac{im}{2(t_F - t_2)} ((q_F - q_2)^2 + (q'_F - q'_2)^2) \right. \\ \left. + \frac{im}{2(t_2 - t_1)} ((q_2 - q_1)^2 + (q'_2 - q'_1)^2) + \frac{im}{2(t_1 - t_0)} ((q_1 - q_0)^2 + (q'_1 - q'_0)^2) \right] \quad (3.83)$$

3.4 The delta function collision problem

The potential for the delta function collision problem is

$$V = V_0 \delta(q - q') \quad (3.84)$$

where V_0 is the strength of the delta function, and is generally considered to be negative.

3.4.1 First order perturbation term

We evaluate the perturbation term similarly to how we did it in section 3.3.1, starting with the form:

$$K_1(F, 0; F', 0) \quad (3.85) \\ = -i \int_0^{t_F} \left[\iint K_0(q_F, q_1; t_F - t) K_0(q'_F, q'_1; t_F - t) K_0(q_1, q_0; t) K_0(q'_1, q'_0; t) V_0 \delta(q_1 - q'_1) dq_1 dq'_1 \right] dt \\ = -iV_0 \int_0^{t_F} \left[\iint K_0(q_F, q_1; t_F - t) K_0(q'_F, q'_1; t_F - t) K_0(q_1, q_0; t) K_0(q'_1, q'_0; t) \delta(q_1 - q'_1) dq_1 dq'_1 \right] dt$$

Let I_1 represents the integral in the square bracket; thus, $K_1 = -iV_0 \int_0^{t_F} I_1 dt$. Then,

$$I_1 = \iint K_0(q_F, q_1; t_F - t) K_0(q'_F, q'_1; t_F - t) K_0(q_1, q_0; t) K_0(q'_1, q'_0; t) \delta(q_1 - q'_1) dq_1 dq'_1 \quad (3.86)$$

$$= \int K_0(q_F, q'_1; t_F - t) K_0(q'_F, q'_1; t_F - t) K_0(q'_1, q_0; t) K_0(q'_1, q'_0; t) dq'_1 \quad (3.87)$$

3.5 The coulomb problem

CHAPTER 4

Correlation function

Implementation of simulations

To represent the joint probability distribution of a two particle system, we need four axes. Two for the position of each Hilbert space that those particles belong in, one for the probability amplitude, and one for the time.

$$P(q_F, q'_F; t_F) = |\langle q_F, q'_F | \eta(t_F) \rangle|^2 \quad (5.1)$$

$$= \left| \iint K(q_F, q'_F, t_F; q_0, q'_0, t_0) \eta(q_0, q'_0, t_0) dq_0 dq'_0 \right|^2 \quad (5.2)$$

$$= \left| \iint \left\{ \sum_{i=0}^{\infty} K_i(q_F, q'_F, t_F; q_0, q'_0, t_0) \right\} \eta(q_0, q'_0, t_0) dq_0 dq'_0 \right|^2 \quad (5.3)$$

For this equation to work with a discrete simulation, we need to exchange the integral sign with the summation sign:

$$= \left| \sum_{q_0} \sum_{q'_0} \left\{ \sum_{i=0}^{\infty} K_i(q_F, q'_F, t_F; q_0, q'_0, t_0) \right\} \eta(q_0, q'_0, t_0) \right| \quad (5.4)$$

The sum of propagators can then be truncated to our liking. I've written the code in Julia to test for the free particle propagator, and it looks something like this:

```

1 using Plots; plotlyjs()
2 using LinearAlgebra
3 using SymPy
4 using BenchmarkTools
5
6 q, q0, q1, q2, qf = symbols("q q_0 q_1 q_2 q_F", real = true)
7 m, t, t0, t1, t2, tf = symbols("m t t_0 t_1 t_2 t_F", real = true, positive = true)
8 q0p, q1p, q2p, qfp = symbols("q^{\\prime}_0 q^{\\prime}_1 q^{\\prime}_2 q^{\\prime}_F",
  ↪ real = true)
```

```

9  a, b, c, d, e, f = symbols("a b c d e f")
10 s1, s2, p1, p2, σ1, σ2 = symbols("s_1 s_2 p_1 p_2 σ_1 σ_2")
11
12 freePropagator(finPos, startPos, finTime, startTime = 0, m = 1) = sqrt(m / (2 * pi * im *
    ↪ (finTime - startTime))) * exp(im * m / (2 * (finTime - startTime)) * (finPos -
    ↪ startPos)^2)
13 freePropagatorC(qf, qfp, q0, q0p, tf, t0) = freePropagator(qf, q0, tf, t0) +
    ↪ freePropagator(qfp, q0p, tf, t0)
14
15 initStateFunction(q0, q0p, σ1, s1, p1, σ2, s2, p2) = (1//2 * pi * σ1)^(1//4) * exp(-(q0 -
    ↪ s1)^2 / (4 * σ1^2) + im * p1 * q0) * (1//2 * pi * σ2)^(1//4) * exp(-(q0p - s2)^2 / (4 *
    ↪ σ2^2) + im * p2 * q0p)
16
17 maxPos = 5
18 minPos = -5
19 stepPos = 0.25
20
21 pos1Vect = collect(minPos:stepPos:maxPos)
22 pos2Vect = collect(minPos:stepPos:maxPos)
23 posVectSize = size(pos1Vect, 1)
24 posMat = [(i, j) for i in pos1Vect, j in pos2Vect]
25
26 posToIndex(pos) = Int32((pos - minPos) / stepPos + 1)
27
28 initState(q) = initStateFunction(q[1], q[2], 1, 2, 2, 1, -2, -2)
29 initMat = initState.(posMat)
30 initMat = round.(initMat, digits = 7)
31
32 surface(pos1Vect, pos2Vect, abs.(initMat))
33
34 finalTime = 1
35 finalMat = Matrix{ComplexF32}(undef, posVectSize, posVectSize)
36
37 for xf in pos1Vect, xfp in pos2Vect
38     sumPos = 0
39     for i in 1:posVectSize, j in 1:posVectSize

```

```
40     x0 = pos1Vect[i]
41     x0p = pos2Vect[j]
42     sumPos += freePropagatorC(xf, xfp, x0, x0p, finalTime, 0) * initMat[i, j]
43 end
44 finalMat[posToIndex(xf), posToIndex(xfp)] = sumPos
45 end
46
47 surface(pos1Vect, pos2Vect, abs2.(finalMat))
```


CHAPTER 6

The Schrödinger's equation for two particles

Integrals evaluation code

All of these codes that I've written are in the Julia language, which I've imported three packages: `SymPy`, `OffsetArrays`, and `Plots`; `plotlyjs()`

A.1 Polynomial extraction

Since we're going to be doing a lot of polynomials rearranging, I've implemented the polynomial extraction function as follows:

```
1 function extractPolynomial(expr, arg)
2     expr isa Sym ? nothing : expr = sympify(1)
3     expr = expand(expr)
4     polyDegree = Int(degree(expr, arg))
5     extractOneSet = []
6     for i in 0:polyDegree
7         temp = expr.coeff(arg, i)
8         push!(extractOneSet, temp)
9     end
10    polyExt = OffsetVector(extractOneSet, 0:polyDegree)
11    return polyExt
12 end
```

This function accepts two arguments: `expr`, which is the expression you want to extract, and `arg`, the variable that you extract with respect to. For example, inputting $ax^2 + bx + c$, `x` would give

out

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix}, \quad (\text{A.1})$$

which is a zero index matrix.

A.2 Code for the spring problem

A.2.1 First order perturbation

The first order perturbation uses the following symbols:

```

1 @syms q0 q1 q2
2 @syms m::(real, positive) t::(real, positive)
3 q0', q1', q2', qf', qf = symbols("q_{0}^{\prime}, q_{1}^{\prime}, q_{2}^{\prime},
   ↪ q_{F}^{\prime}, q_F")
4 t0, t1, t2, tf = symbols("t_0, t_1, t_2, t_F", real = true, positive = true)

```

The following code is used to aid the evaluation of integrals.

```

1 # Simplifying the product of propagators in eq. 2.25
2 idenFunc = IM*m/(2 * (tf - t)) * (qf - q1)^2 + IM*m/(2*t) * (q1 - q0)^2
3
4 idenPoly = extractPolynomial(idenFunc, q1)
5 idenPolyA = -idenPoly[2]
6 idenPolyB = -idenPoly[1]
7 idenPolyC = -idenPoly[0]
8
9 display(idenPolyA)
10 display(idenPolyB)
11 display(idenPolyC)
12

```