

# CS220 Programming Principles

## Homework No. 7

**Due: December 19, 2018 10:30AM ( 4x9 points)**

Submit your racket file to TA.

Use the given hw07-code.scm file as the starting point of your homework.

Before you submit it, rename it to "hw07-your\_student\_number.scm."

### Part 1 : Exercises

**Exercise 1:** Do exercise 3.51 of the SICP book, which takes a closer look at delayed evaluation and stream-enumerate-interval.

**Exercise 2:** Describe the streams produced by the following definitions. Assume that integers is the stream of non-negative integers (starting from 1):

```
(define A (stream-cons 1 (scale-stream 2 A)))

(define (mul-streams a b)
  (stream-cons
    (* (stream-first a) (stream-first b))
    (mul-streams (stream-rest a)
                  (stream-rest b))))

(define B (stream-cons 1 (mul-streams B integers)))
```

**Exercise 3:** Given a stream *s* the following procedure returns the stream of all pairs of elements from *s*:

```
(define (stream-pairs s)
  (if (stream-empty? s)
      empty-stream
      (stream-append
        (stream-map
          (lambda (sn) (list (stream-first s) sn))
          (stream-rest s))
        (stream-pairs (stream-rest s)))))
```

- Suppose that integers is the (finite) stream {1, 2, 3, 4, 5}. What is (stream-pairs s)?
- Give the clearest explanation that you can of how stream-pairs works.
- Suppose that *s* is the stream of positive integers. What are the first few elements of (stream-pairs s)? Can you suggest a modification of stream-pairs that would be more appropriate in dealing with infinite streams?

### Part 2: Working with Power Series

In section 2.5.3, we saw how polynomials could be represented as lists of terms. In a similar way, we can represent power series, such as

$$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{3 \cdot 2} + \frac{x^4}{4 \cdot 3 \cdot 2} + \dots$$

$$\cos x = 1 - \frac{x^2}{2} + \frac{x^4}{4 \cdot 3 \cdot 2} - \dots$$

$$\sin x = x - \frac{x^3}{3 \cdot 2} + \frac{x^5}{5 \cdot 4 \cdot 3 \cdot 2} - \dots$$

as streams of infinitely many terms. That is, the power series

$$a_0 + a_1x + a_2x^2 + a_3x^3 + \dots$$

will be represented as the infinite stream whose elements are  $\{a_0, a_1, a_2, a_3, \dots\}$ <sup>1</sup>

## Implementing a Power Series as a Stream

We provide two ways to construct a series: `coeffs->series` and `proc->series`. The procedure `coeffs->series` takes a list of initial coefficients and pads it with zeroes to produce a powers series. For example, `(coeff->series '(1 3 4))` produces the power series  $1 + 3x + 4x^2 + 0x^3 + 0x^4 + \dots$

```
(define (coeffs->series list-of-coeffs)
  (define zeros (stream-cons 0 zeros))
  (define (iter lst)
    (if (null? lst)
        zeros
        (stream-cons (car lst)
                      (iter (cdr lst)))))
  (iter list-of-coeffs))
```

The other constructor, `proc->series`, takes as argument a procedure `p` of one numeric argument and returns the series

$$p(0) + p(1)x + p(2)x^2 + p(3)x^3 + \dots$$

In the definition of `proc->series`, the stream `non-neg-integers`, which you will define in this problem set, is the stream of non-negative integers:  $\{0, 1, 2, 3, \dots\}$ .

```
(define (proc->series proc)
  (stream-map proc non-neg-integers))
```

## Other Stream Operations

With this representation, we can use basic stream operations such as:

```
(define (add-streams s1 s2)
  (cond ((stream-empty? s1) s2)
        ((stream-empty? s2) s1)
        (else
         (stream-cons (+ (stream-first s1) (stream-first s2))
                       (add-streams (stream-rest s1)
                                     (stream-rest s2)))))
```

---

<sup>1</sup> In this representation, all streams are infinite: a finite polynomial will be represented as a stream with an infinite number of trailing zeroes.

```

                                (stream-rest s2))))))
(define (scale-stream c stream)
  (stream-map (lambda (x) (* x c)) stream))

```

to implement series operations, and thus arrive at a complete power series data abstraction:

```

(define add-series add-streams)

(define scale-series scale-stream)

(define (negate-series s)
  (scale-series -1 s))

(define (subtract-series s1 s2)
  (add-series s1 (negate-series s2)))

(define (show-series s nterms)
  (if (= nterms 0)
      'done
      (begin (write-line (stream-first s))
              (show-series (stream-rest s) (- nterms 1))))))

(define (series-coeff s n)
  (stream-ref s n))

```

You will find the `show-series` procedure helpful because it allows you to examine the series that you will generate in this problem set. You can also examine an individual coefficient (of  $x^n$ ) in a series using `series-coeff`.

**Note:** After loading the code for this problem set, you will find that Scheme's basic arithmetic operations `+`, `-`, `*`, and `/` will work with rational numbers. For instance, `(/ 3 4)` will produce `3/4` rather than `.75`. You'll find this useful in doing the exercises below.

**Problem 1:** Load the code for homework 7 (file `hw07-code.scm`). Define the stream `non-neg-integers`. Show how to define the series:

$$S_1 = 1 + x + x^2 + x^3 + \dots$$

$$S_2 = 1 + 2x + 3x^2 + 4x^3 + \dots$$

Turn in your definitions and a couple of coefficient printouts to demonstrate that they work.

**Problem 2:** Complete the definition of the following procedure, which multiplies two series:

```

(define (mul-series s1 s2)
  (stream-cons < ??E1?? >
    (add-series < ??E2?? >
      < ??E3?? > )))

```

To test your procedure, demonstrate that the product of  $S_1$  (from problem 1) and  $S_1$  is  $S_2$ . What is the coefficient of  $x^{10}$  in the product of  $S_2$  and  $S_2$ ? Turn in your definition of `mul-series`. (Optional: Give a general formula for the coefficient of  $x^n$  in the product of  $S_2$  and  $S_2$ .)

## Inverting a power series

Let  $S$  be a power series whose constant term is 1. We'll call such a power series a "unit power series." Suppose we want to find the *inverse* of  $S$ ,  $1/S$ , which is the power series  $X$  such that  $S \bullet X = 1$ . Write  $S = 1 + S_R$  where  $S_R$  is the rest of  $S$  after the constant term. Then we can solve for  $X$  as follows:

$$S \bullet X = 1$$

$$(1 + S_R) \bullet X = 1$$

$$\begin{aligned}X + S_R \bullet X &= 1 \\X &= 1 - S_R \bullet X\end{aligned}$$

In other words,  $X$  is the power series whose constant term is 1 and whose rest is given by the negative of  $S_R$  times  $X$ .

**Problem 3:** Use this idea to write a procedure `invert-unit-series` that computes  $1/S$  for a unit power series  $S$ . You will need to use `mul-series`. To test your procedure, invert the series  $S_1$  (from problem 1) and show that you get the series  $1 - x$ . (Convince yourself that this is the correct answer.) Turn in a listing of your procedure. This is a very short procedure, but it is very clever. In fact, to someone looking at it for the first time, it may seem that it can't work—that it must go into an infinite loop. Write a few sentences of explanation explaining why the procedure does in fact work, and does not go into a loop.

**Problem 4:** Write a procedure `div-series` that divides two power series. `div-series` should work for any two series, provided that the denominator series begins with a non-zero constant term. (If the denominator has a zero constant term, then `div-series` should signal an error.) You may find it useful to reuse some answer from a previous exercise in this problem set. Turn in a listing of your procedure along with three or four well-chosen test cases (and demonstrate why the answers given by your division are indeed the correct answers).

**Problem 5:** Define a procedure `integrate-series-tail` that, given a series

$$a_0 + a_1x + a_2x^2 + a_3x^3 + \cdots$$

returns the integral of the series (except for the constant term)

$$a_0x + \frac{1}{2}a_1x^2 + \frac{1}{3}a_2x^3 + \frac{1}{4}a_3x^4 + \cdots$$

Turn in a listing of your procedure and demonstrate that it works by computing `integrate-series-tail` of the series  $S_2$  from problem 1.

**Problem 6:** Demonstrate that you can generate the series for  $e^x$  as

```
(define exp-series
  (stream-cons 1 (integrate-series-tail exp-series)))
```

Explain the reasoning behind this definition. Show how to generate the series for sine and cosine, in a similar way, as a pair of mutually recursive definitions.