



* Injections / Implants (under Skin) * Oral Pills oprogesterone progesterone-estrogen combination in tablet form. · progestinone alone or with estragen. · Taken daily for 21 days, starting within first 5 days of cycle. · Mode of action similar to pills. Effective period is long. · Break of 7 days, and then again. * Emergency Contraceptives. • Inhibits Ovulation & Implantation. TUD's within 72 howes very effective. · Alter cervical mucus to retard sperms. · Very effective, less Side effects. Saheli → CDRI Lukhnow (non-steroidal)
 Once a week pill. · Terminal Method. * Surgical Methods. (Sterlisation) Tubectomy Blocks gamete transport. · Highly effective. Vasectomy (Males) (females) · Poor Reversibility. Small part of Vas-Small part of pallopian tube removed or tied up by small incision descence is removed or tied up by small incision on soutum. *Some ill effects >> -Nausea, abdominal pain, breakthrough, bleeding, irregular bleeding, breast cancer. in abdomen through Vagina > Not very significant. * Medical Termination of Pregnancy (MTP). * Sexually Transmitted Infections G Intentional /Voluntary termination. (Induced)

G 45-50 million MTP's in Year (inworld)

1/5th of total conceived pregnancies. · Diseases through Sexual Intercourse · Veneral Diseases CAMTP was allowed after considering emotional, ethical, religious and Social issues. · Reproductive Tract Infections. (1) Hepatitu-B >#1971 (India) → MTP legalised with struck () Gonoruhoea condition. (as fimale forticide 11) 8 AIDS @ Syphilis Com be terminated within 12 weeks - registered practitioner. 3) Genital hexpes Due to shaving CoMore than 12 weeks-24 weeks - 2 registered practitioner 4) Genital Warts

O Woman Life Risk.

Trijury to physical Mental Health. injection needles, Surgical instrument Traspusion of blood 6) Truchomoniasis from mother - child. (2) Child born - Physical Mental abnormalities.

Seriously handicapped. Hepatitis-B ? Not curable. (3) Get sid of Unwanted Prugnancies. Genital Herps } All others Carable, if detected Intercourse HIV detected early. · Early Symptoms → Itching, fluid Discharge, Slight pain, swelling in genital tract. -> Failure of Contraceptive. · Sometimes Undetected / Social Stigma for STD's can lead to pelvic inflammatory diseases aboutions, Still brains, ectopic pregnancies, infertility, Cancer. · 15-24 yrs (Most Vulnerable) → Avoid Sex with multiple partners. → Use Condoms. → Go to qualified doctors.

