

[Skip to main content](#)

SPRINGER LINK

[Log in](#)

 [Menu](#)

[Find a journal](#) [Publish with us](#)

 [Search](#)

 [Cart](#)

1. [Home](#) >
2. [Waste and Biomass Valorization](#) >
3. Article

Valorization of Fish Waste Compost as a Fertilizer for Agricultural Use

- Original Paper
- [Open access](#)
- [Published: 16 April 2018](#)
- **volume 10**, pages 2537–2545 (2019)

[Download PDF](#) 

✓ You have full access to this [open access](#) article



[Waste and Biomass Valorization Aims and scope](#) → [Submit manuscript](#) →



Valorization of Fish Waste Compost as a Fertilizer for Agricultural Use

[Download PDF](#) 

- [Maja Radziemska](#) [ORCID: orcid.org/0000-0002-3526-5944](https://orcid.org/0000-0002-3526-5944)¹,

- [Magdalena Daria Vaverková](#) [ORCID: orcid.org/0000-0002-2384-6207](https://orcid.org/0000-0002-2384-6207)²,
- [Dana Adamcová](#)²,
- [Martin Brtnický](#)³ &
- ...
- [Zbigniew Mazur](#)⁴✉

▼ Show authors


-  20k Accesses
-  54 Citations
- [Explore all metrics](#) →


[Cite this article](#)

Abstract


The manuscript presents results of the evaluation of compost from fish waste (FW) as a fertilizer for agricultural use. A pot experiment was conducted to compare the effects of compost from FW on the yield and macro and microelemental composition of ice lettuce (*Lactuca sativa* L.). In addition, the phytotoxicity degree of the compost and compost effects on seed germination and primary root growth were determined with white mustard (*Sinapis alba* L.). Compost used in the study consisted of FW and pine bark. Results of the evaluation enable concluding that the compost from FW is non-phytotoxic, mature, stable, and suitable for use in agriculture. Its addition to soil caused an increase in fresh and dry matter yield of leaves of ice lettuce (*L. sativa* L.). Fertilization had a significant effect on increased contents of nitrogen, phosphorus, potassium, sodium, calcium, and magnesium in leaves of the test plant. The average accumulation of microelements in ice lettuce (*L. sativa* L.) grown in the soil fertilized with compost from FW followed the descending order Fe > Cu > Ni > Zn > Mn, respectively. Soil fertilization with compost from FW improved the K:(Mg + Ca), K:Mg and K:Ca ratios but, simultaneously, deteriorated the Ca:P ratio.

Graphical Abstract






Similar content being viewed by others



Impact of Biochar from Rice Husk on Nutrient Distribution and Rice Growth and Yield: A Soil Column Experiment

Article 27 November 2023

Chenda Lai, Ratha Muon, ... Vannak Ann



Vermicomposting of different organic materials using the epigeic earthworm *Eisenia foetida*

Article Open access 25 October 2018


Yvonne Indrani Ramnarain, Abdullah Adil Ansari & Lydia Ori



Preparation of nano-fertilizer blend from banana peels

Article Open access 12 February 2019

H S Hussein, H H Shaarawy, ... S I Hawash



[Use our pre-submission checklist](#)

Avoid common mistakes on your manuscript.



Introduction

Directive 2008/98/EC was adopted to introduce a new approach to the handling of waste and its management. Worldwide solid waste (SW) is an important and emerging environmental problem [1]. Accordingly, the Waste Framework Directive (WFD) and hierarchy priorities the prevention of waste generation, followed by processing for reuse and recycling, with disposal as the least favored stage of waste management (WM). Food waste was considered a special case in the WFD, focusing on three key points: the separate collection of bio-waste, treatment of bio-waste to ensure maximum environmental protection (composting and digestate), and the development of techniques to produce environmentally safe materials from bio-waste [2]. Food waste comprises the main fraction (45%) of total municipal solid waste (MSW) in Europe [3]. Sustainable management of food waste is a momentous research area that has rapidly grown over recent years [4]. Worldwide there are several techniques for waste processing into value-added products in the form of compost (organic fertilizer), biogas, animal feed and chemicals [5]. Composting is a natural process [6], an eco-friendly biochemical technique, and a viable alternative for a sustainable MSW management [7].

The Regulation (EC) of the European Parliament and the Council on health rules concerning animal by-products not intended for human consumption describes products derived from aquatic animals, including fish, as removed e.g. by incineration or co-incineration, fermentation, composting or transformation into biogas [8]. Fish waste (FW) can be suitable material for composting. One such material is, which is yielded as a by-product of fish production and fish processing industries. Depending on the type of transformation, the waste may represent between 30 and 45% of the initial weight of the product [9].

FW is suitable for agricultural use owing to high contents of nutrients, such as N, P, and Ca [9]. Several fertilizers made of fishmeal are now commercially available and some are authorized for use in organic agriculture [10]. Even fish effluent can be used to irrigate cherry tomato plants [9, 11]. Composting initiatives using FW have been carried out in various parts of the world in search of

alternative and viable techniques for transforming fish waste into useful agricultural products [12].

Considering the benefits of composting, the composting itself is a biotransformation process of organic materials into stable and complex macromolecules under the action of microorganisms including fungi, bacteria, and corresponding enzyme(s) [13,14,15]. It includes four phases: (i) an initial decomposition phase; (ii) a thermophilic phase of intense microbial decomposition; (iii) the second thermophilic phase; and (iv), a maturation phase. Rapidly multiplying thermophilic bacterial species dominate during the thermophilic phase till the moment when the bulk of the easily decomposable substrate is exhausted. The majority of the remaining material is woody with lignin as the dominating part, which is also stabilized by humic acids and fungi [15, 16]. The final product, compost, can be used as a soil amendment that improves soil texture and fertility and thus reduces the use of synthetic fertilizers applied to the soil. In this case, the conversion should be performed by implementation of novel technologies for the recycling of waste in the form of compost for their use in agriculture [17]. In addition to the usage of compost as a fertilizer, applying compost to the soil may increase the carbon storage capacity within the soil, which reduces greenhouse gas (GHG) emissions into the atmosphere [3, 18, 19].

The above-mentioned material is of great potential use in agriculture. However, its stabilization is recommended prior to its use to prevent problems associated with the appearance of phytotoxic substances [9, 20]. When compost is land applied, finished compost with no phytotoxin can be regarded as a source of nutrients to improve soil structural properties and accelerate germination. Therefore, assessing the phytotoxicity degree of the compost products is essential to achieve high quality compost [21].

The aim of the present paper was to assess the fertilizing power of compost consisting of FW and pine bark. The phytotoxicity degree of the compost and effects of the compost on seed germination and primary root growth were determined with white mustard (*Sinapis alba* L.). Additionally, the effect of the compost application on ice lettuce (*Lactuca sativa* L.) was determined. The authors hypothesized that FW compost might be a suitable for agricultural use.

Materials and Methods

Compost used in the experiment was prepared from (w/w) 80% FW and 20% pine bark (10–15 mm chip size) as a bulking agent and carbon source with the following composition: organic carbon – 566.2 ± 7.3 (g/kg d.m.), total N – 1.92 ± 0.1 (g/kg d.m.), phosphorous – 0.40 ± 0.1 (g/kg d.m.), potassium – 0.94 ± 0.1 (g/kg d.m.), magnesium – 1.12 ± 0.1 (g/kg d.m.), and calcium – 1.22 ± 0.1 (g/kg d.m.). Contents of macro- and micro-elements in compost dry matter were presented in Table 1.

Table 1 Chemical composition of fish waste compost, in dry matter

[Full size table](#) >

Phytotoxicity Test

Maturity and toxicity of the compost were determined to establish its usability for agricultural

purposes. The seeds of white mustard (*S. alba* L.) were used in a seed germination experiment to assess the phytotoxicity [22] of the final compost [23]. The Phytotoxkit makes use of flat and shallow transparent test plates composed of two compartments, the lower one which contains soil saturated to the water holding capacity. The phytotoxkit measures the decrease (or the absence) of seed germination and of the growth of young roots after 3 days of the exposure of selected seeds of higher plants to a contaminated matrix, in comparison to the controls in a reference soil. Water saturation is calculated according to the user's manual. Distilled water was spread over the entire surface of the soil in the test plate. Ten seeds of *S. alba* L. were positioned at equal distances near the middle ridge of the test plate on a filter paper placed on the top of the hydrated soil/compost mixture. After closing, the test plates were placed vertically in a holder and incubated at 25 °C for 3 days. At the end of the incubation period a digital picture was taken of the test plates with the germinated plants. The following concentrations of compost were tested in three replications: 25, 50, and 100%. The analyses and the length measurements were performed using the Image Tool 3.0 for Windows (UTHSCSA, San Antonio, USA). The bioassays were performed in three replicates [24].

Pot Experiment

The fertilizing efficiency of the compost was assessed by its application to ice lettuce (*L. sativa* L.) crop. Tests using plants such as *L. sativa* L., present several advantages. Firstly, these tests are simple, quick, and reliable. Secondly, they are inexpensive and do not require major equipment. Thirdly, plants can be more sensitive to environmental stress over other organisms [25]. The vegetation pot experiment was conducted at a steady temperature of 22 ± 2 °C in the greenhouse facility of the University of Warmia and Mazury in Olsztyn (Poland). The pots were maintained under natural day/night conditions; during the day (14 h), with a relative humidity of $75 \pm 5\%$. Soil used in the experiment was obtained from the arable layer (0–20 cm) of a farm field. PCV pots filled with 10 kg of soil that had been previously mixed with the compost in accordance with the study procedure were used for this purpose. The soil used in the experiment was characterized by a pH value of 5.7 and granulometric composition of 86% sand (2.0–0.05 mm), 11.2% dust (0.05–0.002 mm), and 2.2% content of fractions under 0.002 mm in diameter (S according to USDA). The content of organic carbon and nitrogen in this soil was: Corg. – 6.33 g/kg, N-total – 0.55 g/kg, absorbable elements: phosphorus – 84.86 mg/kg, magnesium – 77.48 mg/kg, and potassium – 57.89 mg/kg soil. The soil samples were air-dried and passed through a 5-mm sieve prior to the greenhouse pot experiment.

The experiment was conducted in four replicates. Compost was applied in the amount of 2 g N compost per pot. Non-fertilized pots served as control objects. The effect of fertilizations was tested on ice lettuce (*L. sativa* L.) of the Dobra variety. The density was one plant per pot. During the vegetation, soil humidity was maintained at the level of 60% capillary water capacity. The lettuce was picked upon reaching full maturity (after 51 days of vegetation).

Analytical Methods

Compost samples were mineralized in concentrated sulfuric(VI) acid with hydrogen peroxide added as a catalyst. After mineralization of the compost samples, the content of phosphorus and potassium

was analyzed with the vanadium–molybdenum method, and that of calcium and sodium with atomic emission spectroscopy—AES on FLAPHO 4 CZJ model. Magnesium content was assayed with atomic absorption spectroscopy—AAS and that of total nitrogen using Kjeldahl's method.

The contents of micronutrients: copper, cadmium, chromium, lead, nickel, and zinc were determined in an air–acetylene flame, following the flame atomic absorption spectrophotometric method (FAAS) on a SpectrAA 240FS atomic absorption spectrophotometer (VARIAN, Australia) using a Sample Introduction Pump System (SIPS) in extracts obtained after microwave digestion of compost samples in nitric acid hydrogen (analytically pure HNO_3) at a concentration of 1.40 g/cm, poured into HP500 Teflon vessels and placed in an MARS-5 microwave oven (CEM Corporation, USA). The mineralization conditions, i.e. volume of nitrogen acid and mineralization temperature, were set as described in the methodology given in the Mars 5 Operation Manual (2001). Compost samples were digested and analyzed using the same reagents. Double deionized water (Milli-Q Millipore 0.055 $\mu\text{S}/\text{cm}$ resistivity) was used for all dilutions. Organic carbon content of compost was determined with the method of non-dispersive infrared (NDIR) spectrometry using a TOC–VCPN analyzer with an SSM-5000A module for carbon content assay in solid samples by Shimadzu.

Calculations and Data Analysis

The percent inhibition of root growth (RI) were calculated as follows [24, 26]:

$$\text{RI}(\%) = \frac{\{\text{Mean root length in the control}\} - \{\text{Mean root length in the test}\}}{\{\text{Mean root length in the control}\}} \times 100$$

$$\text{GI}(\%) = \left(\frac{\{\text{Seed germination}\}}{\{\text{Seed germination}\}} \right) \times \left(\frac{\{\text{Root elongation}\}}{\{\text{Root elongation}\}} \right) / 100$$


The results were subjected to statistical calculations using the Statistica 10 software package [27] (StatSoft Inc. 2010), with results of each series subjected to statistical analysis using Tukey's analysis of variance. The series were compared with the t test for dependent variables, assessing differences between the average results.

Results and Discussion

At the end of the 3-day toxicity bioassay—Phytotoxkit™ (Fig. 1), the length of roots was measured in particular test plants grown in soil with various concentrations of the compost from FW.

Fig. 1






Phytotoxicity test in laboratory conditions—the experiment setup

[Full size image](#) >

Figure 2 presents results of measurements of root length of plants grown in soil fertilized with a 25% dose of compost for all three replicates. This concentration of compost stimulated root growth of white mustard (*S. alba* L.) seeds and ensured the highest values of root elongation (of all compost concentrations analyzed). The mean root length was 52.27 mm.

Fig. 2




Length of roots at compost concentration of 25%

[Full size image](#) >

Further analyzed were the samples of plants grown in soil fertilized with 50% dose of the fish compost. Figure 3 depicts length of roots of test plants grown at this compost concentration for all three replicates. At this concentration of compost, roots of test plants were shorter compared to root lengths determined at compost concentration of 25%. The mean root length reached 42.42 mm and

was, on average, by 9.85 mm lesser than in the samples grown at compost concentration of 25%.

Fig. 3




Length of roots at compost concentration of 50%

[Full size image](#) >

The toxicity symptoms become severe with the increased dose of compost. Figure 4 presents the mean root length of plants grown at 100% compost concentration, which was significantly lower compared to plants fertilized with 25 and 50% of compost. The mean root length accounted for 37.56 mm and was lower by 14.71 mm and by 4.86 mm than in plants grown at 25 and 50% compost concentration.

Fig. 4




Length of roots at compost concentration of 100%

[Full size image](#) >

Afterwards, root growth inhibition was computed for particular compost concentrations and respective results were presented in Fig. 5. Compost concentration of 25% had a stimulating effect on root growth and caused root elongation by 5.23%, whereas the other two compost doses inhibited root growth by 13.69% (50% compost) and 31.12% (100% compost).

Fig. 5



Root growth inhibition at various compost concentrations

[Full size image](#) >

The germination index (GI) is a factor of relative seed germination and relative root elongation. It was proved to be one of the most sensitive parameters that may take account of both low toxicity affecting root growth as well as high toxicity affecting germination, and finally the degree of compost maturity [28]. In general, the decrease of phytotoxicity during composting results from the degradation of phytotoxic substances by microorganisms [29]. Aggelis et al. [30] proposed that: if the $GI < 25$ then the substrate is characterized as very phytotoxic, if $26 < GI < 65$ then the substrate is characterized as phytotoxic, if $66 < GI < 100$ then the substrate is characterized as non-phytotoxic; stable and can be used for agricultural purposes as a fertilizer. According to that suggestion, the compost analyzed in our study can be deemed as phytotoxin-free, (GI values obtained ranged between 69.4% for concentration 25 and 67.7% for concentration 50%), and safe for application as soil fertilizer.

There is no widespread consensus on the influence of organic fertilizers on crop production. Several studies have evaluated the fertilizing effects of composts and have suggested composting as one of the most appropriate techniques for producing organic fertilizers [9, 31, 32]. Composting FW has also been suggested as a valid method of transforming this waste into useful soil amendments for agricultural purposes [9]. Results of determination of the fresh and dry matter yield of leaves of ice lettuce (*L. sativa* L.) are presented in Table 2.

Table 2 Effect of fish compost application on the fresh and dry matter yield of leaves of ice lettuce (*Lactuca sativa* L.)

[Full size table](#) >

Compared to the control pot, fertilization with FW compost caused almost twofold increase in fresh

matter yield and nearly 50% increase in dry matter yield of ice lettuce (*L. sativa* L.) leaves. The beneficial effect of compost made of FW, bark, and straw on fresh and dry matter yield of vegetables was earlier confirmed by [8, 12, 33, 34].

Soil fertilization with FW compost increased contents of macroelements in lettuce leaves (Table 3) by 78.6% in the case of nitrogen, by 61.8% in the case of phosphorus, by 56.3% in the case of potassium, by 44.4% in the case of sodium, and by 38.5% in the case of calcium and magnesium.

Table 3 Macroelement content in ice lettuce (*Lactuca sativa* L.) leaves, g/kg d.m.

[Full size table](#) >

The obtained results can be compared to studies carried out by Krzebietke [35] in which the author showed the content of macroelements in the leaves of butter lettuce subjected to nitrogen fertilization to be as follows: 4.9–6.0 g/kg phosphorus, 40.2–45.2 g/kg potassium, 17.1–23.0 g/kg calcium, 0.8–7.3 g/kg sodium, and 2.8–4.0 g/kg d.m. magnesium. The quality of the edible plant parts depends not only on the concentration of macro and microelements but also on their ratios, which may be affected not only by the species of plant but also by the proportions of cations in the fertilizers [36] (Ekholm et al. 2007). The optimal proportions between the individual elements should be higher than Ca:P—2; Ca:Mg—3; K:(Ca + Mg)—1.6–2.2; K: Mg—6; and K: Ca—2 [37, 38]. The studies confirmed that the leaves of ice lettuce were characterized by a wider ranges of K:(Mg + Ca), K:Mg and K:Ca (Table 4). Soil fertilization with compost from fish waste improved the above ratios but, simultaneously, deteriorated the Ca:P ratio.

Table 4 Mass ratios between macroelements in ice lettuce (*Lactuca sativa* L.) leaves

[Full size table](#) >

Analyses showed an increase in contents of all analyzed microelements in ice lettuce (*L. sativa* L.) leaves (Table 5). The percentage increase in the concentration of the analyzed microelements ranged from 83.2% (Fe) to 35.7% (Mn) and followed the descending order: Fe > Cu > Ni > Zn > Mn.

Table 5 Microelement content in ice lettuce (*Lactuca sativa* L.) leaves, mg/kg d.m.

[Full size table](#) >

Radziemska and Mazur [8], who analyzed contents of heavy metals in aerial parts of corn fertilized with various composts made of FW, demonstrated the uptake of nickel, copper, and zinc by plant to depend on the chemical composition of composts and additional mineral fertilization. The concentration of macro- and micro-elements in leaves of lettuce (*L. sativa* L.) results from their contents in the compost in forms available to plants, and this availability depends on, among other things, pH value, air–water ratio, and prevalence of microorganisms [39, 40].

Conclusions

Composting of fish waste with pine bark allows for a significant reduction in the volume of fisheries by-products and waste. The stability and maturity of the compost are essential for its successful

application, particularly for composts used in agriculture. However, they are not easy to define and cannot be described in a single parameter. The conducted experiments indicate the compost product to be non-phytotoxic, mature, stable, and suitable for use in agriculture. Soil fertilization with the compost from fish waste caused an increase in leaf yield of ice lettuce (*L. sativa* L.) and had a significant effect on increased contents of nitrogen, phosphorus, potassium, sodium, calcium, and magnesium in leaves of the test plant. The average accumulation of microelements in ice lettuce (*L. sativa* L.) grown in soil fertilized with the compost from fish waste followed the descending order Fe > Cu > Ni > Zn > Mn, respectively. The proposed fish waste compost can be a useful fertilizer in agriculture.

References

1. Adamcová, D., Vaverková, M.D., Bartoň, S., Havlíček, Z., Broušková, E.: Soil contamination in landfills: a case study of a landfill in Czech Republic. *Solid Earth* **7**, 239–247 (2016).
<https://doi.org/10.5194/se-7-239-2016>
[Article](#) [Google Scholar](#)
2. Ravindran, R., Jaiswal, A.K.: Exploitation of food industry waste for high-value products. *Trends Biotechnol.* **34**, 58–69 (2016)
[Article](#) [Google Scholar](#)
3. Cerda, A., Artola, A., Font, X., Barrena, R., Gea, T., Sánchez, A.: Composting of food wastes: status and challenges. *Biores. Technol.* (2017). <https://doi.org/10.1016/j.biortech.2017.06.133>
[Google Scholar](#)
4. Garcia-Garcia, G., Woolley, E., Rahimifard, S., Colwill, J., White, R., Needham, L.: A methodology for sustainable management of food waste. *Waste Biomass Valor.* **8**, 1–19 (2017).
<https://doi.org/10.1007/s12649-016-9720-0>
[Article](#) [Google Scholar](#)
5. Awasthi, M.K., Pandey, A.K., Bundela, P.S., Wong, J.W., Li, R., Zhang, Z.: Co-composting of gelatin industry sludge combined with organic fraction of municipal solid waste and poultry waste employing zeolite mixed with enriched nitrifying bacterial consortium. *Bioresour. Technol.* **213**, 181–189 (2016)
[Article](#) [Google Scholar](#)
6. Vaverková, M.D., Adamcová, D., Kotovicová, J., Toman, F.: Evaluation of biodegradability of plastics bags in composting conditions. *Ecol. Chem. Eng.* **S21**, 45–57 (2014)
[Google Scholar](#)

7. Jara-Samaniego, J., Pérez-Murcia, M.D., Bustamante, M.A., Pérez-Espinosa, A., Paredes, C., López, M., Moral, R.: Composting as sustainable strategy for municipal solid waste management in the Chimborazo Region, Ecuador: suitability of the obtained composts for seedling production. *J. Clean. Prod.* **141**, 1349–1358 (2017)

[Article](#) [Google Scholar](#)

8. Radziemska, M., Mazur, Z.: Effect of compost from by-product of the fishing industry on crop yield and microelement content in maize. *J. Ecol. Eng.* **16**, 168–175 (2015)

[Article](#) [Google Scholar](#)

9. Illera-Vives, M., Labandeira, S.S., Brito, L.M., López-Fabal, A., López-Mosquera, M.E.: Evaluation of compost from seaweed and fish waste as a fertilizer for horticultural use. *Sci. Hortic.* **186**, 101–107 (2015)

[Article](#) [Google Scholar](#)

10. EC Regulation 2092/91: Regulation 2092/91/EEC of 22 July 1991 on organic production of agricultural products and indications referring thereto on agricultural products and foodstuffs. *Off. J. Eur. Commun. L198*, **15**, 198 (1991)

[Google Scholar](#)

11. Castro, R.S., Borges Azevedo, C.M.S., Bezerra-Neto, F.: Increasing cherry tomato yield using fish effluent as irrigation water in Northeast Brazil. *Sci. Hortic.* **110**, 44–50 (2006)

[Article](#) [Google Scholar](#)

12. López-Mosquera, M.E., Fernández-Lema, E., Villares, R., Corral, R., Alonso, B., Blanco, C.: Composting fish waste and seaweed to produce a fertilizer for use in organic agriculture. *Proc. Environ. Sci.* **9**, 113–117 (2011)

[Article](#) [Google Scholar](#)

13. Zeng, G.M., Yu, M., Chen, Y.N., Huang, D.L., Zhang, J.C., Huang, H.L., Jiang, R.Q., Yu, Z.: Effects of inoculation with *Phanerochaete chrysosporium* at various time points on enzyme activities during agricultural waste composting. *Bioresour. Technol.* **101**, 222–227 (2010)

[Article](#) [Google Scholar](#)

14. Sun, D., Lan, Y., Xu, E.G., Meng, J., Chen, W.: Biochar as a novel niche for culturing microbial communities in composting. *Waste Manage.* **54**, 93–100 (2016)

[Article](#) [Google Scholar](#)

15. Voběrková, S., Vaverková, M.D., Burešová, A., Adamcová, D., Vršanská, M., Kynický, J., Brtnický, M., Adam, V.: Effect of inoculation with white-rot fungi and fungal consortium on the

composting efficiency of municipal solid waste. *Waste Manage.* **61**, 157–164 (2017)

[Article](#) [Google Scholar](#)

16. Belyaeva, O.N., Haynes, R.J., Sturm, E.C.: Chemical, physical and microbial properties and microbial diversity in manufactured soils produced from co-composting green waste and biosolids. *Waste Manage.* **32**, 2248–2257 (2012)

[Article](#) [Google Scholar](#)

17. Vandecasteele, B., Sinicco, T., D'Hose, T., Nest, V., Mondini, T.C.: Biochar amendment before or after composting affects compost quality and N losses, but not P plant uptake. *J. Environ. Manage.* **168**, 200–209 (2016)

[Article](#) [Google Scholar](#)

18. Saer, A., Lansing, S., Davitt, N.H., Graves, R.E.: Life cycle assessment of a food waste composting system: environmental impact hotspots. *J. Cleaner Prod.* **52**, 234–244 (2013)

[Article](#) [Google Scholar](#)

19. Mu, D., Horowitz, N., Casey, M., Jones, K.: Environmental and economic analysis of an in-vessel food waste composting system at Kean University in the U.S. *Waste Manage.* **59**, 476–486 (2017)

[Article](#) [Google Scholar](#)

20. Michalak, I., Chojnacka, K.: Algal compost—toward sustainable fertilization. *Rev. Inorg. Chem.* **33**, 161–172 (2013)

[Article](#) [Google Scholar](#)

21. Cui, H.-Y., Zhao, Y., Chen, Y.-N., Zhang, X., Wang, X.-Q., Lu, Q., Jia, L.-M., Wei, Z.-M.: Assessment of phytotoxicity grade during composting based on EEM/PARAFAC combined with projection pursuit regression. *J. Hazard. Mater.* **326**, 10–17 (2017)

[Article](#) [Google Scholar](#)

22. Phytotoxkit: Seed germination and early growth microbiotest with higher plants. Standard operation procedure, pp. 1–24. MicroBioTests Inc., Nazareth (2004)

[Google Scholar](#)

23. Zhang, L., Sun, X.Y.: Addition of fish pond sediment and rock phosphate enhances the composting of green waste. *Bioresour. Technol.* **233**, 116–126 (2017)

[Article](#) [Google Scholar](#)

24. Adamcová, D., Vaverková, M.D., Břoušková, E.: The toxicity of two types of sewage sludge from wastewater treatment plant for plants. J. Ecol. Eng. **17**, 33–37 (2016)

[Article](#) [Google Scholar](#)

25. Valerio, M.E., Garcia, J.F., Peinado, F.M.: Determination of phytotoxicity of soluble elements in soils, based on a bioassay with lettuce (*Lactuca sativa* L.). Sci. Total Environ. **378**, 63–66 (2007)

[Article](#) [Google Scholar](#)

26. Unuofin, F.O., Mnkeni, P.N.S.: Optimization of *Eisenia fetida* stocking density for the bioconversion of rock phosphate enriched cow dung-waste paper mixtures. Waste Manage. **34**, 2000–2006 (2014)

[Article](#) [Google Scholar](#)

27. StatSoft, Inc.: Electronic Statistics Textbook. StatSoft, Tulsa (2010).

<http://www.statsoft.com/textbook/>.

28. Rashad, F.M., Saleh, W.D., Moselhy, M.A.: Bioconversion of rice straw and certain agro-industrial wastes to amendments for organic farming systems: 1. Composting, quality, stability and maturity indices. Bioresour. Technol. **101**(15), 5952–5960 (2010)

[Article](#) [Google Scholar](#)

29. Aparna, C., Saritha, P., Himabindu, V., Anjaneyulu, Y.: Techniques for the evaluation of maturity for composts. Waste Manage. (2007). <https://doi.org/10.1016/j.wasman.2007.07.008>

[Google Scholar](#)

30. Aggelis, G., Ehaliotis, C., Nerud, F., Stoychev, I., Lyberatos, G., Zervakis, G.: Evaluation of white-rot fungi for detoxification and decolorization of effluents from the green olive debittering process. Appl. Microbiol, Biotechnol, **59**(2–3), 353–360 (2002)

[Google Scholar](#)

31. Han, W., Clarke, W., Pratt, S.: Composting of waste algae: a review. Waste Manage. **34**, 1148–1155 (2014)

[Article](#) [Google Scholar](#)

32. Piriz, M.L., Eyra, M.C., Rostagno, C.M.: Changes in biomass and botanical composition of beach-cast seaweeds in a disturbed coastal area from Argentine Patagonia. J. Appl. Phycol. **15**, 67–74 (2003)

[Article](#) [Google Scholar](#)

33. Illera-Vives, M., Seoane Labandeira, S., Iglesias Loureiro, L., López-Mosquerana M. E.:

Agronomic assessment of a compost consisting of seaweed and fish waste as an organic fertilizer for organic potato crops. *J. Appl. Phycol.* **29**, 1663–1671 (2017)

[Article](#) [Google Scholar](#)

34. Mazur, Z., Radziemska, M.: Influence of compost from fish by-products on nutrient supply in radish. *Ecol. Chem. Eng. A* **21**, 231–240 (2014)

[Google Scholar](#)

35. Krzebietke, S.: Response of butter lettuce (*Lactuca sativa* L.) to different forms of nitrogen fertilizers with chlorine and sulphates. *J. Elementol.* **13**, 581–588 (2008)

[Google Scholar](#)

36. Ekholm, P., Reinivuo, H., Mattila, P., Pakkala, H., Koponen, J., Happonen, A., Hellström, J., Ovaskainen, M.L.: Changes in the mineral and trace element contents of cereals, fruits and vegetables in Finland. *J. Food Compos. Anal.* **20**, 487–495 (2007)

[Article](#) [Google Scholar](#)

37. Francke, A.: The effect of potassium fertilization on the macronutrient content of pepino dulce (*Solanum muricatum* Ait.) fruit. *Acta Sci. Pol. Hortorum Cultus.* **9**, 51–57 (2010)

[Google Scholar](#)

38. Wierzbicka, B., Majkowska-Gadomska, J., Nowak, M.: Concentrations of some bionutrients in parthenocarpic cucumber fruits in forced cultivation. *Acta Sci. Pol. Hortorum Cultus* **6**, 3–8 (2007)

[Google Scholar](#)

39. Hernández, A., Castillo, H., Ojeda, D., Arras, A., López, J., Sánchez, E.: Effect of vermicompost and compost on lettuce production. *Chilean J. Agric. Res.* **70**, 583–589 (2010)

[Article](#) [Google Scholar](#)

40. Abou-El-Hassan, S., Desoky, A.H.: Effect of compost and compost tea on organic production of head lettuce. *J. Appl. Sci. Res.* **9**, 5650–5655 (2013)

[Google Scholar](#)

[Download references](#) ↓

Author information

Authors and Affiliations

1. Department of Environmental Improvement, Faculty of Civil and Environmental Engineering, Warsaw University of Life Sciences, Nowoursynowska 159, 02-776, Warsaw, Poland

Maja Radziemska

2. Department of Applied and Landscape Ecology, Faculty of AgriSciences, Mendel University in Brno, Zemědělská 1, 613 00, Brno, Czech Republic

Magdalena Daria Vaverková & Dana Adamcová

3. Department of Geology and Pedology, Faculty of Forestry and Wood Technology, Mendel University in Brno, Zemědělská 1/1665, 613 00, Brno, Czech Republic

Martin Brtnický

4. Department of Environmental Chemistry, Faculty of Environmental Management and Agriculture, University of Warmia and Mazury in Olsztyn, Pl. Łódzki 4, 10-727, Olsztyn, Poland

Zbigniew Mazur

Authors

1. Maja Radziemska

[View author publications](#)

You can also search for this author in [PubMed](#) [Google Scholar](#)

2. Magdalena Daria Vaverková

[View author publications](#)

You can also search for this author in [PubMed](#) [Google Scholar](#)

3. Dana Adamcová

[View author publications](#)

You can also search for this author in [PubMed](#) [Google Scholar](#)

4. Martin Brtnický

[View author publications](#)

You can also search for this author in [PubMed](#) [Google Scholar](#)

5. Zbigniew Mazur

[View author publications](#)

You can also search for this author in [PubMed](#) [Google Scholar](#)

Corresponding author

Correspondence to [Zbigniew Mazur](#).

Rights and permissions

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

[Reprints and Permissions](#)

About this article



Check for
updates

Cite this article

Radziemska, M., Vaverková, M.D., Adamcová, D. *et al.* Valorization of Fish Waste Compost as a Fertilizer for Agricultural Use. *Waste Biomass Valor* **10**, 2537–2545 (2019).


<https://doi.org/10.1007/s12649-018-0288-8>

[Download citation](#) 

- Received: 17 November 2017
- Accepted: 31 March 2018
- Published: 16 April 2018
- Issue Date: 01 September 2019
- DOI: <https://doi.org/10.1007/s12649-018-0288-8>

Keywords

- [Composting](#)
- [Fish waste](#)
- [Toxicity of compost](#)
- [Organic fertilizer](#)
- [Sinapis alba L.](#)
- [Lactuca sativa L.](#)



[Use our pre-submission checklist](#)


Avoid common mistakes on your manuscript.



Advertisement

search

Search by keyword or author



Search

Navigation

- [Find a journal](#)
- [Publish with us](#)

Discover content

- [Journals A-Z](#)
- [Books A-Z](#)

Publish with us

- [Publish your research](#)

- [Open access publishing](#)

Products and services

- [Our products](#)
- [Librarians](#)
- [Societies](#)
- [Partners and advertisers](#)

Our imprints

- [Springer](#)
- [Nature Portfolio](#)
- [BMC](#)
- [Palgrave Macmillan](#)
- [Apress](#)
- Your privacy choices/Manage cookies
- [Your US state privacy rights](#)
- [Accessibility statement](#)
- [Terms and conditions](#)
- [Privacy policy](#)
- [Help and support](#)

121.126.206.200

Not affiliated

© 2023 Springer Nature