




Instantly Bioavailable Nano Calcium and Magnesium with Nanogel Technology





Instantly Bioavailable Nano Calcium and Magnesium with Nanogel Technology

Gelmil Nano is instantly bioavailable oral anionic Calcium and Magnesium gel fortified with phosphorus and gluconeogenic precursors. Each bottle provides calcium equivalent to 5 bottles of I/V calcium. Serum calcium reaches to significant levels within 15-30 minutes and stays for extended periods. Self-dispensing squeezable bottle provides ease of administration.

Composition:

Each 300 gm Contains:

| Calcium | 40 gm |
|--------------------|----------|
| Phosphorus | 13 gm |
| Magnesium | 6 gm |
| Phyto bio- Calphos | |
| Vitamin D | 50000 IU |

Gluconeogenic precursors (Propylene glycol & Glycerol) eq. to Metabolizable energy value of 3027 Kilojoules

Benefits:

- Instantly bio available oral Calcium gel fortified with phosphorus and magnesium.
- Phytobio-Calphos significantly improves the serum level of Calcium, Phosphorus and Magnesium.
- Gluconeogenic precursors provide instant energy to alleviate negative energy balance.
- Self-dispensing squeezable bottle provides ease of administration.

Indications:

- Comprehensive nutrition supplement during milk Fever and for negative energy balance management.
- Reduce the chances of milk fever relapse after I/V Calcium administration.
- Recommended during anti-hemoprotozoan treatment, rumen impaction, uterine inertia, retention of placenta and to support normal parturition.

Recommended Usage:

- Calving support- Administer half bottle at the first sign of calving to support normal parturition, alleviate negative energy balance, reduce the chance of milk fever & associated complications.
- Lactation support- Administer half bottle on the first day followed by the balance quantity on the second day to alleviate negative energy balance and increase milk production.
- Milk Fever Management- Administer one bottle post parturition (upon observing milk fever symptoms or follow-up therapy after I/V calcium administration) by oral route, repeat after 24-48 hrs if required.

















