

Multiple Systems

Henry Yu

Contents

Chapter 1

Multiple Systems

Page 2

1.1 Classical Information

2

Classical states — 2 • Probabilistic states — 3 • Tensor products of vectors — 5

Chapter 1

Multiple Systems

1.1 Classical Information

1.1.1 Classical states

Suppose that we have two systems:

- X is a system having classical state set Σ .
- Y is a system having classical state set Γ .

Imagine that X and Y are placed side-by-side, with X on the left and Y on the right, and viewed together as if they form a single system.

We denote this new compound system by (X, Y) , or XY .

Question 1

What are the classical states of (X, Y) ?

Solution: The classical state set of (X, Y) is the **Cartesian product**:

$$\Sigma \times \Gamma = \{(x, y) \mid x \in \Sigma, y \in \Gamma\}$$

Note:

It's not important that X is on the left and Y is on the right; we could have just as well placed Y on the left and X on the right. The important thing is that the two systems are distinguishable.

Example 1.1.1 (Card suits)

If $\Sigma = \{0, 1\}$ and $\Gamma = \{\clubsuit, \diamond, \heartsuit, \spadesuit\}$, then:

$$\Sigma \times \Gamma = \{(0, \clubsuit), (0, \diamond), (0, \heartsuit), (0, \spadesuit), (1, \clubsuit), (1, \diamond), (1, \heartsuit), (1, \spadesuit)\}$$

This description generalizes to more than two systems in a natural way.

Suppose X_1, \dots, X_n are systems having classical state sets $\Sigma_1, \dots, \Sigma_n$, respectively.

The classical state set of the n -tuple (X_1, \dots, X_n) , viewed as a single compound system, is the Cartesian product:

$$\Sigma_1 \times \dots \times \Sigma_n = \{(x_1, \dots, x_n) \mid x_1 \in \Sigma_1, \dots, x_n \in \Sigma_n\}$$

Example 1.1.2

If $\Sigma_1 = \Sigma_2 = \Sigma_3 = \{0, 1\}$, then the classical state set of (X_1, X_2, X_3) is:

$$\Sigma_1 \times \Sigma_2 \times \Sigma_3 = \{(0, 0, 0), (0, 0, 1), (0, 1, 0), (0, 1, 1), (1, 0, 0), (1, 0, 1), (1, 1, 0), (1, 1, 1)\}$$

An n -tuple (x_1, \dots, x_n) may also be written as a string $x_1 \cdots x_n$.

Example 1.1.3 (Binary alphabet)

Suppose X_1, \dots, X_n are bits, so their classical state sets are all the same:

$$\Sigma_1 = \Sigma_2 = \dots = \Sigma_{10} = \{0, 1\}$$

The classical state set of (X_1, \dots, X_n) is the Cartesian product:

$$\Sigma_1 \times \Sigma_2 \times \dots \times \Sigma_{10} = \{0, 1\}^{10}$$

You can also think about this as a 10-bit register in a classical computer. Written as strings, these classical states look like this:

```
0000000000
0000000001
0000000010
0000000011
      ⋮
1111111111
```

Note:

Cartesian products of classical state sets are ordered lexicographically (i.e., in dictionary order):

- We assume the individual classical state sets are already ordered.
- Significance decreases from left to right.

Example 1.1.4

The Cartesian product $\{1, 2, 3\} \times \{0, 1\}$ is ordered like this:

$$(1, 0), (1, 1), (2, 0), (2, 1), (3, 0), (3, 1)$$

When n -tuples are written as strings and ordered in this way, we observe familiar patterns, such as $\{0, 1\} \times \{0, 1\}$ being ordered as 00, 01, 10, 11.

1.1.2 Probabilistic states

Probabilistic states of compound systems associates probabilities with the Cartesian product of the classical state sets of individual systems.

Example 1.1.5

This is a probabilistic state pair of bits (X, Y) :

$$Pr((X, Y) = (0, 0)) = \frac{1}{2}$$

$$Pr((X, Y) = (0, 1)) = 0$$

$$Pr((X, Y) = (1, 0)) = 0$$

$$Pr((X, Y) = (1, 1)) = \frac{1}{2}$$

An alternate notation using vector notation:

$$\begin{pmatrix} \frac{1}{2} \\ 0 \\ 0 \\ \frac{1}{2} \end{pmatrix} \begin{array}{l} \leftarrow \text{probability associated with state 00} \\ \leftarrow \text{probability associated with state 01} \\ \leftarrow \text{probability associated with state 10} \\ \leftarrow \text{probability associated with state 11} \end{array}$$

For a given probabilistic state of (X, Y) , we say that X and Y are **statistically independent** if

$$Pr((X, Y) = (x, y)) = Pr(X = x)Pr(Y = y)$$

for all $x \in \Sigma$ and $y \in \Gamma$.

Suppose that a probabilistic state of (X, Y) is expressed as a vector:

$$|\pi\rangle = \sum_{(x,y) \in \Sigma \times \Gamma} p_{ab} |xy\rangle$$

The systems X and Y are independent if there exist probability vectors

$$|\phi\rangle = \sum_{x \in \Sigma} q_x |x\rangle \quad \text{and} \quad |\psi\rangle = \sum_{y \in \Gamma} r_y |y\rangle$$

such that $p_{xy} = q_x r_y$ for all $x \in \Sigma$ and $y \in \Gamma$.

Example 1.1.6 (Independent bits)

The probabilistic state of a pair of bits (X, Y) represented by the vector

$$|\pi\rangle = \frac{1}{6} |00\rangle + \frac{1}{12} |01\rangle + \frac{1}{2} |10\rangle + \frac{1}{4} |11\rangle$$

is one in which X and Y are independent. The required condition is true for these probability vectors:

$$|\phi\rangle = \frac{1}{4} |0\rangle + \frac{3}{4} |1\rangle \quad \text{and} \quad |\psi\rangle = \frac{2}{3} |0\rangle + \frac{1}{3} |1\rangle$$

Example 1.1.7 (Dependent bits)

For the probabilistic state

$$\frac{1}{2} |00\rangle + \frac{1}{2} |11\rangle$$

of two bits (X, Y) , we have that X and Y are not independent.

If they were, we would have numbers q_0, q_1, r_0, r_1 such that

$$q_0 r_0 = \frac{1}{2}$$

$$q_0 r_1 = 0$$

$$q_1 r_0 = 0$$

$$q_1 r_1 = \frac{1}{2}$$

But if $q_0 r_1 = 0$, then either $q_0 = 0$ or $r_1 = 0$ (or both), contradicting either the first or last equality since any number multiplied by 0 is 0.

1.1.3 Tensor products of vectors

The tensor product of two vectors

$$|\phi\rangle = \sum_{x \in \Sigma} \alpha_x |x\rangle \quad \text{and} \quad |\psi\rangle = \sum_{y \in \Gamma} \beta_y |y\rangle$$

is the vector

$$|\phi\rangle \otimes |\psi\rangle = \sum_{(x,y) \in \Sigma \times \Gamma} \alpha_x \beta_y |xy\rangle$$

Equivalently, the vector $|\phi\rangle \otimes |\psi\rangle$ is defined by this condition:

$$\langle xy | \pi \rangle = \langle x | \phi \rangle \langle y | \psi \rangle \quad (\text{for all } x \in \Sigma \text{ and } y \in \Gamma)$$

Note:

In essence, this is the same operation for probabilistic states of a pair of independent bits; we are just giving a name to it now.