

Perfect Writeup: Assignment 13 Extra Credit

Henry Yu

Exercise 5

Assume that $a_1, \dots, a_n > 0, k > 0$, and $p > 1$. In the previous Exercise 4, it was found that $k \left(\sum_{j=1}^n a_j^{p/(p-1)} \right)^{(p-1)/p}$ is the maximum value of $f(x_1, \dots, x_n) = a_1x_1 + \dots + a_nx_n$ subject to $x_1^p + \dots + x_n^p = k^p$ and $x_1, \dots, x_n > 0$; this occurs when each $x_i = ka_i^{1/(p-1)} \left(\sum_{j=1}^n a_j^{p/(p-1)} \right)^{-1/p}$. Note how this obviously does not extend to the case $p = 1$. So here is that separate case $p = 1$:

Prove that $k \max \{a_1, \dots, a_n\}$ is the maximum value of $f(x_1, \dots, x_n) = a_1x_1 + \dots + a_nx_n$ subject to $x_1 + \dots + x_n = k$ where $k, a_1, \dots, a_n > 0$, in two different ways:

- (a) First prove this using only careful "common sense". Notice that the argument must have two parts: first, show how, given any a_1, \dots, a_n , corresponding values x_1, \dots, x_n can be chosen so that $x_1 + \dots + x_n = k$ and $a_1x_1 + \dots + a_nx_n = kA$; second, show that for any values x_1, \dots, x_n with $x_1 + \dots + x_n = k$, it is always true that $a_1x_1 + \dots + a_nx_n \leq kA$.
- (b) Second prove this by carefully showing that $A = \lim_{p \rightarrow 1^+} \left(\sum_{j=1}^n a_j^{p/(p-1)} \right)^{(p-1)/p}$.

(a) The first case is, given a_1, \dots, a_n , corresponding values x_1, \dots, x_n can be chosen. By setting all $a_i = A$, the maximum value of $f(x)$ occurs when x_1, \dots, x_n is as large as possible while still under the constraints of $x_1 + \dots + x_n = k$, so $x_i = \frac{k}{n}$.

$$\begin{aligned} f(x) &= x_1a_1 + \dots + x_na_n \\ &= a(x_1 + \dots + x_n) \\ &= Ak \\ &= kA. \end{aligned}$$

The second case is, given x_1, \dots, x_n with $x_1 + \dots + x_n = k$, it is always true that $a_1x_1 + \dots + a_nx_n \leq kA$. This means that not all $a_i = A$. Without loss of generality, suppose that $a_i = A$ and $a_i = 0$ for all $i > 1$,

$$\begin{aligned} f(x) &= A * x_1 + a_2x_2 + \dots + a_nx_n \\ &= Ak \\ &= kA. \end{aligned}$$

(b) We want to prove that $A = \lim_{p \rightarrow 1^+} \left(\sum_{j=1}^n a_j^{p/(p-1)} \right)^{(p-1)/p}$. To do this, we start by letting $q = \frac{p}{p-1}$, where $q \rightarrow +\infty$ iff $p \rightarrow 1^+$,

$$A = \lim_{q \rightarrow +\infty} \left(\sum_{j=1}^n a_j^q \right)^{\frac{1}{q}}.$$

Dividing both sides by A,

$$1 = \lim_{q \rightarrow +\infty} \left(\frac{a_1^q}{A^q} + \frac{a_2^q}{A^q} + \cdots + \frac{a_n^q}{A^q} \right)^{\frac{1}{q}}.$$

Let $b_j = \frac{a_j}{A}$. Now, taking the natural log of both sides,

$$\ln 1 = \lim_{q \rightarrow +\infty} \frac{\ln (b_1^q + \cdots + b_n^q)}{q}.$$

Taking the limit as q approaches $+\infty$, the expression goes to the value of $\ln 1$, which is 0.

$$\therefore A = \lim_{p \rightarrow 1^+} \left(\sum_{j=1}^n a_j^{p/(p-1)} \right)^{(p-1)/p}.$$