

EXPLORING DATA VIA INVARIANT COORDINATE SELECTION IN R, PYTHON, AND JULIA

Aurore Archimbaud, Colombe Becquart, Andreas Alfons,
Klaus Nordhausen & Valentin Todorov
joint w/ Anne M. Ruiz

ICS and Related Methods Conference
May 28, 2026



OUTLINE OF THE PRESENTATION

Invariant coordinate selection

- Motivation

- General principle

- Illustrations

- ICS - methodology

- Applications


ICS - implementations

Conclusion

Tutorial


INVARIANT COMPONENT SELECTION (ICS)

ICS Tyler et al. (2009) - GPCA Caussinus and Ruiz-Gazen (1990)



INVARIANT COMPONENT SELECTION (ICS)


ICS Tyler et al. (2009) - GPCA Caussinus and Ruiz-Gazen (1990)



- **Affine invariant** multivariate method.


INVARIANT COMPONENT SELECTION (ICS)

ICS Tyler et al. (2009) - GPCA Caussinus and Ruiz-Gazen (1990)



- **Affine invariant** multivariate method.
- Can recover the **Fisher's linear discriminant subspace** under some elliptical mixture models, without knowing the class labels (Tyler et al., 2009; Becquart et al., 2024).

EXAMPLE WITH 2 GROUPS - 2 DIMENSIONS




Linear Discriminant Analysis

- ▶ **Supervised** learning
- ▶ Finds a lower-dimensional subspace that maximizes the separation between different classes (FDS)
- ▶ Maximizes the ratio:

$$\arg \max_{v \in \mathbb{R}^p} \frac{v' B v}{v' W v}$$

EXAMPLE WITH 2 GROUPS - 2 DIMENSIONS




Linear Discriminant Analysis

- **Supervised** learning
- Finds a lower-dimensional subspace that maximizes the separation between different classes (FDS)
- Maximizes the ratio:

$$\arg \max_{v \in \mathbb{R}^p} \frac{v' B v}{v' W v}$$

EXAMPLE WITH 2 GROUPS - 2 DIMENSIONS



Invariant Coordinate Selection

- **Unsupervised** learning
- Finds a lower-dimensional subspace that maximizes the generalized kurtosis - it corresponds to the FDS in certain models.
- Maximizes the ratio:


$$\arg \max_{v \in \mathbb{R}^p} \frac{v' \text{COV}_4 v}{v' \text{COV} v}$$

EXAMPLE WITH 2 GROUPS - 2 DIMENSIONS


Invariant Coordinate Selection

- **Unsupervised** learning
- Finds a lower-dimensional subspace that maximizes the generalized kurtosis - it corresponds to the FDS in certain models.
- Maximizes the ratio:


$$\arg \max_{v \in \mathbb{R}^p} \frac{v' \text{COV}_4 v}{v' \text{COV} v}$$



TOY EXAMPLE FROM NORDHAUSEN AND RUIZ-GAZEN (2021)



ANOTHER TOY EXAMPLE



1. Components calculation


1. Two scatter matrices $\mathbf{V}_1$ and $\mathbf{V}_2$.
2. Eigendecomposition of $\mathbf{V}_1^{-1}\mathbf{V}_2$.
3. Decreasing eigenvalues: kurtosis.
4. Projection of the centered data on the eigenvectors $\rightarrow$ ICs.

1. Components calculation

1. V_1 : covariance matrix.
2. V_2 : fourth moment matrix.
3. Eigendecomposition of $V_1^{-1}V_2$
4. Projection of the centered data on the eigenvectors.


1. Components calculation

1. V_1 : **covariance** matrix.
2. V_2 : **fourth moment** matrix.
3. Eigendecomposition of $V_1^{-1}V_2$
4. Projection of the centered data on the eigenvectors.



1. Components calculation

1. V_1 : **covariance** matrix.
2. V_2 : **fourth moment** matrix.
3. Eigendecomposition of $V_1^{-1}V_2$
4. Projection of the centered data on the eigenvectors.




2. Components selection

1. Based on the eigenvalues.
2. Based on the components.
3. **First and/or last $k \leq p$ components retained.**

1. Components calculation

1. V_1 : **covariance** matrix.
2. V_2 : **fourth moment** matrix.
3. Eigendecomposition of $V_1^{-1}V_2$
4. Projection of the centered data on the eigenvectors.




2. Components selection

1. Based on the eigenvalues.
2. Based on the components.
3. **First and/or last $k \leq p$ components retained.**


1. Components calculation

1. V_1 : **covariance** matrix.
2. V_2 : **fourth moment** matrix.
3. Eigendecomposition of $V_1^{-1}V_2$
4. Projection of the centered data on the eigenvectors.



2. Components selection

1. Based on the eigenvalues.
2. Based on the components.
3. **First and/or last $k \leq p$ components retained.**



ICS - SUMMARY

Main characteristics

→ Only the **largest and/or smallest eigenvalues** are of interest to recover the FDS.

ICS - SUMMARY

Main characteristics

→ Only the **largest and/or smallest eigenvalues** are of interest to recover the FDS.

Challenges

- ▶ Choice of scatters
- ▶ Choice of components

ICS - SUMMARY

Main characteristics


→ Only the **largest and/or smallest eigenvalues** are of interest to recover the FDS.

Challenges

- ▶ Choice of scatters
- ▶ Choice of components

→ **Depends on the application.**

MAIN APPLICATIONS



APPLICATIONS

Outlier detection

1. Euclidian distance of the observations calculated using the k selected ICs.

APPLICATIONS

Outlier detection

1. Euclidian distance of the observations calculated using the k selected ICs.

Remark: $k = p \rightarrow$ Mahalanobis distance w.r.t. $\mathbf{V}_1$

APPLICATIONS

Outlier detection

1. Euclidian distance of the observations calculated using the *k selected ICs*.

Remark: $k = p \rightarrow$ Mahalanobis distance w.r.t. V_1

2. Cutoff based on quantiles from simulations.


APPLICATIONS

Outlier detection

1. Euclidian distance of the observations calculated using the *k* selected ICs.

Remark: $k = p \rightarrow$ Mahalanobis distance w.r.t. V_1

2. Cutoff based on quantiles from simulations.




APPLICATIONS

Outlier detection

1. Euclidian distance of the observations calculated using the k selected ICs.

Remark: $k = p \rightarrow$ Mahalanobis distance w.r.t. V_1

2. Cutoff based on quantiles from simulations.



Tandem clustering with ICS

1. Dimension reduction to retain k selected ICs.
2. Clustering method.
3. Identification of clusters.


APPLICATIONS

Outlier detection

1. Euclidian distance of the observations calculated using the k selected ICs.


Remark: $k = p \rightarrow$ Mahalanobis distance w.r.t. V_1

2. Cutoff based on quantiles from simulations.



Tandem clustering with ICS

1. Dimension reduction to retain k selected ICs.
2. Clustering method.
3. Identification of clusters.



OUTLINE OF THE PRESENTATION

Invariant coordinate selection

ICS - implementations

Ecosystem

Software design

R

Python

Julia

Conclusion

Tutorial

ICS ECOSYSTEM DEVELOPMENT



SOFTWARE DESIGN

Main common features:

- ▶ the `ICS` class;
- ▶ several scatter matrices;
- ▶ supported algorithms: `"whiten"`, `"standard"`, `"QR"`;
- ▶ same parameters: `fix_signs`, `center`.

SOFTWARE DESIGN

Main common features:

- ▶ the ICS class;
- ▶ several scatter matrices;
- ▶ supported algorithms: "whiten", "standard", "QR";
- ▶ same parameters: `fix_signs`, `center`.

Main differences:

- ▶ **R**: coexistence of S3 and S4 class system
- ▶ **Python**: computing the invariant coordinates is divided into `fit` and `transform` methods, in line with popular machine learning frameworks such as scikit-learn:
- ▶ **Julia**: same approach as Python with a `ICSModel` class.

ICS - ALGORITHMS

Let $\mathbf{X}_n = (\mathbf{x}_1, \dots, \mathbf{x}_n)'$ be a p -variate dataset.

Simultaneous diagonalization of two scatter matrices $\mathbf{S}_{1,n}$ and $\mathbf{S}_{2,n}$:

$$\mathbf{W}_n \mathbf{S}_{1,n} \mathbf{W}_n^\top = \mathbf{I}_p \quad \text{and} \quad \mathbf{W}_n \mathbf{S}_{2,n} \mathbf{W}_n^\top = \mathbf{D}_n$$

where the diagonal matrix $\mathbf{D}_n$ contains the eigenvalues

$\rho_1, \dots, \rho_p$ of $\mathbf{S}_{1,n}^{-1} \mathbf{S}_{2,n}$ in decreasing order and

$\mathbf{W}_n = (\mathbf{w}_1, \dots, \mathbf{w}_p)'$ contains the corresponding eigenvectors as its rows.

ICS - ALGORITHMS

Let $\mathbf{X}_n = (\mathbf{x}_1, \dots, \mathbf{x}_n)'$ be a p -variate dataset.

Simultaneous diagonalization of two scatter matrices $\mathbf{S}_{1,n}$ and $\mathbf{S}_{2,n}$:

$$\mathbf{W}_n \mathbf{S}_{1,n} \mathbf{W}_n^\top = \mathbf{I}_p \quad \text{and} \quad \mathbf{W}_n \mathbf{S}_{2,n} \mathbf{W}_n^\top = \mathbf{D}_n$$

where the diagonal matrix $\mathbf{D}_n$ contains the eigenvalues $\rho_1, \dots, \rho_p$ of $\mathbf{S}_{1,n}^{-1} \mathbf{S}_{2,n}$ in decreasing order and $\mathbf{W}_n = (\mathbf{w}_1, \dots, \mathbf{w}_p)'$ contains the corresponding eigenvectors as its rows.

The (affine) invariant coordinates or scores are:

$$\mathbf{Z}_n = \mathbf{X}_n^c \mathbf{W}_n^\top.$$

$\mathbf{X}_n^c$ can be the centered version of $\mathbf{X}_n$ with respect to the location estimator associated with $\mathbf{S}_{1,n}$.

ICS - ALGORITHMS

- ▶ `whiten`: whitens X_n with respect to $\mathbf{S}_1$ before computing $\mathbf{S}_2$ (should be a function).
- ▶ `standard`: performs the spectral decomposition of the symmetric matrix $M(X_n)$.

ICS - ALGORITHMS

- ▶ **whiten**: whitens X_n with respect to S_1 before computing S_2 (should be a function).
- ▶ **standard**: performs the spectral decomposition of the symmetric matrix $M(X_n)$.

whiten:

- ▶ $Y_n = X_n S_1(X_n)^{-1/2}$
- ▶ $S_2(Y_n)$
- ▶ $S_2(Y_n) = UDU'$
- ▶ $W = U' S_1(X_n)^{-1/2}$
- ▶ $Z = X_n^c W^\top$

standard:

- ▶ $S_1(X_n), S_2(X_n)$
- ▶ $M(X_n) = S_1(X_n)^{-1/2} S_2(X_n) S_1(X_n)^{-1/2}$
- ▶ $M(X_n) = UDU'$
- ▶ $W = U' S_1(X_n)^{-1/2}$
- ▶ $Z = X_n^c W^\top$

ICS - ALGORITHMS

The signs of W can be fixed with `fix_signs`:

- ▶ **"scores"**: the generalized skewness values of all components are positive.
Common for **ICS** framework.
- ▶ **"W"**: the maximum element in each row of W is positive and each row has norm 1.
Common with **ICA** framework.

Z can be centered through the boolean `center` input parameter.

ICS - **QR** - BASED ON PIVOTED QR DECOMPOSITION

Archimbaud et al. (2023b), numerically stable algorithm for $\text{COV} - \text{COV}_w$ focusing on:

$$M(X_n) = \text{COV}^{-1/2} \text{COV}_w \text{COV}^{-1/2}.$$

ICS - **QR** - BASED ON PIVOTED QR DECOMPOSITION

Archimbaud et al. (2023b), numerically stable algorithm for COV – COV_w focusing on:

$$M(X_n) = \text{COV}^{-1/2} \text{COV}_w \text{COV}^{-1/2}.$$

Algorithm:

- ▶ Pivoted QR factorization: $\Pi_2^\top \left(\frac{1}{\sqrt{n-1}} X_n^c \right) \Pi_1 = QR$
- ▶ $Q = \Pi_2 Q, R = R \Pi_1^\top$
- ▶ Leverage scores $q_i = \|Q(i, :)\|_2^2, i = 1, \dots, n$
- ▶ SVD of $\text{Diag} \left(\sqrt{w((n-1)q_i)} \right)_{i=1}^n Q$ to obtain

$$\tilde{M}(X_n) = \frac{n-1}{n} Q^\top \text{Diag}(w((n-1)q_i))_{i=1}^n Q = \tilde{U}_2 \tilde{D}_2 \tilde{U}_2^\top$$

- ▶ $W = (R^{-1} \tilde{U}_2)^\top$
- ▶ $Z = \sqrt{n-1} Q \tilde{U}_2$ or equivalently $Z = X_n^c \Pi_1 W^\top$.

ICS - SUMMARY

Main attributes

Language	R	R	Python	Julia
Package	ICS - <code>ics()</code> , <code>ics2()</code>	ICS - <code>ICS()</code>	ICSpyLab	ICSTools.jl
Class	'ICS'	'ICS'	'ICS'	'ICSModel'
Generalized eigenvalues	@gKurt	\$gen.kurtosis, <code>kurtosis_</code> <code>gen.kurtosis()</code>	<code>kurtosis_</code>	<code>kurtosis_</code>
Unmixing matrix	@UnMix	\$W	<code>components_</code>	<code>W_</code>
Scores	@Scores	\$scores, <code>components()</code>		<code>scores_</code>


ICS - SUMMARY

Methods

Language Package	R ICS	Python ICSpyLab	Julia ICSTools.jl
Print basic information	<code>print()</code>		
Summary	<code>summary()</code>	<code>describe()</code>	<code>show()</code>
Coefficient matrix of ICS	<code>coef()</code>		
Plots the kurtosis measures	<code>screeplot()</code>	<code>plot_kurtosis()</code>	<code>scree_plot()</code>
Scatterplot of ICs	<code>plot()</code>		
Extract the transformed data matrix	<code>fitted()</code>		
Transform data using the fitted transformation matrix		<code>transform()</code>	<code>predict()</code>
Fit and transform		<code>fit_transform()</code>	<code>fit_predict!()</code>
Transform data back to its original space		<code>inverse_transform()</code>	

CHOICE OF SCATTER MATRICES

- ▶ $\text{COV} - \text{COV}_4$ or FOBI Peña et al. (2010); Cardoso (1989); Nordhausen and Virta (2019); Fischer et al. (2017a, 2020)
- ▶ $\text{MCD}_\alpha - \text{COV}$, $\text{RMCD}_\alpha - \text{COV}$ with values of $\alpha \leq 0.5$ Rousseeuw (1985)
- ▶ $\text{TCOV} - \text{UCOV}$, $\text{TCOV} - \text{COV}$ Caussinus and Ruiz-Gazen (1993a)
- ▶ $\text{LCOV} - \text{COV}$ Hennig (2009)



CHOICE OF SCATTER MATRICES

Applications:

- For **outlier detection**: Archimbaud et al. (2018a) for a small proportion of outliers.

CHOICE OF SCATTER MATRICES

Applications:

- ▶ For **outlier detection**: Archimbaud et al. (2018a) for a small proportion of outliers.
- ▶ For **clustering**: Becquart et al. (2024) and Alfons et al. (2024). One scatter matrix should reflect the **within (local) cluster structure** and one the **between (global) cluster structure**.

CLASS AND RELEVANT COMPONENTS FOR SCATTER MATRICES

Language Package	R ICS	Python ICSpyLab	Julia ICSTools.jl	
Class	'ICS_scatter'	'Scatter'	'Scatter'	
Components	location	location	location	Estimated location vector (optional)
	scatter	scatter	scatter	Estimated scatter matrix
	label	label	label	Label for the scatter matrix (used for printing output)

Multivariate ICS		ICS - implementations		Conclusion	Tutorial
Ecosystem		Software design		Python	Julia
Language Class		R	R 'ICS_scatter'	Python 'Scatter'	Julia 'Scatter'
Covariance		<code>ICS::cov()</code>	<code>ICS::ICS_cov()</code>	<code>cov()</code>	<code>cov2()</code>
Fourth-moment co-variance	<code>co-</code>	<code>ICS::cov4()</code>	<code>ICS::ICS_cov4()</code>	<code>cov4()</code>	<code>cov4()</code>
One-step estimator	<code>M-</code>	<code>ICS::covW()</code>	<code>ICS::ICS_covW()</code>	<code>covW()</code>	<code>covW()</code>
One-step Tyler shape matrix		<code>ICS::covAxis()</code>	<code>ICS::ICS_covAxis()</code>	<code>covAxis()</code>	<code>covAxis()</code>
Multivariate t-distribution estimator	<code>t-</code>	<code>ICS::tM()</code>	<code>ICS::ICS_tM()</code>	<code>tM()</code>	<code>tM()</code>
Supervised (quantiles)	<code>scatter</code>	<code>ICS::scovq()</code>	<code>ICS::ICS_scovq()</code>		<code>mlc()</code>
MCD		<code>rrcov::CovMcd()</code>	<code>ICSClust::ICS_mcd_raw(),</code> <code>ICS_mcd_rwt()</code>	<code>mcd()</code>	<code>mcd_raw(),</code> <code>mcd_rwt()</code>
Cauchy location and scatter		<code>ICS::tM()</code>	<code>ICSClust::ICS_mlc()</code>		
Pairwise one-step M-estimate		<code>ICSClust::tcov()</code>	<code>ICSClust::ICS_tcov()</code>	<code>tcov()</code>	<code>tcov()</code>
Simple robust estimates		<code>ICSClust::ucov()</code>	<code>ICSClust::ICS_ucov()</code>		
Local shape scatter		<code>ICSClust::lcov()</code>	<code>ICSClust::ICS_lcov()</code>		<code>lcov()</code>

CHOICE OF COMPONENTS WITH $q < p$

Archimbaud et al. (2018a), Alfons et al. (2024), Nordhausen et al. (2022), Radojicic and Nordhausen (2020)

CHOICE OF COMPONENTS WITH $q < p$

Archimbaud et al. (2018a), Alfons et al. (2024), Nordhausen et al. (2022), Radojicic and Nordhausen (2020)

Based on the Invariant Components:

- Keep only non-gaussian components using marginal normality tests.

CHOICE OF COMPONENTS WITH $q < p$

Archimbaud et al. (2018a), Alfons et al. (2024), Nordhausen et al. (2022), Radojicic and Nordhausen (2020)

Based on the Invariant Components:

- Keep only non-gaussian components using marginal normality tests.

In this context of particular sequential multiple testing, we apply the Bonferroni correction on the significance level: $\alpha_i = \alpha/i$ for $i = 1, \dots, p$ with $\alpha = 5\%$ (see Dray (2008)).

CHOICE OF COMPONENTS WITH $q < p$

Based on the eigenvalues and eigenvectors:

- ▶ Visually, using a **scree plot**.
- ▶ Using the number q of clusters a priori and the fact that some should be almost equal (**med_criterion** and **var_criterion**).
- ▶ Using asymptotic distribution of the eigenvalues or bootstrapping (**ICSboot**, **FOBIasymp**, **FOBIboot**).
- ▶ Using quasi inferential procedures (**parallel analysis**).
- ▶ Using ladle and data augmentation (**FOBIladle**).

CHOICE OF COMPONENTS WITH $q < p$

For outlier detection:

- ▶ **First** components

CHOICE OF COMPONENTS WITH $q < p$

For outlier detection:

- ▶ **First** components


For clustering:

- ▶ **First** and/or **last** components

SELECTION OF COMPONENTS - SUMMARY IN R


Criterion		Function	Notes
Marginal tests	normality	<code>ICSClust::normal_crit()</code>	First and last few components are investigated
Median-based criterion		<code>ICSClust::med_crit()</code>	Requires a priori knowledge
Variance-based criterion		<code>ICSClust::var_crit()</code>	From <code>ICtest::ICSboot()</code>
Discriminatory power criterion		<code>ICSClust::discriminatory_crit()</code>	Supervised, requires a priori knowledge
Parallel analysis (simulation)		<code>ICSOutlier::comp_simu_test()</code> , <code>comp.simu.test()</code>	Only first components are investigated
Marginal tests	normality	<code>ICSOutlier::comp_norm_test()</code> , <code>comp.norm.test()</code>	Only first components are investigated
Asymptotic and resampling based approaches		<code>ICtest::ICSboot()</code> , <code>FOBIboot()</code> , <code>FOBIasympt()</code> , <code>FOBIladle()</code>	Focus on identifying Gaussian subspace

GENESIS: 2007/2008 - KLAUS NORDHAUSEN + HANNU OJA + DAVID E. TYLER




2016-2018: + AURORE ARCHIMBAUD AND ANNE RUIZ-GAZEN

+ Joris May



2016-2018: + AURORE ARCHIMBAUD AND ANNE RUIZ-GAZEN

+ Joris May



2023: + ANDREAS ALFONS + ZLATKO DRMAČ

ICSClust (v1.0)

ICSClust(S3)

ICS_scatter: ICS_lcov, ICS_med, ICS_mlc, ICS_teov, ICS_ucov

1. *Computation of ICS*: ICS

2. *Selection of ICS and visualization*:

"med_crit", "normal_crit", "var_crit", "discriminatory_crit"

3. *Clustering and visualization*:

"kmeans_clust", "tkmeans_clust", "pam_clust", "mclust_clust",
"rmclust_clust", "rimle_clust"

ICSOutlier (v4.0)

ICS_outlier(S3)

0. *Computation of ICS*: ICS

1. *Selection of ICS*: comp_simu_test comp_norm_test

2. *ICS distances*: ics_distances

3. *Identification of outliers*: dist_simu_test

ICS (v1.4-2)

```
ICS(
  X,
  S1 = ICS_cov,
  S2 = ICS_cov4,
  S1_args = list(),
  S2_args = list(),
  algorithm = c("whiten", "standard", "QR"),
  center = FALSE,
  fix_signs = c("scores", "W"),
  na.action = na.fail
)
```

```
@gKurt   → $gen_kurtosis
@UnMix   → $W
@Scores  → $scores
```

ICS (S3)


1. *ICS_scatter* class

2. *Different algorithms*

3. *Centering*

4. *Fixing signs*

ICS - R ECOSYSTEM



PYTHON - 2023: COLOMBE BECQUART + ABDALLAH ABDELSAMEIA

Package **ICSpyLab**

- ▶ version 0.1.0: May 2025
- ▶ version 1.0.0: May 2026
- ▶ Main features:
 - ▶ the ICS class;
 - ▶ several scatter matrices;
 - ▶ supported algorithms:
"whiten", "standard",
"QR" and "eigh".
 - ▶ same parameters: fix_signs,
center.



JULIA 2026: VALENTIN TODOROV

Package **ICSTools.jl**

- ▶ Main features:
 - ▶ the `ICSTools` class;
 - ▶ several scatter matrices (**Robustbase.jl**);
 - ▶ supported algorithms: "whiten", "standard", "QR";
 - ▶ same parameters: `fix_signs`, `center`.



OUTLINE OF THE PRESENTATION

Invariant coordinate selection

ICS - implementations

Conclusion

Tutorial

CONCLUSION

ICS is an attractive **unsupervised multivariate method:**

- ▶ Designed to **find structure** in a low-dimensional subspace:
 - ▶ for outlier detection,
 - ▶ as pre-processing for clustering
- ▶ Affine invariant.

CONCLUSION

ICS is an attractive **unsupervised multivariate method:**

- ▶ Designed to **find structure** in a low-dimensional subspace:
 - ▶ for outlier detection,
 - ▶ as pre-processing for clustering
- ▶ Affine invariant.

Still presents some challenges:

- ▶ Choice of scatter matrices.
- ▶ Choice of components.

SOME PERSPECTIVES

Implementation:

- ▶ **R ecosystem**: algorithms for functional, compositional, collinear or HDLSS data.
- ▶ **Python** and **Julia**: utilities for outlier detection and clustering

SOME PERSPECTIVES

Implementation:

- ▶ **R ecosystem**: algorithms for functional, compositional, collinear or HDLSS data.
- ▶ **Python** and **Julia**: utilities for outlier detection and clustering

Github repository:

- ▶ <https://github.com/AuroreAA/ICS-implementation>

Working paper:

- ▶ Alfons, A., Archimbaud, A., Becquart, C., Nordhausen, K., & Todorov, V. (2026). *Exploring data via invariant coordinate selection in R, Python, and Julia*.

All contributions are welcomed.

OUTLINE OF THE PRESENTATION

Invariant coordinate selection

ICS - implementations

Conclusion

Tutorial

REFERENCES I

- Alashwali, F. and Kent, J. T. (2016). The use of a common location measure in the invariant coordinate selection and projection pursuit. Journal of Multivariate Analysis, 152:145–161.
- Alfons, A., Archimbaud, A., Nordhausen, K., and Ruiz-Gazen, A. (2024). Tandem clustering with invariant coordinate selection. Econometrics and Statistics.
- Archimbaud, A. (2026). Generalized implementation of invariant coordinate selection with positive semi-definite scatter matrices. Journal of Multivariate Analysis, 211:105520.
- Archimbaud, A., Alfons, A., Nordhausen, K., and Ruiz-Gazen, A. (2023a). ICSCluster: Tandem Clustering with Invariant Coordinate Selection. R package version 0.1.0.
- Archimbaud, A., Boulfani, F., Gendre, X., Nordhausen, K., Ruiz-Gazen, A., and Virta, J. (2025). ICS for multivariate functional anomaly detection with applications to predictive maintenance and quality control. Econometrics and Statistics, 33:282–303.
- Archimbaud, A., Drmač, Z., Nordhausen, K., Radojičić, U., and Ruiz-Gazen, A. (2023b). Numerical considerations and a new implementation for ICS. SIAM Journal on Mathematics of Data Science (SIMODS), 5(1):97–121.
- Archimbaud, A., Nordhausen, K., and Ruiz-Gazen, A. (2018a). ICS for multivariate outlier detection with application to quality control. Computational Statistics & Data Analysis, 128:184–199.
- Archimbaud, A., Nordhausen, K., and Ruiz-Gazen, A. (2018b). ICSCluster: Unsupervised outlier detection for low-dimensional contamination structure. The R Journal, 10(1):234–250.
- Arias, V. B., Garrido, L., Jenaro, C., Martínez-Molina, A., and Arias, B. (2020). A little garbage in, lots of garbage out: Assessing the impact of careless responding in personality survey data. Behavior Research Methods, 52:2489–2505.
- Becquart, C. and Abdelsameia, A. (2025). ICSPyLab: Python implementation of invariant coordinate selection. <https://icspylab.readthedocs.io/>. Python package version 0.1.0.
- Becquart, C., Archimbaud, A., Ruiz-Gazen, A., Prilč, L., and Nordhausen, K. (2024). Invariant coordinate selection and fisher discriminant subspace beyond the case of two groups. arXiv preprint arXiv:2409.17631.

REFERENCES II

- Becquart, C., Archimbaud, A., Ruiz-Gazen, A., Prilć, L., and Nordhausen, K. (2026). Invariant coordinate selection and fisher discriminant subspace beyond the case of two groups. Journal of Multivariate Analysis, 211:105521.
- Cardoso, J.-F. (1989). Source separation using higher order moments. In Proceedings of the IEEE International Conference on Acoustics, Speech and Signal Processing, pages 2109–2112. IEEE.
- Caussinus, H., Fekri, M., Hakam, S., and Ruiz-Gazen, A. (2003). A monitoring display of multivariate outliers. Computational Statistics & Data Analysis, 44(1):237–252.
- Caussinus, H. and Ruiz-Gazen, A. (1990). Interesting projections of multidimensional data by means of generalized principal component analyses. In Compstat, pages 121–126. Springer.
- Caussinus, H. and Ruiz-Gazen, A. (1993a). Projection pursuit and generalized principal component analysis. In Morgenthaler, S., Ronchetti, E., and Stahel, W. A., editors, New directions in statistical data analysis and robustness., Monte Verita, Proceedings of the Centro Stefano Franciscini Ascona Series. Springer.
- Caussinus, H. and Ruiz-Gazen, A. (1993b). Projection pursuit and generalized principal component analysis. New Directions in Statistical Data Analysis and Robustness, pages 35–46.
- Caussinus, H. and Ruiz-Gazen, A. (1995). Metrics for finding typical structures by means of principal component analysis. In Data Sci. and its Applications, pages 177–192, Japan. Harcourt Brace.
- Caussinus, H. and Ruiz-Gazen, A. (2007). Classification and generalized principal component analysis. In Selected contributions in data analysis and classification, pages 539–548. Springer.
- Chakraborty, B. and Chaudhuri, P. (1998). On an adaptive transformation-retransformation estimate of multivariate location. Journal of the Royal Statistical Society. Series B, 60(1):145–157.
- Chakraborty, B., Chaudhuri, P., and Oja, H. (1998). Operating transformation retransformation on spatial median and angle test. Statistica Sinica, 8(3):767–784.
- Claassen, C. (2023). Generalising invariant coordinate selection to a non-linear dimensionality reduction method. Master's thesis.

REFERENCES III

- Dray, S. (2008). On the number of principal components: A test of dimensionality based on measurements of similarity between matrices. Computational Statistics and Data Analysis, 52(4):2228 – 2237.
- Dümbgen, L., Gysel, K., and Perler, F. (2021). Refining invariant coordinate selection via local projection pursuit. arXiv preprint arXiv:2112.11998.
- Fekri, M. and Ruiz-Gazen, A. (2015). A B-robust non-iterative scatter matrix estimator: Asymptotics and application to cluster detection using invariant coordinate selection. In Nordhausen, K. and Taskinen, S., editors, Modern Nonparametric, Robust and Multivariate Methods: Festschrift in Honour of Hannu Oja, pages 395–423. Springer International Publishing, Cham.
- Fischer, D., Honkatukia, M., Tuiskula-Haavisto, M., Nordhausen, K., Cavero, D., Preisinger, R., and Vilkki, J. (2017a). Subgroup detection in genotype data using invariant coordinate selection. BMC Bioinformatics, 18:173–181.
- Fischer, D., Honkatukia, M., Tuiskula-Haavisto, M., Nordhausen, K., Cavero, D., Preisinger, R., and Vilkki, J. (2017b). Subgroup detection in genotype data using invariant coordinate selection. BMC Bioinformatics, 18:173–181.
- Fischer, D., Nordhausen, K., and Oja, H. (2020). On linear dimension reduction based on diagonalization of scatter matrices for bioinformatics downstream analyses. Heliyon, 6:e05732.
- Goldberg, L. R. (1992). The development of markers for the big-five factor structure. Psychological assessment, 4(1):26.
- Heinonen, L. and Virta, J. (2026). A method for sparse and robust independent component analysis. Journal of Multivariate Analysis, 213:105587.
- Hennig, C. (2009). Discussion of “Invariant Co-ordinate Selection”, by D. E. Tyler, F. Critchley, L. Dümbgen, and H. Oja. Journal of the Royal Statistical Society B, 71:579–583.
- Hyvärinen, A., Karhunen, J., and Oja, E. (2001). Independent Component Analysis. John Wiley & Sons.
- Liski, E., Nordhausen, K., and Oja, H. (2014). Supervised invariant coordinate selection. Statistics: A Journal of Theoretical and Applied Statistics, 4:711–731.

REFERENCES IV

- Miettinen, J., Taskinen, S., Nordhausen, K., and Oja, H. (2015). Fourth moments and independent component analysis. Statistical Science, 30(3):372–390.
- Mondon, C., Trinh, H. T., Ruiz-Gazen, A., and Thomas-Agnan, C. (2025). Ics for complex data with application to outlier detection for density data. Journal of Multivariate Analysis, page 105522.
- Nordhausen, K., Alfons, A., Archimbaud, A., Oja, H., Ruiz-Gazen, A., and Tyler, D. E. (2023). ICS: Tandem Clustering with Invariant Coordinate Selection. R package version 1.4-2.
- Nordhausen, K. and Oja, H. (2018a). Independent component analysis: A statistical perspective. WIREs: Computational Statistics, 10:e1440.
- Nordhausen, K. and Oja, H. (2018b). Robust nonparametric inference. Annual Review of Statistics and Its Application, 5:473–500.
- Nordhausen, K., Oja, H., and Ollila, E. (2008a). Robust independent component analysis based on two scatter matrices. Austrian Journal of Statistics, 37:91–100.
- Nordhausen, K., Oja, H., and Tyler, D. E. (2006). On the efficiency of invariant multivariate sign and rank test. In Liski, E. P., Isotalo, J., Niemelä, J., Puntanen, S., and Styan, G. P. H., editors, Festschrift for Tarmo Pukkila on his 60th Birthday, pages 217–231. University of Tampere, Tampere.
- Nordhausen, K., Oja, H., and Tyler, D. E. (2008b). Tools for exploring multivariate data: The package ICS. Journal of Statistical Software, Articles, 28(6):1–31.
- Nordhausen, K., Oja, H., and Tyler, D. E. (2022). Asymptotic and Bootstrap Tests for Subspace Dimension. Journal of Multivariate Analysis, 188:104830.
- Nordhausen, K., Oja, H., Tyler, D. E., and Virta, J. (2017). Asymptotic and bootstrap tests for the dimension of the non-gaussian subspace. IEEE Signal Processing Letters, 24(6):887–891.
- Nordhausen, K. and Ruiz-Gazen, A. (2021). On the usage of joint diagonalization in multivariate statistics. Journal of Multivariate Analysis, page 104844.

REFERENCES V

- Nordhausen, K. and Virta, J. (2019). An overview of properties and extensions of FOBI. Knowledge-Based Systems, 173:113–116.
- Oja, H., Sirkiä, S., and Eriksson, J. (2006). Scatter matrices and independent component analysis. Austrian Journal of Statistics, 35:175–189.
- Peña, D., Prieto, F. J., and Viladomat, J. (2010). Eigenvectors of a kurtosis matrix as interesting directions to reveal cluster structure. Journal of Multivariate Analysis, 101(9):1995–2007.
- Peres-Neto, P. R., Jackson, D. A., and Somers, K. M. (2005). How many principal components? stopping rules for determining the number of non-trivial axes revisited. Computational Statistics and Data Analysis, 49(4):974 – 997.
- Radojicic, U. and Nordhausen, K. (2020). Non-gaussian component analysis: Testing the dimension of the signal subspace. In Maciak, M., Pesta, M., and Schindler, M., editors, Analytical Methods in Statistics. AMISTAT 2019, pages 101–123. Springer, Cham.
- Rousseeuw, P. J. (1985). Multivariate estimation with high breakdown point. Mathematical statistics and applications, 8(283-297):37.
- Rousseeuw, P. J., Raymaekers, J., and Hubert, M. (2018). A measure of directional outlyingness with applications to image data and video. Journal of Computational and Graphical Statistics, 27(2):345–359.
- Ruiz-Gazen, A. (1996). A very simple robust estimator of a dispersion matrix. Computational Statistics & Data Analysis, 21:149–162.
- Ruiz-Gazen, A., Thomas-Agnan, C., Laurent, T., and Mondon, C. (2022). Detecting outliers in compositional data using invariant coordinate selection. In Robust and Multivariate Statistical Methods: Festschrift in Honor of David E. Tyler, pages 197–224. Springer.
- Tyler, D. E., Critchley, F., Dümbgen, L., and Oja, H. (2009). Invariant coordinate selection. Journal of the Royal Statistical Society: Series B, 71(3):549–592.

NOTATIONS

- ▶ $\mathbf{X}$ is a p -dimensional random vector, $F_{\mathbf{X}}$ its cumulative distribution function and $\mathbf{m}(F_{\mathbf{X}})$ an affine equivariant location estimator.
- ▶ $\mathbf{X}_n = (\mathbf{x}_1, \dots, \mathbf{x}_n)'$ a p -variate dataset. $\mathbf{x}_1, \dots, \mathbf{x}_n$ follow the same distribution as $\mathbf{X}$.

NOTATIONS

- ▶ $\mathbf{X}$ is a p -dimensional random vector, $F_{\mathbf{X}}$ its cumulative distribution function and $\mathbf{m}(F_{\mathbf{X}})$ an affine equivariant location estimator.
- ▶ $\mathbf{X}_n = (\mathbf{x}_1, \dots, \mathbf{x}_n)'$ a p -variate **dataset**. $\mathbf{x}_1, \dots, \mathbf{x}_n$ follow the same distribution as $\mathbf{X}$.
- ▶ $\mathcal{P}_p$ is the set of all symmetric positive definite matrices of order p .
- ▶ A **scatter functional** is defined as a matrix $\mathbf{S}(F_{\mathbf{X}}) \in \mathcal{P}_p$, uniquely defined at $F_{\mathbf{X}}$, which is **affine equivariant** in the sense that:

$$\mathbf{S}(F_{\mathbf{A}\mathbf{X}+\boldsymbol{\gamma}}) = \mathbf{A}\mathbf{S}(F_{\mathbf{X}})\mathbf{A}',$$

for all $p \times p$ non-singular matrices $\mathbf{A}$ and all $\boldsymbol{\gamma} \in \mathbb{R}^p$.

For sake of simplicity, the dependence on $F_{\mathbf{X}}$ is dropped.

CHOICE OF SCATTER MATRICES I

- $\text{COV} - \text{COV}_4$ or FOBI Peña et al. (2010); Cardoso (1989); Nordhausen and Virta (2019); Fischer et al. (2017a, 2020)

$$\text{COV}_4(\mathbf{X}_n) = \frac{1}{n} \sum_{i=1}^n r^2(\mathbf{x}_i)(\mathbf{x}_i - \bar{\mathbf{x}}_n)(\mathbf{x}_i - \bar{\mathbf{x}}_n)^\top,$$

where $r^2(\mathbf{x}_i) = (\mathbf{x}_i - \bar{\mathbf{x}}_n)^\top \text{COV}(\mathbf{x}_n)^{-1}(\mathbf{x}_i - \bar{\mathbf{x}}_n)$ is the squared Mahalanobis distance.

CHOICE OF SCATTER MATRICES II

- $\text{MCD}_\alpha - \text{COV}$ and $\text{RMCD}_\alpha - \text{COV}$: (Reweighted) Minimum Covariance Determinant **Rousseeuw (1985)**.

$$\text{MCD}_\alpha(\mathbf{X}_n) = c_\alpha \frac{1}{n_\alpha} \sum_{j=1}^{n_\alpha} (\mathbf{x}_{i_j} - \bar{\mathbf{x}}_{\alpha,n})(\mathbf{x}_{i_j} - \bar{\mathbf{x}}_{\alpha,n})^\top,$$

where $n_\alpha = \lceil \alpha n \rceil$ observations for which the sample covariance matrix has the smallest determinant and usually $\alpha \in [0.5, 1]$.

In our ecosystem it is also interesting to consider **values of α that are smaller than 0.5**.

CHOICE OF SCATTER MATRICES III




Figure: Shapes of different scatter matrices for a sample from a bivariate Gaussian mixture distribution with three balanced clusters.

CHOICE OF SCATTER MATRICES IV

- $SCOV - COV$, $TCOV - UCOV$ Caussinus and Ruiz-Gazen (1993a, 1995, 2007); Ruiz-Gazen (1996); Caussinus and Ruiz-Gazen (1990); Fekri and Ruiz-Gazen (2015)

CHOICE OF SCATTER MATRICES IV

- **SCOV – COV, TCOV – UCOV** Caussinus and Ruiz-Gazen (1993a, 1995, 2007); Ruiz-Gazen (1996); Caussinus and Ruiz-Gazen (1990); Fekri and Ruiz-Gazen (2015) $\Rightarrow$ **TCOV – COV**:

$$\text{TCOV}_{\beta}(\mathbf{X}_n) = \frac{\sum_{i=1}^{n-1} \sum_{j=i+1}^n w(\beta r^2(\mathbf{x}_i, \mathbf{x}_j)) (\mathbf{x}_i - \mathbf{x}_j)(\mathbf{x}_i - \mathbf{x}_j)^{\top}}{\sum_{i=1}^{n-1} \sum_{j=i+1}^n w(\beta, r^2(\mathbf{x}_i, \mathbf{x}_j))},$$

where $r^2(\mathbf{x}_i, \mathbf{x}_j) = (\mathbf{x}_i - \mathbf{x}_j)^{\top} \text{COV}(\mathbf{x}_n)^{-1} (\mathbf{x}_i - \mathbf{x}_j)$,
 $w(x) = \exp(-x/2)$, $\beta = 4$.

CHOICE OF SCATTER MATRICES IV

- **SCOV – COV, TCOV – UCOV** Caussinus and Ruiz-Gazen (1993a, 1995, 2007); Ruiz-Gazen (1996); Caussinus and Ruiz-Gazen (1990); Fekri and Ruiz-Gazen (2015) $\Rightarrow$ TCOV – COV:

$$\text{TCOV}_{\beta}(\mathbf{X}_n) = \frac{\sum_{i=1}^{n-1} \sum_{j=i+1}^n w(\beta r^2(\mathbf{x}_i, \mathbf{x}_j)) (\mathbf{x}_i - \mathbf{x}_j)(\mathbf{x}_i - \mathbf{x}_j)^{\top}}{\sum_{i=1}^{n-1} \sum_{j=i+1}^n w(\beta, r^2(\mathbf{x}_i, \mathbf{x}_j))},$$

where $r^2(\mathbf{x}_i, \mathbf{x}_j) = (\mathbf{x}_i - \mathbf{x}_j)^{\top} \text{COV}(\mathbf{x}_n)^{-1} (\mathbf{x}_i - \mathbf{x}_j)$,
 $w(x) = \exp(-x/2)$, $\beta = 4$.

- **LCOV – COV Hennig (2009)**, a local shape matrix based on the aggregation of covariance matrices computed based on 10% of the nearest neighbors (with regard to the Mahalanobis distance).

PARALLEL ANALYSIS (PA)

As in [Peres-Neto et al. \(2005\)](#) for PCA or in [Caussinus et al. \(2003\)](#).

Computation of cut-offs:

- 10 000 simulations of $\mathcal{N}_n(0, I_p)$
- ICS with $\text{COV}(\mathbf{X}_n)^{-1} \text{COV}_4(\mathbf{X}_n)$
- Quantiles of the ICS eigenvalues at level $1 - \frac{\alpha}{i}$ for each component i , with $\alpha = 5\%$ and $i = 1, \dots, p$.

PARALLEL ANALYSIS (PA)

As in [Peres-Neto et al. \(2005\)](#) for PCA or in [Caussinus et al. \(2003\)](#).

Computation of cut-offs:

- 10 000 simulations of $\mathcal{N}_n(0, I_p)$
- ICS with $\text{COV}(\mathbf{X}_n)^{-1} \text{COV}_4(\mathbf{X}_n)$
- Quantiles of the ICS eigenvalues at level $1 - \frac{\alpha}{i}$ for each component i , with $\alpha = 5\%$ and $i = 1, \dots, p$.

Test:

- Sequentially testing if the ICS eigenvalues are higher than corresponding quantiles.
- Stop as soon as one is lower than the cut-off.

NORMALITY TESTS

Finding the first i^{th} coordinate with no longer information about the structure of the data through normality tests.

NORMALITY TESTS

Finding the first i^{th} coordinate with no longer information about the structure of the data through normality tests.

Normality tests:

- The **D'Agostino** test of skewness (**DA**),
- The **Anscombe-Glynn** (**AG**) test of kurtosis,
- The **Bonett-Seier** (**BS**) test of Geary's kurtosis,
- The **Jarque-Bera** (**JB**) test for normality which is based on both skewness and kurtosis measures,
- The **Shapiro-Wilk** (**SW**) normality test.

NORMALITY TESTS

Finding the first i^{th} coordinate with no longer information about the structure of the data through normality tests.

Normality tests:

- The **D'Agostino** test of skewness (**DA**),
- The **Anscombe-Glynn** (**AG**) test of kurtosis,
- The **Bonett-Seier** (**BS**) test of Geary's kurtosis,
- The **Jarque-Bera** (**JB**) test for normality which is based on both skewness and kurtosis measures,
- The **Shapiro-Wilk** (**SW**) normality test.

Stopping rule

- Sequentially testing if the ICs are gaussian or not.
- Stop as soon as one is gaussian based on the corrected level of 5%.