

# Euler Totient Function

Shivansh Subramanian

22<sup>nd</sup> May 2020

## 1 Introduction

$\phi(x)$  is defined as the number of integers  $d$  less than or equal to  $x$  such that  $\gcd(d, x) = 1$

## 2 Theorems

### 2.1

$$\phi(p) = p - 1$$

### 2.2

$$\phi(p^k) = p^k - p^{k-1} = p^k \times [1 - \frac{1}{p}]$$

Proof

We know that for  $\gcd(n, p^k) = 1$ ,  $n$  must not be divisible by  $p$

$1p, 2p, \dots, (p^{k-1})p$  are all divisible by  $p$

Therefore, there are  $p^{k-1}$  numbers divisible by  $p$ , hence

$$\phi(p^k) = p^k - p^{k-1} = p^k \times [1 - \frac{1}{p}]$$

### 2.3 Multiplicativeness

If  $\gcd(m, n) = 1$ , then

$$\phi(mn) = \phi(m)\phi(n)$$

Proof:

All numbers between 1 and  $mn$  can be written as

$$\begin{array}{cccccc} 1 & 2 & \dots & r & \dots & m \\ m+1 & m+2 & \dots & m+r & \dots & 2m \\ \vdots & \vdots & \dots & \vdots & \dots & \vdots \\ (n-1)m+1 & (n-1)m+2 & \dots & (n-1)m+r & \dots & nm \end{array}$$

We know  $\phi(mn)$  is equal to number of entries in this array such that the element is relatively prime to  $mn$

$$\gcd(km + r, m) = \gcd(r, m)$$

This implies that the numbers in a given column are relatively prime to  $m$  iff  $r$  is relatively prime to  $m$ , therefore  $\phi(m)$  such numbers exist in each row. Now in such a row where  $\gcd(r, m) = 1$ , we have  $n$  elements

$$r, m + r, 2m + r, \dots, (n-1)m + r$$

Here, we have to show that there exist no two numbers such that they give the same remainder on being divided by  $n$ .

Assume

$$km + r \equiv (lm + r) \pmod{n}$$

$$km \equiv (lm) \pmod{n}$$

Since  $m$  and  $n$  are relatively prime

$$k \equiv l \pmod{n}$$

Therefore it is not possible for different  $k$  and  $l$  to give same remainder. This implies that all the elements in row map to  $1, 2, \dots, (n-1)$  as modulo  $n$ . Say element  $s$  maps to  $t$  modulo  $n$ .

$$s \equiv t \pmod{n}$$

We can easily prove that  $\gcd(s, n) = 1$  iff  $\gcd(t, n) = 1$ . Which implies  $\phi(n)$  such integers exist in each row. Hence we conclude that

$$\phi(mn) = \phi(m)\phi(n)$$

## 2.4

If

$$n = p_1^{k_1} p_2^{k_2} \dots p_r^{k_r}$$

Then

$$\phi(n) = n \prod_{x=1}^r \left[1 - \frac{1}{p_x}\right]$$

Proof:

Since we know  $\phi(n)$  is multiplicative

$$\phi(n) = \prod_{x=1}^r \phi(p_x^{k_x})$$

From Theorem 2.2

$$\phi(n) = \prod_{x=1}^r p_x^{k_x} \times [1 - \frac{1}{p_x}]$$

$$\phi(n) = n \prod_{x=1}^r [1 - \frac{1}{p_x}]$$

## 2.5

For any positive integer n

$$\sqrt{\frac{n}{2}} \leq \phi(n) \leq n$$

Proof:

The part of proving  $\phi(n) \leq n$  is trivial, we look at the other part then

Let

$$n = p_1 p_2 \dots p_k q_1^{a_1} q_2^{a_2} \dots q_l^{a_l}$$

Let

$$s = p_1 p_2 \dots p_k$$

$$t = q_1^{a_1} q_2^{a_2} \dots q_l^{a_l}$$

We know

$$\phi(n) = \phi(st) = \phi(s)\phi(t)$$

$$\frac{\phi(n)}{\sqrt{n}} = \frac{\phi(s)}{\sqrt{s}} \frac{\phi(t)}{\sqrt{t}}$$

Then

$$\frac{\phi(s)}{\sqrt{s}} = \prod_{x=1}^k \frac{p_x - 1}{\sqrt{p_x}}$$

For all  $p \geq 2$ , we have  $\frac{p-1}{\sqrt{p}} > 1$

Therefore there are two cases, where m contains only 2 and where n contains 2 and some other factors

The former cases is trivial, and since in the latter case the fraction for other factors other than 2 is greater than 1, we can easily prove

$$\frac{\phi(s)}{\sqrt{s}} \geq \frac{1}{\sqrt{2}}$$

$$\frac{\phi(t)}{\sqrt{t}} = \prod_{x=1}^l q_x^{a_x-1} (q_x - 1) \geq 1$$

Therefore we arrive at the conclusion

$$\sqrt{\frac{n}{2}} \leq \phi(n) \leq n$$

## 2.6

If

$$n = p_1^{a_1} p_2^{a_2} \dots p_r^{a_r}$$

Then

$$\phi(n) \geq \frac{n}{2^r}$$

Proof:

Since we know  $\phi$  is multiplicative

$$\phi(n) = \phi(p_1^{a_1}) \phi(p_2^{a_2}) \dots \phi(p_r^{a_r})$$

Since for any p

$$\phi(p^a) = p^a \left[1 - \frac{1}{p}\right]$$

We know that

$$\frac{1}{2} \geq \frac{1}{p}$$

Therefore

$$\phi(p^a) = p^a \left[1 - \frac{1}{p}\right] \geq \frac{p^a}{2}$$

$$\phi(n) \geq \frac{n}{2^r}$$

## 2.7

If n is a composite number,

$$\phi(n) \leq n - \sqrt{n}$$

Proof:

Let p be the smallest prime divisor of n, then

Let

$$n = p_1^{a_1} p_2^{a_2} \dots p_r^{a_r}$$

Where  $p = p_1$  Then

$$\phi(n) = n \prod_{k=1}^r \left[1 - \frac{1}{p_k}\right]$$

Assume such a n' that  $p \nmid n$ . Then

$$\phi'(n') = n' \prod_{k=2}^r \left[1 - \frac{1}{p_k}\right]$$

And

$$\phi'(n') \leq n'$$

$$\phi'(n') \times p^{a_1} \leq n' \times p^{a_1}$$

$$\phi'(n') \times p^{a_1} \leq n$$

$$\phi'(n') \times p^{a_1} \times [1 - \frac{1}{p}] \leq n \times [1 - \frac{1}{p}]$$

$$\phi(n) \leq n \times [1 - \frac{1}{p}]$$

And since

$$p \leq \sqrt{n}$$

$$\phi(n) \leq n \times [1 - \frac{1}{p}] \leq n \times [1 - \frac{1}{\sqrt{n}}]$$

$$\phi(n) \leq n - \sqrt{n}$$