

Table 1: Exhibit check

	Weak ($r = 1.2$)	Strong ($r = 3$)
Proceeds without a challenger, t_0	0.291667	0.416667
Proceeds with a high-value challenger, t_H	0.704167	1.541667
Proceeds with a low-value challenger, t_L	0.687500	0.875000
Gross profit of a high-value challenger, g_H	9.295833	8.458333
Gross profit of a low-value challenger, g_L	0.312500	0.125000
Information spread, $\Delta_T = t_H - t_L$	0.016667	0.666667
Expected gross profit at the prior, $B_r(1/2)$	4.804167	4.291667

Benchmark primitives $h = 10$, $\ell = 1$, $p = 0.5$; values are incremental value per target share. Δ_T is the spread in expected target proceeds across challenger types, not total merger value and not the trader's expected profit. Closed forms in the text agree with direct integration of the realized sale rule to within 10^{-9} (Online Appendix C.1).

Table 2: Exhibit check

	q_H	q_L	E	O_H	$\mathcal{R}_T$	Evidence	Unique
<i>Panel A. Equilibrium outcomes</i>							
Weak incumbent ($r = 1.2$)	0	0	0.250000	0.125000	0.392708	analytical	yes
Strong incumbent ($r = 3$)	1	-1	0.522757	0.324192	0.872392	analytical	yes
Very strong incumbent ($r = 3.6$)	1	-1	0.250000	0.125000	0.664236	analytical	yes
<i>Panel B. Information controls</i>							
Frozen informative orders, $r = 1.2$	1	-1	0.562178	0.350606	0.520039	numerical diagnostic	n/a
Frozen informative orders, $r = 3$	1	-1	0.522757	0.324192	0.872392	numerical diagnostic	n/a
Price hidden, $r = 1.2$	0	0	0.250000	0.125000	0.392708	analytical	yes
Price hidden, $r = 3$	1	-1	0.250000	0.125000	0.614583	analytical	yes
<i>Panel C. Access to prices at $r = 3$</i>							
				Price observed		Price hidden	Gain
Target proceeds $\mathcal{R}_T$				0.872392		0.614583	0.257809
Net acquisition surplus $\mathcal{W}$				2.382301		2.302083	0.080218

Benchmark primitives, Laplace noise, cost atoms. E is the preparation probability, O_H the probability that a high-value challenger acquires the target, $\mathcal{R}_T$ expected target proceeds. Evidence is the result status of the row (analytical, computer-assisted, or numerical diagnostic); Unique records separately whether the continuation is the unique one under the stated bound. Panel A rows are analytical outcomes of Proposition 2.

Panel B rows are information controls: control is an experiment role, not an evidence class. The frozen rows are a fixed-profile control that imposes full orders at both strengths and lets the challenger reoptimize; the profile is not an investor equilibrium at $r = 1.2$ and coincides with the equilibrium profile at $r = 3$ while its role remains that of a control, so its Unique field is not applicable. The price-hidden rows are the reoptimized equilibria of the economy in which the price is withheld from the challenger. The matched-dividend invariance diagnostic of Section 6.1 is reported in the online matched-price panel.

Panel C: $\mathcal{W}$ is allocation value net of paid preparation costs, not target revenue. The five strict margins of Proposition 2 at $(r_0, r_1) = (1.2, 3)$ are $\zeta_L = 1.37 \times 10^0$, $\zeta_{H0} = 1.20 \times 10^0$, $\zeta_{H1} = 2.17 \times 10^{-1}$, $\zeta_0 = 3.33 \times 10^{-3}$, $\zeta_1 = 2.41 \times 10^{-3}$.

Table 3: Exhibit check

	E weak	E strong	ΔE (pp)	O_H weak	O_H strong	Evidence	Unique
<i>Panel A. Noise and preparation costs</i>							
Laplace, cost atoms	0.250000	0.522757	27.28	0.125000	0.324192	analytical	yes
Laplace, cost mixture	0.250000	0.522715	27.27	0.125000	0.324148	analytical	yes
logistic, cost atoms	0.250000	0.301509	5.15	0.125000	0.161994	analytical	yes
logistic, cost mixture	0.250000	0.301374	5.14	0.125000	0.161854	analytical	yes
<i>Panel B. Moderate values ($h = 2, \ell = 1$)</i>							
Moderate values	0.250000	0.526805	27.68	0.125000	0.327032	analytical	yes
<i>Panel C. Complementary private information</i>							
$a = 0.70, d = 0.75$	0.850000	0.879438	2.94	0.425000	0.448942	analytical	yes

E is the preparation probability and O_H the probability of high-value challenger ownership, both probabilities. $\Delta E = 100(E_{\text{strong}} - E_{\text{weak}})$ is the change in preparation in percentage points, computed from the same validated probabilities as the two preparation columns, not a percentage change.

Each row is a distinct parameter vector (Appendix A.8): Panel A uses the benchmark vector with strengths (1.2, 3); Panel B uses the moderate-value vector with (1.05, 1.5); Panel C uses the complementary-signal vector with (1.1, 2.3), investor accuracy a and buyer accuracy d . Panels B and C are separate illustrations, not one joint calibration. The cost mixture uses half-width $\varepsilon_C = 0.1$; noise laws share the scale b , not the variance.

Evidence is the result status of the comparison; Unique records whether the sufficient uniqueness conditions of the relevant proposition hold (all five margins strictly positive). The individual margins and their minimum are in the online table of sufficient-condition margins; the complete accuracy grid is the online accuracy-grid table.

Table 4: Complementary private information: complete accuracy grid.

Investor a	Buyer d	E weak	E strong	ΔE (pp)	O_H weak	O_H strong	Minimum margin	Evidence	Comparison condition
0.68	0.73	0.850000	0.850000	0.00	0.425000	0.425000	-1.49×10^{-1}	numerical diagnostic	not met
0.68	0.74	0.850000	0.850000	0.00	0.425000	0.425000	-7.71×10^{-2}	numerical diagnostic	not met
0.68	0.75	0.850000	0.850000	0.00	0.425000	0.425000	-5.30×10^{-3}	numerical diagnostic	not met
0.68	0.76	0.925000	0.880132	-4.49	0.482000	0.449575	-3.75×10^{-2}	numerical diagnostic	not met
0.68	0.77	0.925000	0.882492	-4.25	0.482750	0.451741	-1.27×10^{-1}	numerical diagnostic	not met
0.69	0.73	0.850000	0.850000	0.00	0.425000	0.425000	-1.22×10^{-1}	numerical diagnostic	not met
0.69	0.74	0.850000	0.850000	0.00	0.425000	0.425000	-5.08×10^{-2}	numerical diagnostic	not met
0.69	0.75	0.850000	0.878570	2.86	0.425000	0.448157	9.57×10^{-4}	analytical	met
0.69	0.76	0.925000	0.880909	-4.41	0.482000	0.450290	-3.75×10^{-2}	numerical diagnostic	not met
0.69	0.77	0.925000	0.883109	-4.19	0.482750	0.452331	-1.27×10^{-1}	numerical diagnostic	not met
0.70	0.73	0.850000	0.850000	0.00	0.425000	0.425000	-9.54×10^{-2}	numerical diagnostic	not met
0.70	0.74	0.850000	0.850000	0.00	0.425000	0.425000	-2.48×10^{-2}	numerical diagnostic	not met
0.70*	0.75	0.850000	0.879438	2.94	0.425000	0.448942	1.45×10^{-3}	analytical	met
0.70	0.76	0.925000	0.881610	-4.34	0.482000	0.450945	-3.75×10^{-2}	numerical diagnostic	not met
0.70	0.77	0.925000	0.883674	-4.13	0.482750	0.452879	-1.27×10^{-1}	numerical diagnostic	not met
0.71	0.73	0.850000	0.850000	0.00	0.425000	0.425000	-6.88×10^{-2}	numerical diagnostic	not met
0.71	0.74	0.850000	0.878066	2.81	0.425000	0.447687	9.59×10^{-4}	analytical	met
0.71	0.75	0.850000	0.880215	3.02	0.425000	0.449658	7.77×10^{-4}	analytical	met
0.71	0.76	0.925000	0.882247	-4.28	0.482000	0.451551	-3.75×10^{-2}	numerical diagnostic	not met
0.71	0.77	0.925000	0.884195	-4.08	0.482750	0.453392	-1.27×10^{-1}	numerical diagnostic	not met
0.72	0.73	0.850000	0.850000	0.00	0.425000	0.425000	-4.24×10^{-2}	numerical diagnostic	not met
0.72	0.74	0.850000	0.878915	2.89	0.425000	0.448457	6.16×10^{-4}	analytical	met
0.72	0.75	0.850000	0.880919	3.09	0.425000	0.450315	1.00×10^{-4}	analytical	met
0.72	0.76	0.925000	0.882833	-4.22	0.482000	0.452115	-3.75×10^{-2}	numerical diagnostic	not met
0.72	0.77	0.925000	0.884680	-4.03	0.482750	0.453877	-1.27×10^{-1}	numerical diagnostic	not met

Notes. All 25 accuracy combinations in the declaration in Section C.3 are reported; none is dropped. Other parameters are $h = 10$, $\ell = 1$, $p = 0.5$, $\rho = 0.85$, $c_L = 1$, $c_H = 7.14$, $b = 2$, $k = 0.015$, with strengths $r_0 = 1.1$ and $r_1 = 2.3$. E and O_H are probabilities; ΔE is the change in preparation in percentage points from the same validated probabilities. The minimum margin is the smallest of the five conditions in (OA.29); the comparison condition is met when it is strictly positive (6 rows), which places the row inside the analytical uniqueness region. A condition that is not met is not a rejection: every row is a validated equilibrium of the finite check, and its evidence is then numerical diagnostic unless a separate analytical or interval argument supports more. The asterisk marks the example used in the main text. All rows pass the declared numerical acceptance checks.