

MATH 135 - Algebra for Honours Mathematics

Winter 2014

Contents

1	Injections and Bijections	2
1.1	Injective(One-to-One)	2
1.1.1	Definition	2
1.1.2	Simple Example	2
1.1.3	Hard Example	2
1.1.4	Frequently Asked Questions	2
1.2	Surjective	3
1.2.1	Definition	3
1.3	Bijections	3
1.3.1	Definition	3
1.3.2	Simple Example	3
1.4	Summary	3
2	Counting	4
2.1	Bijection and Cardinality	4
2.1.1	Definition	4
2.1.2	Guidelines	4
2.2	Finite Sets	4
2.2.1	Definitions	4
2.2.2	Propositions	4
2.2.3	Example	5
2.3	Infinite Sets	5
2.3.1	Propositions	5
2.3.2	Example	6

Chapter 1

Injective, Surjective and Bijections

1.1 Injective(One-to-One)

1.1.1 Definition

Injective: Let S and T be two sets. A function $f : S \rightarrow T$ is **one-to-one**(or **injective**) iff for every $x_1 \in S$, $f(x_1) = f(x_2)$ implies that $x_1 = x_2$ and $|S| \leq |T|$. When trying to prove that a function is one-to-one, start off with $f(x_1) = f(x_2)$ and try to use algebraic manipulation to obtain $x_1 = x_2$.

1.1.2 Simple Example

Proposition: Let $m \neq 0$ and b be fixed real numbers. The function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = mx + b$ is one to one

Proof: Let $x_1, x_2 \in S$. Suppose that $f(x_1) = f(x_2)$. Now we show that $x_1 = x_2$. Since $f(x_1) = f(x_2)$, $mx_1 + b = mx_2 + b$. Subtracting b from both sides and dividing by m gives $x_1 = x_2$ as required.

1.1.3 Hard Example

Proposition: Let $f : T \rightarrow U$ and $g : S \rightarrow T$ be one-to-one functions. Then $f \circ g$ is a one-to-one function.

Proof: Let $x_1, x_2 \in S$. Suppose that $(f \circ g)(x_1) = (f \circ g)(x_2)$. Since $(f \circ g)(x_1) = (f \circ g)(x_2)$, we know that $f(g(x_1)) = f(g(x_2))$. Since f is one-to-one, we know that $g(x_1) = g(x_2)$. And since g is one-to-one, $x_1 = x_2$ as required.

1.2 Surjective

1.2.1 Definition

Surjective: A function $f : S \rightarrow T$ is **surjective**(or **onto**) if and only if for every $y \in T$ there exists an $x \in S$ so that $f(x) = y$. This implies that $|S| \geq |T|$.

When trying to prove that a function is onto, try to find a function $g(x)$ such that $f(g(x)) = y$ to prove that each y in the codomain is mapped to.

1.3 Bijections

1.3.1 Definition

Bijection: A function $f : S \rightarrow T$ is **bijection** iff f is both surjective and injective.

1.3.2 Simple Example

We have already shown that for $m \neq 0$ and b a fixed real number, the function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = mx + b$ is both surjective and injective. Hence, f is bijective.

1.4 Summary

- $f : S \rightarrow T$ is a function iff $\forall s \in S \exists! t \in T, f(s) = t$ where $!$ means unique
- $f : S \rightarrow T$ is surjective iff $\forall t \in T \exists s \in S, f(s) = t$, meaning for each element $t \in T$, there is at least one element $s \in S$ so that $f(s) = t$
- $f : S \rightarrow T$ is injective iff $\forall x_1 \in S \forall x_2 \in S, f(x_1) = f(x_2) \Rightarrow x_1 = x_2$ or $x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$, meaning for each element $t \in T$, there is at most one element $s \in S$ so that $f(s) = t$

1.4.1 Frequently Asked Questions

Questions to be added

Chapter 2

Counting

2.1 Bijection and Cardinality

2.1.1 Definition

Cardinality: If there exists a bijection between the sets S and T , we say that the sets have the same and we write $|S| = |T|$.

Number of Elements, Finite, Infinite: If there exists a bijection between a set S and \mathbb{N}_n , we say that the **number of elements** in S is n and we write $|S| = n$. Moreover, we also say that S is a **finite set**. If no bijection exists between a set S and \mathbb{N}_n for any n , we say that S is an **infinite set**.

Countable: A set S is **countable** if there exists an injective function f from S to the natural numbers \mathbb{N} .

2.1.2 Guidelines

Proposition: Let $S = \dots$ Let $T = \dots$ Then there exists a bijection $f : S \rightarrow T$. Hence, $|S| = |T|$.

To do this, we must prove that f is both surjective and injective.

Consider the function $f : S \rightarrow T$ defined by $f(s) = \dots$. We show that f is surjective. Let $t \in T$. Consider $s = \dots$. We show that $s \in S$. Now we show that $f(s) = t$.

We then show that f is injective. Let $s_1, s_2 \in S$ and suppose that $f(s_1) = f(s_2)$. Now we show that $s_1 = s_2$.

Hence, $f : S \rightarrow T$ is a bijection and $|S| = |T|$.

2.2 Finite Sets

2.2.1 Definitions

Disjoint: Set S and T are **disjoint** if $S \cap T = \emptyset$

2.2.2 Propositions

Cardinality of Intersecting Sets(CIS): If S and T are any finite sets, then

$$|S \cup T| = |S| + |T| - |S \cap T|$$

Cardinality of Disjoint Sets(CDS): If S and T are disjoint finite sets, then

$$|S \cup T| = |S| + |T|$$

2.2.3 Example

Proof of CDS:

1. Since S is a finite set, there exists a bijection $f : S \rightarrow \mathbb{N}_m$ for some non negative integer m , and $|S| = m$
2. Since T is a finite set, there exists a bijection $g : T \rightarrow \mathbb{N}_n$ for some non negative integer n , and $|T| = n$
3. Construct function $h : S \cup T \rightarrow \mathbb{N}_{m+n}$ as follows:
 $h(x) = f(x)$ if $x \in S$ else $g(x) + m$ if $x \in T$
4. To show that h is surjective, let $y \in \mathbb{N}_{m+n}$. If $y \leq m$, then because f is surjective there exists an element $x \in S$ so that $f(x) = y$, hence $h(x) = y$. If $m + 1 \leq y \leq m + n$, then because g is surjective, there exists an element $x \in T$ so that $g(x) = y - m$ and so $h(x) = (y - m) + m = y$.
5. To show that h is injective, let $x_1, x_2 \in S \cup T$ and suppose that $h(x_1) = h(x_2)$. If $h(x) \leq m$ then $h(x) = f(x)$ so if $h(x_1) \leq m$ we have
 $h(x_1) = h(x_2) \Rightarrow f(x_1) = f(x_2)$
But since f is a bijection $f(x_1) = f(x_2)$ implies $x_1 = x_2$ as needed. If $h(x) > m$ then $h(x) = g(x)$ so if $h(x_1) > m$ we have
 $h(x_1) = h(x_2) \Rightarrow g(x_1) + m = g(x_2) + m \Rightarrow g(x_1) = g(x_2)$
But since g is a bijection $g(x_1) = g(x_2)$ implies $x_1 = x_2$ as needed. Since h is a function which is both injective and surjective, h is bijective.

6. Thus

$$|S \cup T| = |\mathbb{N}_{m+n}| = m + n = |\mathbb{N}_m| + |\mathbb{N}_n| = |S| + |T|$$

If it wasn't clear, $f(x)$ is mapped to $1, 2, \dots, m$ and $g(x) + m$ is mapped to $m + 1, m + 2, \dots, m + n$.

2.3 Infinite Sets

2.3.1 Propositions

Cardinality of Subsets of Finite Sets(CSFS): If S and T are finite sets, and $S \subset T$, then $|S| < |T|$

$|\mathbb{N}| = |2\mathbb{N}|$: Let $2\mathbb{N}$ be the set of positive even natural numbers. Then $|\mathbb{N}| = |2\mathbb{N}|$

$$|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$$

Even-Odd Factorization of Natural Numbers(EOFNN): Any natural number n can be written uniquely as $n = 2^i q$ where i is a non-negative integer and q is an odd natural number. Note: use EOFNN to prove $|\mathbb{N} \times \mathbb{N}| = |\mathbb{N}|$.
Note: Not all infinite sets have the same size

2.3.2 Example

Proof of $|\mathbb{N}| = |2\mathbb{N}|$

We want to prove that there's a bijection between both sets

1. Consider the function $f : \mathbb{N} \rightarrow 2\mathbb{N}$ defined by $f(s) = 2s$
2. We show that f is surjective. Let $t \in 2\mathbb{N}$. Consider $s = \frac{1}{2}t$. We show that $s \in \mathbb{N}$ since $f(\frac{1}{2}t) = t$ and therefore is surjective
3. We show that f is injective. Let $s_1, s_2 \in \mathbb{N}$ and suppose that $f(s_1) = f(s_2)$. Now we show that $s_1 = s_2$. Since $f(s_1) = 2s_1$ and $f(s_2) = 2s_2$, $s_1 = s_2$.
4. Hence, $f : \mathbb{N} \rightarrow 2\mathbb{N}$ is a bijection and $|\mathbb{N}| = |2\mathbb{N}|$.