

Quiz 2 Practice

The following are some practice problems for the second quiz. You can use computer program to help with calculation but no access to the internet.

Rule: Quiz 2 is on Monday November 18 from 4-5pm (the regular lecture starts from 5.10-5.50pm). The cheat sheet of one A4 double-sided is allowed (just like in quiz 1).

Format: 10% of total grade.

One programming problem 5%, Two non-programming problems (like in quiz 1) 5%

Sample programming problems

- (Simulation) Define a bias coin to have a probability of getting head =0.75 and probability of getting tail =0.25. Then let a random variable X to be the number of heads after tossing this bias coin twice.
 - Generate 10,000 random values of X . Then, use the data to find the mean (expected value), standard deviation. Also plot the distribution of X .
 - Let the sample mean $\bar{X} = \frac{X_1 + X_2 + \dots + X_{25}}{25}$. Generate 10,000 random values of \bar{X} . Then, use the data to find the mean (expected value), standard deviation. Also plot the distribution of \bar{X} .
- (Benford's law) Investigate the distribution of the leading digits from the 200-by-200 multiplication table. Compare the distribution with the Benford's law.

For example, 4-by-4 multiplication table:

m/n	1	2	3	4
1	1	2	3	4
2	2	4	6	8
3	3	6	9	12
4	4	8	12	16

The list of leading digits is [1, 2, 3, 4, 2, 4, 6, 8, 3, 6, 9, 1, 4, 8, 1, 1].

- (Alarm clock) Recall that you can generate $X = \text{Exp}(\lambda)$, the exponential random variable of mean λ by

$$x = -\frac{1}{\lambda} \log(1 - U),$$

where U is a uniform $(0, 1)$.

Let $X_1 \sim \text{Exp}(1)$, $X_2 \sim \text{Exp}(2)$, $X_3 \sim \text{Exp}(3)$.

- a) Simulate 10,000 values of X_1 . Plot the distribution and compare it with the probability density function of $Exp(1)$ i.e. $f(x) = e^{-x}, x \geq 0$.
- b) Use the simulation to approximate the probability $\mathbb{P}(X_1 = \min\{X_1, X_2, X_3\})$.
- c) Use the simulation to approximate the probability $\mathbb{P}(X_1 = \max\{X_1, X_2, X_3\})$.

Sample non-programming problems

1. (Exponential distribution) Show the memoryless property of exponential function:
Given $X \sim Exp(\lambda)$,

$$\mathbb{P}\{X \geq a \mid X \geq b\} = \mathbb{P}\{X \geq a - b\}.$$

2. Let X_1 and X_2 be independent and identically distributed $Unif\{3, 6, 9\}$. Define the random variable

$$Y := \begin{cases} 5 & \text{if } X_2 \leq 7, \\ 8 & \text{if } X_2 > 7. \end{cases}$$

Compute the following.

- (a) $Cov(X_1, X_2)$ and $corr(X_1, X_2)$.
 - (b) $Cov(X_1, Y)$ and $corr(X_1, Y)$.
 - (c) $Cov(X_2, Y)$ and $corr(X_2, Y)$.
3. (Descent) We define the indicator variable of $w \in S_n$ at position i , where $i = 1, 2, \dots, n-1$ as

$$I_i = \begin{cases} 1, & \text{if } w_i > w_{i+1} \\ 0, & \text{otherwise.} \end{cases}$$

$$\text{Show that } Cov(I_i, I_j) = \begin{cases} 0, & \text{if } |i - j| \geq 2 \\ \frac{1}{4}, & \text{if } i = j \\ -\frac{1}{12}, & \text{if } |i - j| = 1. \end{cases}$$

4. (Toilet problem) Calculate the probability of getting the best toilet when there are 4 toilets and the learning phase is of length 1. Show your work.