Discrete Simulation: Homework 1

Austin J. Maddison

Mahidol University International College

September 20, 2024

1. Birthday Problem

The simulations ran were set to $seed = 27$ and $N = 100000$.

a)

$$p_{10} = 0.1169$$
$$CI = [0.1169, 0.1189]$$

b)

$$p_{20} = 0.4113$$
$$CI = [0.4113, 0.4143]$$

c)

$$p_{30} = 0.7059$$
$$CI = [0.7059, 0.7088]$$

d) Found n that satisfies the condition $p_n \leq 0.5$ from extracting the smallest difference from varying n's of p_n and the function of $p = 0.5$.



Figure 1: Closest point to $p = 0.5$ was $p_{23} = 0.54$ which satisfies the condition.

2. Alice and Bob Play a Game

The strategy I let Alice have is the following...

Case 1: Found Exclusive Output
She presses the button recording n output values $x_1, x_2, ...x_n$. As she records each output she checks whether x_i is exclusive to one of the buttons output range. If x_i is exclusive she returns the corresponding button as the answer.

Case 2: No Exclusive Output
After she presses the button n times with no exclusive output appearing. She calculates $\bar{x}$ and finds the minimum difference between the mean output range of the 2 buttons. She returns the corresponding button that gets the minimum distance.

Psuedocode
Although the source code differs because loops are removed for speed, the idea is the same.

```
xs = []
for i in range(0, n):
    x_n = button_unknown.get_next_value()
    xs.append(x_n)

    # Case 1: Found exclusive output.
    if(x_n == 1)
        return 1  # it is button 1
    if(x_n == 100)
        return 2  # it is button 2

# Case 2: No exclusive output. Evaluate
    the minimum distance of means.
x_mean = sum(xs) / n
button_1_mean = (1 + 99)/2
button_2_mean = (2 + 100)/2

return argmin([abs(x_mean - button_1_mean)
    , abs(x_mean - button_2_mean)]) + 1
```

Find n such that Alice is correct ≤ 0.9

Similar to problem 1.Birthday Problem, I sampled Alice's strategy over some reasonable range of n

$[0, 500]$ and extract the n that results in probability p that Alice is correct is atleast 0.99. I set the seed=27 and ran 1000 trials for each all n's.



Figure 2: Closest point to $p = 0.99$ was $p_{375} = 0.990$ which satisfies the condition.



Figure 3: y-lim[0.8, 1.0] of Figure 2.

3. Practice with Uniform and Geometric Distributions

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

4. Source Code

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.