

Contents

1	Countability	2
----------	---------------------	----------

Countability

Definition 1.0.1 ► Second-Countable Space

topological space X is called second-countable if it has a countable basis for its topology.

Definition 1.0.2 ► Lindelöf Space

A topological space X is called a Lindelöf space if every open cover of X has a countable subcover.

Theorem 1.0.3 ► Lindeöf Theorem

Every second-countable space is Lindeöf.

Proof. Let X be a second-countable space, $\mathcal{B} = \{B_n\}_{n=1}^{\infty}$ is a countable basis of X . Let $\mathcal{U} = \{U_{\alpha}\}_{\alpha \in A}$ be any open cover of X . Let

$$\mathcal{B}' := \{B_n \in \mathcal{B} : \exists U_{\alpha} \in \mathcal{U}, \text{ s.t. } B_n \subseteq U_{\alpha}\}$$

For each $x \in X$, $\exists U_{\alpha} \in \mathcal{U}$ such that $x \in U_{\alpha}$. Since $\mathcal{B}$ is a basis, there $\exists B_n \in \mathcal{B}$ such that $x \in B_n \subseteq U_{\alpha}$. $B_n \in \mathcal{B}'$ by definition. Thus $\mathcal{B}'$ covers X . For each $B_{n_k} \in \mathcal{B}'$, we choose an open set U_k in $\mathcal{U}$, such that $B_{n_k} \subseteq U_k$. Then $\{U_k\}_{k=1}^{\infty}$ is a subcover of $\mathcal{U}$. $\square$