

Introduction

1.1 Arithmetic in $[0, \infty]$

1.2 Metric Space

Lemma 1.2.1

X is a metric space, $x \in X$, B_1, B_2 are two balls in X . If $x \in B_1 \cap B_2$, then x is the center of an open ball $B \subseteq B_1 \cap B_2$.

Proof. We may assume that B_1, B_2 have different center. Let $y_1 \neq y_2$,

$$B_1 = \{x : d(x, y_1) < r_1\}, \quad B_2 = \{x : d(x, y_2) < r_2\}$$

Take $x \in B_1 \cap B_2$, we claim that there exists $r_0 > 0$ such that $B := \{y : d(x, y) < r_0\} \subseteq B_1 \cap B_2$. Otherwise, for all $r > 0$, there exists a point y_r such that the following two hold at the same time

1. $d(x, y_r) < r$
2. $d(y_r, y_1) \geq r_1$ or $d(y_r, y_2) \geq r_2$.

Thus

$$d(x, y_1) \geq d(y_r, y_1) - d(x, y_r) > r_1 - r, \quad \text{or} \quad d(x, y_2) > r_2 - r$$

The above shows that

$$r > \min\{r_1 - d(x, y_1), r_2 - d(x, y_2)\}, \quad \forall r > 0$$

which is a contradiction. □

1.3 Limits of Set Sequences

Definition 1.3.1 ▶ Limit Inferior and Limit Superior

Let $\{A_n\}_{n=1}^{\infty}$ be a sequence of sets.

1. By the *Limit Inferior of the sequence*, we mean

$$\liminf_{n \rightarrow \infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=N}^{\infty} A_n = \{x : \exists N_0 \in \mathbb{N}, \forall n > N_0, x \in A_n\}$$

2. By the *Limit Superior of the sequence*, we mean

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} A_n = \{x : \forall N \in \mathbb{N}, \exists n \geq N, x \in A_n\}$$

Note.

1. Limit Inferior 包含那些“最终稳定下来”的元素，即从某个点之后就永远属于序列中所有后续集合的元素。
2. Limit Superior 包含那些“反复出现”的元素，即在无限多个集合中出现的元素。

Proposition 1.3.2

For any sequence of sets $\{A_n\}$, it always holds that:

$$\liminf_{n \rightarrow \infty} A_n \subseteq \limsup_{n \rightarrow \infty} A_n$$

Proof sketch.

直觉上是显然的，因为“最终稳定下来”的元素一定会“反复出现”。

Proof. It is obvious by the $\{x : x \in P\}$ form representation of the Limit Inferior and Superior. □

Definition 1.3.3 ▶ Limit of a Set Sequence

If the limit inferior and limit superior of a set sequence $\{A_n\}_{n=1}^{\infty}$ are equal, i.e., $\liminf_{n \rightarrow \infty} A_n = \limsup_{n \rightarrow \infty} A_n$, then we say the *limit* of the se-

quence exists, and it is defined as:

$$\lim_{n \rightarrow \infty} A_n = \liminf_{n \rightarrow \infty} A_n = \limsup_{n \rightarrow \infty} A_n$$

Proposition 1.3.4

Let $\{A_n\}_{n=1}^{\infty}$ be a sequence of sets.

1. If $\{A_n\}$ is an increasing sequence, then the limit of $\{A_n\}$ exists and

$$\lim_{n \rightarrow \infty} A_n = \bigcup_{n=1}^{\infty} A_n$$

2. If $\{A_n\}$ is a decreasing sequence, then the limit of $\{A_n\}$ exists and

$$\lim_{n \rightarrow \infty} A_n = \bigcap_{n=1}^{\infty} A_n$$

Proof. 1. If $\{A_n\}$ is increasing, then

$$\bigcap_{n=N}^{\infty} A_n = A_N$$

and

$$\bigcup_{n=N}^{\infty} A_n = \bigcup_{n=1}^{\infty} A_n, \forall N \in \mathbb{N}.$$

Thus

$$\liminf_{n \rightarrow \infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=N}^{\infty} A_n = \bigcup_{N=1}^{\infty} A_N$$

and

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=1}^{\infty} A_n = \bigcup_{n=1}^{\infty} A_n$$

which are the same.

2. If $\{A_n\}$ is decreasing, then

$$\bigcap_{n=N}^{\infty} A_n = \bigcap_{n=1}^{\infty} A_n$$

and

$$\bigcup_{n=N}^{\infty} A_n = A_N$$

Thus

$$\liminf_{n \rightarrow \infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=N}^{\infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=1}^{\infty} A_n = \bigcap_{n=1}^{\infty} A_n$$

and

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} A_n = \bigcap_{N=1}^{\infty} A_N$$

which are the same.

□

Proposition 1.3.5

For a sequence of sets $\{A_n\}_{n=1}^{\infty}$ and their corresponding sequence of indicator functions $\{\chi_{A_n}(x)\}_{n=1}^{\infty}$:

- $\chi_{\liminf_{n \rightarrow \infty} A_n}(x) = \liminf_{n \rightarrow \infty} \chi_{A_n}(x)$
- $\chi_{\limsup_{n \rightarrow \infty} A_n}(x) = \limsup_{n \rightarrow \infty} \chi_{A_n}(x)$
- $\lim_{n \rightarrow \infty} A_n$ exists, then $\chi_{\lim_{n \rightarrow \infty} A_n}(x) = \lim_{n \rightarrow \infty} \chi_{A_n}(x)$

Proof sketch.

- 若 x 在 limit inferior 里面, 则 x 是“最终稳定”的, $\chi_{A_n}(x)$ 是关于 n “最终”恒为 1 的.
- 若 x 在 limit superior 里面, 则 x 是“反复出现”的, 即相当于 N 多大, 总会出现之后的某个 n 使得 $\chi_{A_n}(x) = 1$.
- 当极限存在时, 函数列 $\{\chi_{A_n}(x)\}_{n=1}^{\infty}$ 的 limit inferior 和 supperior 根据上两条相等, 等于极限集合的 χ .

1.4 Exercises

Problem 1.4.1 ► Convex Bodies

A convex body is a compact, convex subset $K \subseteq \mathbb{R}^n$ which contains 0 as an interior point and which is symmetric around the origin .

1. Prove that the following formula defines a norm on $\mathbb{R}^n$:

$$N(x) := \inf \left\{ \lambda > 0 : \frac{x}{\lambda} \in K \right\}$$

2. Prove that K is homeomorphic to $\mathbb{D}^n$ and that ∂K is homeomorphic to $\mathbb{S}^{n-1}$.

Note.

1. 若 K 是一个单位球. 则 $N(x) = \|x\|$.
2. 证明 $N(x)$ 是一个 norm, 需要证明: 1. 非负正定 2. 绝对一次齐次 3. 三角不等式.
3. 由于 φ 的形式足够简单, 可以通过直接构造逆映射证明双射.

Lemma 1.4.2

X is compact, Y is Hausdorff . If $\exists f : X \rightarrow Y$ is a continuous bijection, then X, Y are homeomorphic.

Proof. 1. It is obvious that $N(x) \geq 0$. If $N(x) = 0$, then

$$\inf \left\{ \lambda > 0 : \frac{x}{\lambda} \in K \right\} = 0$$

which means that for all $\varepsilon > 0$, $\frac{x}{\varepsilon} \in K$. Since K is compact, $\text{dist}(K, 0) < \infty$. If $x \neq 0$, then $\|x\| > 0$. We take $\varepsilon_0 < \frac{1}{\text{dist}(K, 0) \cdot \|x\|}$. Then $\left\| \frac{x}{\varepsilon_0} \right\| > \text{dist}(K, 0)$. But $\frac{x}{\varepsilon_0} \in K$, which is a contradiction. For the second part, let $x \in \mathbb{R}^n, k \in \mathbb{R} \setminus \{0\}$. For all $\lambda > N(x)$,

$$\frac{x}{\lambda} \in K \implies \frac{kx}{k\lambda} \in K$$

, which shows that $N(kx) \leq kN(x)$. Replace k by $\frac{1}{k}$ and replace x by kx , we have $N(kx) \geq kN(x)$. Thus $N(kx) = kN(x)$. For the triangle

inequality, we need to show that

$$\inf\left\{\lambda > 0 : \frac{x_1 + x_2}{\lambda} \in K\right\} \leq \inf\left\{\lambda > 0 : \frac{x_1}{\lambda} \in K\right\} + \inf\left\{\lambda > 0 : \frac{x_2}{\lambda} \in K\right\}$$

Let $\lambda_0 = N(x_1 + x_2)$. Only need to show that

$$\frac{x_1 + x_2}{N(x_1) + N(x_2)} \in K$$

For all $\delta > 0$, we have

$$\frac{x_1}{N(x_1) + \delta} \in K, \quad \frac{x_2}{N(x_2) + \delta} \in K$$

. Since K is convex, we have the convex combination

$$\frac{1}{N(x_1) + N(x_2) + 2\delta} \left((N(x_2) + \delta) \frac{x_2}{N(x_2) + \delta} + (N(x_1) + \delta) \frac{x_1}{N(x_1) + \delta} \right) \in K$$

That is

$$\frac{x_1 + x_2}{N(x_1) + N(x_2) + 2\delta} \in K$$

for all $\delta > 0$. Then

$$N(x_1 + x_2) \leq N(x_1) + N(x_2) + 2\delta, \quad \forall \delta > 0 \implies N(x_1 + x_2) \leq N(x_1) + N(x_2)$$

2. $x \in K^\circ \iff N(x) < 1$, $x \in \text{Ext}(K) \iff N(x) > 1$, $x \in \partial K \iff N(x) = 1$. We define a map $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$.

$$\varphi(x) := \begin{cases} N(x) \frac{x}{\|x\|}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

We have

$$\|\varphi(x)\| = N(x)$$

Then φ maps K° on to $(\mathbb{D}^n)^\circ$, and maps ∂K on to $\mathbb{S}^{n-1}$. To show φ is continuous, only need to show it is continuous at 0. In fact, $\|\lim_{x \rightarrow 0} \varphi(x)\| = \lim_{x \rightarrow 0} \|\varphi(x)\| = \lim_{x \rightarrow 0} N(x) = 0$ since N is a norm. We define $\psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$:

$$\psi(x) := \begin{cases} \frac{\|x\|}{N(x)} x, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Then $\varphi \circ \psi = \psi \circ \varphi = \text{Id}$. The same skills can be used to show ψ is continuous at 0 as well. Finally, by the lemma above, X, Y are homeomorphic.

□

Problem 1.4.3

Let X, X' be topological spaces and $x \in X, x' \in X'$ be base points. Prove that the wedge sum $X \vee X'$ is homeomorphic to the subspace:

$$(X \times \{x'\}) \cup (\{x\} \times X') \subseteq X \times X'$$

Proof. $X \vee X' = (X \amalg X') / \{x \sim x'\}$. Considering map $\varphi : X \times X' \rightarrow X \vee X'$

$$\varphi(x_1, x_2) = \begin{cases} x_1, & x_2 = x' \\ x_2, & x_1 = x \end{cases}$$

$$\varphi(x_1, x_2) = (1 + x_2 - x')(x_1)$$

□

Problem 1.4.4

The torus $\mathbb{T}$ is the quotient of $[0, 1]^2$ by the equivalence relation generated by $(0, y) \sim (1, y)$ for all $x, y \in [0, 1]$. Prove that $\mathbb{T}$ is homeomorphic to :

1. The product $\mathbb{S}^1 \times \mathbb{S}^1$
2. The quotient of $\mathbb{R}^2$ under the action of the (discrete) group $\mathbb{Z}^2$ by translation.

Problem 1.4.5 ► Projective spaces

Let $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$. The n -dimensional projective space $\mathbb{K}\mathbb{P}^n$ is the orbit space of $\mathbb{K}^{n+1} \setminus \{0\}$ under the action of $\mathbb{K}^*$ by rescaling.

1. Prove that $\mathbb{R}\mathbb{P}^n$ is homeomorphic to the orbit space $\mathbb{S}^n \setminus \{\pm 1\}$, where $\{\pm 1\}$ acts by multiplication on $\mathbb{S}^n \subseteq \mathbb{R}^{n+1}$.
2. Prove that $\mathbb{C}\mathbb{P}^n$ is homeomorphic to the orbit space $\mathbb{S}^{2n+1} \setminus \mathbb{S}^1$, where $\mathbb{S}^1 = \{z \in \mathbb{C} : |z| = 1\}$ acts on $\mathbb{S}^{2n+1} \subseteq \mathbb{C}^{n+1}$ by multiplication.
3. Prove that $\mathbb{R}\mathbb{P}^1$ is homeomorphic to $\mathbb{S}^1$ and that $\mathbb{C}\mathbb{P}^1$ is homeomorphic to $\mathbb{S}^2$.

Problem 1.4.6

1. Let A be a compact subspace of a Hausdorff space X . Prove that X/A is Hausdorff.

Proof. $\forall [x], [y] \in X/A$.

1. $x \notin A, y \notin A$. Let $p : X \rightarrow X/A$ is the quotient map. Take a neighborhood of x, y , U_x, U_y , such that $U_x \cap U_y = \emptyset$, $U_x \cap A \neq \emptyset$, $U_y \cap A \neq \emptyset$.

Lemma 1.4.7

T_2 中的紧子集可以由点分离.

2. $x \in A, y \notin A$

□