

Contents

1	Introduction	2
1.1	Arithmetic in $[0, \infty]$	2
1.2	Metric Space	2
1.3	Limits of Set Sequences	3
2	Abstract Integration	6
2.1	The Concept of Measurability	6
2.2	Simple Functions	9
2.3	Measure	11
2.4	Integration of Positive Functions	13
2.5	Integration of Complex Functions	20
2.6	The Sets of Measure Zero	24
2.7	Exercises	30
3	Measure Theory	40
3.1	Outer Measure	40
	Index	44

Introduction

1.1 Arithmetic in $[0, \infty]$

1.2 Metric Space

Lemma 1.2.1

X is a metric space, $x \in X$, B_1, B_2 are two balls in X . If $x \in B_1 \cap B_2$, then x is the center of an open ball $B \subseteq B_1 \cap B_2$.

Proof. We may assume that B_1, B_2 have different center. Let $y_1 \neq y_2$,

$$B_1 = \{x : d(x, y_1) < r_1\}, \quad B_2 = \{x : d(x, y_2) < r_2\}$$

Take $x \in B_1 \cap B_2$, we claim that there exists $r_0 > 0$ such that $B := \{y : d(x, y) < r_0\} \subseteq B_1 \cap B_2$. Otherwise, for all $r > 0$, there exists a point y_r such that the following two hold at the same time

1. $d(x, y_r) < r$
2. $d(y_r, y_1) \geq r_1$ or $d(y_r, y_2) \geq r_2$.

Thus

$$d(x, y_1) \geq d(y_r, y_1) - d(x, y_r) > r_1 - r, \quad \text{or} \quad d(x, y_2) > r_2 - r$$

The above shows that

$$r > \min\{r_1 - d(x, y_1), r_2 - d(x, y_2)\}, \quad \forall r > 0$$

which is a contradiction. □

1.3 Limits of Set Sequences

Definition 1.3.1 ► Limit Inferior and Limit Superior

Let $\{A_n\}_{n=1}^{\infty}$ be a sequence of sets.

1. By the *Limit Inferior of the sequence*, we mean

$$\liminf_{n \rightarrow \infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=N}^{\infty} A_n = \{x : \exists N_0 \in \mathbb{N}, \forall n > N_0, x \in A_n\}$$

2. By the *Limit Superior of the sequence*, we mean

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} A_n = \{x : \forall N \in \mathbb{N}, \exists n \geq N, x \in A_n\}$$

Note.

1. Limit Inferior 包含那些“最终稳定下来”的元素，即从某个点之后就永远属于序列中所有后续集合的元素。
2. Limit Superior 包含那些“反复出现”的元素，即在无限多个集合中出现的元素。

Proposition 1.3.2

For any sequence of sets $\{A_n\}$, it always holds that:

$$\liminf_{n \rightarrow \infty} A_n \subseteq \limsup_{n \rightarrow \infty} A_n$$

Proof sketch.

直觉上是显然的，因为“最终稳定下来”的元素一定会“反复出现”。

Proof. It is obvious by the $\{x : x \in P\}$ form representation of the Limit Inferior and Superior. □

Definition 1.3.3 ► Limit of a Set Sequence

If the limit inferior and limit superior of a set sequence $\{A_n\}_{n=1}^{\infty}$ are equal, i.e., $\liminf_{n \rightarrow \infty} A_n = \limsup_{n \rightarrow \infty} A_n$, then we say the *limit* of the se-

quence exists, and it is defined as:

$$\lim_{n \rightarrow \infty} A_n = \liminf_{n \rightarrow \infty} A_n = \limsup_{n \rightarrow \infty} A_n$$

Proposition 1.3.4

Let $\{A_n\}_{n=1}^{\infty}$ be a sequence of sets.

1. If $\{A_n\}$ is an increasing sequence, then the limit of $\{A_n\}$ exists and

$$\lim_{n \rightarrow \infty} A_n = \bigcup_{n=1}^{\infty} A_n$$

2. If $\{A_n\}$ is a decreasing sequence, then the limit of $\{A_n\}$ exists and

$$\lim_{n \rightarrow \infty} A_n = \bigcap_{n=1}^{\infty} A_n$$

Proof. 1. If $\{A_n\}$ is increasing, then

$$\bigcap_{n=N}^{\infty} A_n = A_N$$

and

$$\bigcup_{n=N}^{\infty} A_n = \bigcup_{n=1}^{\infty} A_n, \forall N \in \mathbb{N}.$$

Thus

$$\liminf_{n \rightarrow \infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=N}^{\infty} A_n = \bigcup_{N=1}^{\infty} A_N$$

and

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=1}^{\infty} A_n = \bigcup_{n=1}^{\infty} A_n$$

which are the same.

2. If $\{A_n\}$ is decreasing, then

$$\bigcap_{n=N}^{\infty} A_n = \bigcap_{n=1}^{\infty} A_n$$

and

$$\bigcup_{n=N}^{\infty} A_n = A_N$$

Thus

$$\liminf_{n \rightarrow \infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=N}^{\infty} A_n = \bigcup_{N=1}^{\infty} \bigcap_{n=1}^{\infty} A_n = \bigcap_{n=1}^{\infty} A_n$$

and

$$\limsup_{n \rightarrow \infty} A_n = \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} A_n = \bigcap_{N=1}^{\infty} A_N$$

which are the same. □

Proposition 1.3.5

For a sequence of sets $\{A_n\}_{n=1}^{\infty}$ and their corresponding sequence of indicator functions $\{\chi_{A_n}(x)\}_{n=1}^{\infty}$:

- $\chi_{\liminf_{n \rightarrow \infty} A_n}(x) = \liminf_{n \rightarrow \infty} \chi_{A_n}(x)$
- $\chi_{\limsup_{n \rightarrow \infty} A_n}(x) = \limsup_{n \rightarrow \infty} \chi_{A_n}(x)$
- $\lim_{n \rightarrow \infty} A_n$ exists, then $\chi_{\lim_{n \rightarrow \infty} A_n}(x) = \lim_{n \rightarrow \infty} \chi_{A_n}(x)$

Proof sketch.

- 若 x 在 limit inferior 里面, 则 x 是“最终稳定”的, $\chi_{A_n}(x)$ 是关于 n “最终”恒为 1 的.
- 若 x 在 limit superior 里面, 则 x 是“反复出现”的, 即相当于 N 多大, 总会出现之后的某个 n 使得 $\chi_{A_n}(x) = 1$.
- 当极限存在时, 函数列 $\{\chi_{A_n}(x)\}_{n=1}^{\infty}$ 的 limit inferior 和 supperior 根据上两条相等, 等于极限集合的 χ .

Abstract Integration

2.1 The Concept of Measurability

Definition 2.1.1

1. A collection $\mathfrak{M}$ of subsets of a set X is said to be a σ -**algebra** in X if $\mathfrak{M}$ has the following properties:
 - (a) $X \in \mathfrak{M}$.
 - (b) If $A \in \mathfrak{M}$, then $A^c \in \mathfrak{M}$.
 - (c) If $A = \bigcup_{n=1}^{\infty} A_n$ and $A_n \in \mathfrak{M}$ for $n = 1, 2, 3, \dots$, then $A \in \mathfrak{M}$.
2. If $\mathfrak{M}$ is a σ -algebra in X , then X is called a **measurable** provided that $f^{-1}(V)$ is measurable set in X for every open set V in Y .
3. If X is a measurable space, Y is a topological space, and f is a mapping of X into Y , then f is said to be **measurable** provided that $f^{-1}(V)$ is a measurable set in X for every open set V in Y .

Note.

对比 topological space 的定义, σ -algebra 的定义是对称的, 即可测集的补集仍是可测的, 而对于拓扑空间, 开集的补集不是开集, 而是被定义为了闭集这样的对象.

Lemma 2.1.2

1. Let X be a measurable space, Y, Z be topological spaces. If $f : X \rightarrow Y$ is measurable, and if $g : Y \rightarrow Z$ is continuous, then $g \circ f : X \rightarrow Z$ is measurable.
2. Let u, v be real measurable functions on a measurable space X , let Φ be a continuous mapping of the plane into a topological space Y , and define

$$h(x) = \Phi(u(x), v(x))$$

for $x \in X$. Then $h : X \rightarrow Y$ is measurable.

Proof. 1. Take any open subset $V \subseteq Z$. Then $g^{-1}(V)$ is an open subset of

Y . Since f is measurable,

$$(g \circ f)^{-1}(V) = f^{-1}(g^{-1}(V))$$

is a measurable set.

2. Let $V \subseteq Y$ be a open subset. Then $\Phi^{-1}(V)$ is a open subset of $[-\infty, \infty]^2$. There exists countably many open sets of $[-\infty, \infty]$: $U_i, V_i, i \in \mathbb{N}$, such that

$$\Phi^{-1}(V) = \bigcup_i U_i \times V_i$$

Thus

$$\begin{aligned} h^{-1}(V) &= (u, v)^{-1} \left(\bigcup_i U_i \times V_i \right) \\ &= \bigcup_i (u, v)^{-1}(U_i \times V_i) \\ &= \bigcup_i (u^{-1}(U_i) \cap v^{-1}(V_i)) \end{aligned}$$

which is a measurable set.

□

Corollary 2.1.3

Let X be a measurable space. The following propositions hold

1. If $f = u + iv$, where u and v are real measurable functions on X , then f is a complex measurable function on X .
2. If $f = u + iv$ is a complex measurable function on X , then u, v , and $|f|$ are real measurable functions on X .
3. If f and g are complex measurable functions on X , then so are $f + g$ and fg .
4. If E is a measurable set in X and if

$$\chi_E(x) = \begin{cases} 1 & x \in E \\ 0 & x \notin E \end{cases}$$

then χ_E is a measurable function.

5. If f is a complex measurable function on X , there is a complex measurable function α on X such that $|\alpha| = 1$ and $f = \alpha |f|$.

Theorem 2.1.4

If $\mathcal{F}$ is any collection of subsets of X , there exists a smallest σ -algebra $\mathfrak{M}^*$ such that $\mathcal{F} \subseteq \mathfrak{M}^*$.

Definition 2.1.5 ▶ Borel

Let X be a topological space.

1. By **Borel σ -algebra**, we mean the smallest σ -algebra $\mathcal{B}$ in X such that every open set in X belongs to $\mathcal{B}$. The members of $\mathcal{B}$ are called the **Borel sets** of X .
2. All countable unions of closed sets and all countable intersections of open sets are Borel sets, which we called F_σ 's **and** G_δ 's, respectively.
3. By a **Borel function**, we mean a measurable function on the measurable space $(X, \mathcal{B})$.

Remark.

1. The letters F and G were used for closed and open sets, respectively, and σ refers to union, δ to intersection.
2. A continuous function is Borel-measurable, since the preimage of any open set is open and therefore a Borel set.

Theorem 2.1.6

Suppose $\mathfrak{M}$ is a σ -algebra in X , and Y is a topological space. Let f map X into Y .

1. If Ω is the collection of all sets $E \subseteq Y$ such that $f^{-1}(E) \in \mathfrak{M}$, then Ω is a σ -algebra in Y .
2. If f is measurable and E is a Borel set in Y , then $f^{-1}(E) \in \mathfrak{M}$.
3. If $Y = [-\infty, \infty]$ and $f^{-1}((\alpha, \infty]) \in \mathfrak{M}$ for every real α , then f is measurable.
4. If f is measurable, if Z is a topological space, if $g : Y \rightarrow Z$ is a Borel mapping, and if $h = g \circ f$, then $h : X \rightarrow Z$ is measurable.

Theorem 2.1.7

If $f_n : X \rightarrow [-\infty, \infty]$ is measurable, for $n = 1, 2, 3, \dots$, and

$$g = \sup_{n \geq 1} f_n, \quad h = \lim_{n \rightarrow \infty} \sup f_n,$$

then g and h are measurable.

Corollary 2.1.8

1. The limit of every pointwise convergent sequence of complex measurable functions is measurable.
2. If f and g are measurable (with range in $[-\infty, \infty]$), then so are $\max\{f, g\}$ and $\min\{f, g\}$. In particular, this is true of the functions

$$f^+ = \max\{f, 0\}, \quad f^- = -\min\{f, 0\}.$$

which are called the **positive part** and **negative part** of f , respectively.

Remark.

1. There are the standard representation

$$|f| = f^+ + f^-, \quad f = f^+ - f^-$$

2. And easy but (may) useful observation is: If $f = g - h$, $g \geq 0$, $h \geq 0$, then $f^+ \leq g$ and $f^- \leq h$.

2.2 Simple Functions

Definition 2.2.1

A complex function s on a measurable space X whose range consists of only finitely many points will be called a **simple function**. Among these are the nonnegative simple functions, whose range is a finite subset of $[0, \infty)$.

Specifically, if $\alpha_1, \dots, \alpha_n$ are distinct values of a simple function s , and if we set $A_i = \{x : s(x) = \alpha_i\}$, then clearly

$$s = \sum_{i=1}^n \alpha_i \chi_{A_i}.$$

Where χ_{A_i} is the characteristic function of A_i .

Remark.

- Here, we explicitly exclude ∞ from the values of a simple function.
- It is clear that s is measurable if and only if each of the sets A_i is measurable.

Theorem 2.2.2

Let $f : X \rightarrow [0, \infty]$ be measurable space. There exists simple measurable functions s_n on X such that

1. $0 \leq s_1 \leq s_2 \leq \dots \leq f$.
2. $s_n(x) \rightarrow f(x)$ as $n \rightarrow \infty$, for every $x \in X$.

Proof sketch.

We construct a sequence of Borel simple functions $\{\varphi_n(x)\}$ to act as an **identity**. 当 n 增大的同时, 我们同时让 $\varphi_n(x)$ 的单位逼近精度和单位逼近范围随着 n 提升. 并且在舍弃误差时, 总是向下取整, 使得该单位逼近是自下而上的.

Proof. For every $x \in [0, \infty]$, and for every $n \in \mathbb{N}$, there exists a unique integer $k_n(x)$, such that

$$k_n(x) 2^{-n} \leq x < (k_n(x) + 1) 2^{-n}$$

For every $n \in \mathbb{N}$, we define

$$\varphi_n(x) = \begin{cases} k_n(x) 2^{-n}, & 0 \leq x \leq n \\ n, & x \geq n \end{cases}$$

Each $\varphi_n(x)$ is then a Borel simple function. It is not hard to show that $0 \leq \varphi_1 \leq \varphi_2 \leq \dots \leq \text{Id}$. For each n , we define

$$s_n(x) := (\varphi_n \circ f)(x)$$

Then $\{s_n\}$ is a sequence of simple measurable functions such that $0 \leq s_1 \leq s_2 \leq \dots \leq f$. Since $\lim_{n \rightarrow \infty} \varphi_n(x) = x$, then $\lim_{n \rightarrow \infty} s_n(x) = f(x)$. $\square$

2.3 Measure

Definition 2.3.1 ► Measure and Measure Space

- (a) A **positive measure** is a function μ , defined on a σ -algebra $\mathfrak{M}$, whose range is in $[0, \infty]$ and which is **countably additive**. This means that if $\{A_i\}$ is a disjoint countable collection of members of $\mathfrak{M}$, then

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i).$$

To avoid trivialities, we shall also assume that $\mu(A) < \infty$ for at least one $A \in \mathfrak{M}$.

- (b) A **measure space** is a measurable space which has a positive measure defined on the σ -algebra of its measurable sets.
- (c) A **complex measure** is a complex-valued countably additive function defined on a σ -algebra.

Theorem 2.3.2

Let μ be a positive measure on a σ -algebra $\mathfrak{M}$. Then

1. $\mu(\emptyset) = 0$.
2. $\mu(A_1 \cup \dots \cup A_n) = \mu(A_1) + \dots + \mu(A_n)$ if $A_1, \dots, A_n$ are pairwise disjoint members of $\mathfrak{M}$.
3. $A \subset B$ implies $\mu(A) \leq \mu(B)$ if $A \in \mathfrak{M}, B \in \mathfrak{M}$.
4. $\mu(A_n) \rightarrow \mu(A)$ as $n \rightarrow \infty$ if $A = \bigcup_{n=1}^{\infty} A_n, A_n \in \mathfrak{M}$, and

$$A_1 \subset A_2 \subset A_3 \subset \dots$$

5. $\mu(A_n) \rightarrow \mu(A)$ as $n \rightarrow \infty$ if $A = \bigcap_{n=1}^{\infty} A_n, A_n \in \mathfrak{M}$,

$$A_1 \supset A_2 \supset A_3 \supset \dots,$$

and $\mu(A_1)$ is finite.

Proof. 1. Take $A \in \mathfrak{M}$ such that $\mu(A) < \infty$.¹ And let $A_2 = A_3 = \dots = \emptyset$, then $\mu(\emptyset) > 0$ leads to a contradiction to the countably additive.

2. Take $A_{n+1} = A_{n+2} = \dots = \emptyset$.

¹That is what we supposed at the definition of measure.

3. Note that $B = (B \setminus A) \cup A$, then by additivity

$$\mu(B) = \mu(A) + \mu(B \setminus A) \geq \mu(A)$$

4. Let $A_0 = \emptyset$, and let $B_n = A_n \setminus A_{n-1}$ for all $n \in \mathbb{N}$. Then $B_1, \dots, B_n$ are pairwise disjoint members of $\mathfrak{M}$ such that $A = \bigcup_{n=1}^{\infty} B_n$. We have

$$\mu(A) = \sum_{n=1}^{\infty} \mu(B_n) = \sum_{n=1}^{\infty} \mu(A_n \setminus A_{n-1})$$

If one of the $\mu(A_n)$ is ∞ , then $\lim_{n \rightarrow \infty} \mu(A_n)$ and $\mu(A)$ both are ∞ . Otherwise, we have

$$\mu(A_n \setminus A_{n-1}) = \mu(A_n) - \mu(A_{n-1}), \quad \forall n \in \mathbb{N}$$

Thus

$$\mu(A) = \sum_{n=1}^{\infty} \mu(A_n \setminus A_{n-1}) = \lim_{n \rightarrow \infty} \mu(A_n)$$

5. Let $B_n = A_n \setminus A_{n+1}$ for all $n \in \mathbb{N}$. $B_1, \dots, B_n$ are pairwise disjoint members of $\mathfrak{M}$ with finite measure, such that

$$A_1 \setminus A = A_1 \setminus \left(\bigcap_{n=1}^{\infty} A_n \right) = \bigcup_{n=1}^{\infty} (A_1 \setminus A_n) = \bigcup_{n=1}^{\infty} \left(\bigcup_{k=1}^n B_k \right) = \bigcup_{n=1}^{\infty} B_n$$

. We have

$$\mu(A_1 \setminus A) = \mu\left(\bigcup_{n=1}^{\infty} B_n\right),$$

where the RHS is $\mu(A_1) - \mu(A)$, and the LHS is

$$\sum_{n=1}^{\infty} (\mu(A_n) - \mu(A_{n+1})) = \mu(A_1) - \lim_{n \rightarrow \infty} \mu(A_{n+1})$$

Since $\mu(A_1) < \infty$, we have

$$\mu(A) = \lim_{n \rightarrow \infty} \mu(A_n)$$

□

Example 2.3.3

1. For any $E \subset X$, where X is any set, define $\mu(E) = \infty$ if E is an infinite set, and let $\mu(E)$ be the number of points in E if E is finite. This μ is called the **counting measure** on X .
2. Fix $x_0 \in X$, define $\mu(E) = 1$ if $x_0 \in E$ and $\mu(E) = 0$ if $x_0 \notin E$, for any $E \subset X$. This μ may be called the **unit mass concentrated at x_0** .
3. Let μ be the counting measure on the set $\{1, 2, 3, \dots\}$, let $A_n = \{n, n+1, n+2, \dots\}$. Then $\bigcap A_n = \emptyset$ but $\mu(A_n) = \infty$ for $n = 1, 2, 3, \dots$. This shows that the hypothesis

$$\mu(A_1) < \infty$$

is not superfluous in Theorem 2.3.2(5).

2.4 Integration of Positive Functions

Definition 2.4.1 ▶ Integration of Positive Functions

1. If $s : X \rightarrow [0, \infty)$ is a measurable simple function, of the form

$$s = \sum_{i=1}^n \alpha_i \chi_{A_i},$$

where $\alpha_1, \dots, \alpha_n$ are the distinct values of s , and if $E \in \mathfrak{M}$, we define

$$\int_E s \, d\mu = \sum_{i=1}^n \alpha_i \mu(A_i \cap E).$$

The convention $0 \cdot \infty = 0$ is used here; it may happen that $\alpha_i = 0$ for some i and that $\mu(A_i \cap E) = \infty$.

2. If $f : X \rightarrow [0, \infty]$ is measurable, and $E \in \mathfrak{M}$, we define

$$\int_E f \, d\mu = \sup \int_E s \, d\mu,$$

the supremum being taken over all simple measurable functions s such that $0 \leq s \leq f$. The left member above is called the **Lebesgue integral** of f over E , with respect to the measure μ . It is a number in $[0, \infty]$.

Remark.

We apparently have two definitions for $\int_E f d\mu$ if f is simple, they are the same.

Corollary 2.4.2

- (a) If $0 \leq f \leq g$, then $\int_E f d\mu \leq \int_E g d\mu$.
- (b) If $A \subset B$ and $f \geq 0$, then $\int_A f d\mu \leq \int_B f d\mu$.
- (c) If $f \geq 0$ and c is a constant, $0 \leq c < \infty$, then

$$\int_E cf d\mu = c \int_E f d\mu.$$

- (d) If $f(x) = 0$ for all $x \in E$, then $\int_E f d\mu = 0$, even if $\mu(E) = \infty$.
- (e) If $\mu(E) = 0$, then $\int_E f d\mu = 0$, even if $f(x) = \infty$ for every $x \in E$.
- (f) If $f \geq 0$, then $\int_E f d\mu = \int_X \chi_E f d\mu$.

Remark.

We can also regard the $\int_E f d\mu$ as the restricted function $f|_E$ integrates on the induced measure space E .

Proposition 2.4.3

Let s and t be nonnegative measurable simple functions on X . For $E \in \mathfrak{M}$, define

$$\varphi(E) = \int_E s d\mu.$$

Then φ is a measure on $\mathfrak{M}$. Also

$$\int_X (s + t) d\mu = \int_X s d\mu + \int_X t d\mu.$$

Note.

$\varphi(E) = \int_E s d\mu$ 就是依据 s 对测度 μ 进行一个有限的算数的加权. 自然地也就成为一个测度.

Proof. If $E_1, E_2, \dots$ are disjoint members of $\mathfrak{M}$ whose union is E , the countable

additivity of μ shows that

$$\begin{aligned}
 \varphi(E) &= \sum_{i=1}^n \alpha_i \mu(A_i \cap E) \\
 &= \sum_{i=1}^n \sum_{j=0}^{\infty} \alpha_i \mu(A_i \cap E_j) \\
 &= \sum_{j=0}^{\infty} \sum_{i=1}^n \alpha_i \mu(A_i \cap E_j) \\
 &= \sum_{j=0}^{\infty} \varphi(E_j)
 \end{aligned}$$

Also, $\varphi(\emptyset) \neq \infty$, μ is not identically ∞ .

Next, let $\beta_1, \dots, \beta_m$ be distinct values of t . And let $B_j := \{x : t(x) = \beta_j\}$. Then

$$\begin{aligned}
 \int_X (s+t) \, d\mu &= \sum_{i,j} \int_{A_i \cap B_j} (s+t) \, d\mu \\
 &= \sum_{i,j} \alpha_i \mu(A_i \cap B_j) + \sum_{i,j} \beta_j \mu(A_i \cap B_j) \\
 &= \sum_i \alpha_i \mu(A_i) + \sum_j \beta_j \mu(B_j) \\
 &= \int_X s \, d\mu + \int_X t \, d\mu
 \end{aligned}$$

□

Theorem 2.4.4 ► Lebesgue's Monotone Convergence Theorem

Let $\{f_n\}$ be a sequence of measurable functions on X , and suppose that

(a) $0 \leq f_1(x) \leq f_2(x) \leq \dots \leq \infty$ for every $x \in X$,

(b) $f_n(x) \rightarrow f(x)$ as $n \rightarrow \infty$, for every $x \in X$.

Then f is measurable, and

$$\int_X f_n \, d\mu \rightarrow \int_X f \, d\mu \quad \text{as } n \rightarrow \infty.$$

Proof. Since $\int_X f_n d\mu \leq \int_X f_{n+1} d\mu$ for all n , there exists an $\alpha > 0$ such that

$$\int_X f_n d\mu \rightarrow \alpha, \quad \text{as } n \rightarrow \infty$$

By Theorem 2.1.7, f is measurable, then by $\int_X f_n d\mu \leq \int_X f d\mu \forall n$,

$$\alpha \leq \int_X f d\mu$$

Take any simple function $0 \leq s \leq f$, and $0 \leq c < 1$. Define

$$E_n^s := \{x \in X : f_n(x) \geq cs(x)\}$$

Then since $\varphi(E) := \int_E s d\mu$ is a measure,

$$\alpha \geq \lim_{n \rightarrow \infty} \int_X f_n d\mu \geq \lim_{n \rightarrow \infty} \int_{E_n^s} f_n d\mu \geq c \lim_{n \rightarrow \infty} \int_{E_n^s} s d\mu = c \lim_{n \rightarrow \infty} \varphi(E_n^s) = c\varphi(X)$$

That is

$$\alpha \geq c \int_X s d\mu$$

Since s is arbitrary,

$$\alpha \geq c \int_X f d\mu$$

and since c is arbitrary,

$$\alpha \geq \int_X f d\mu$$

which completes the proof. □

Corollary 2.4.5

Let $f, g : X \rightarrow [0, \infty]$ be measurable functions on X . Then $f + g$ is measurable and

$$\int_X (f + g) d\mu = \int_X f d\mu + \int_X g d\mu$$

Proof. Let $\{s_n\}, \{s'_n\}$ be simple function sequences in Theorem 2.2.2 of f, g ,

respectively . Then by the theorem above and the Proposition 2.4.3

$$\begin{aligned}\int_X (f + g) \, d\mu &= \lim_{n \rightarrow \infty} \int_X (s_n + s'_n) \, d\mu = \lim_{n \rightarrow \infty} \int_X s_n \, d\mu + \lim_{n \rightarrow \infty} \int_X s'_n \, d\mu \\ &= \int_X f \, d\mu + \int_X g \, d\mu\end{aligned}$$

□

Corollary 2.4.6

If $f_n : X \rightarrow [0, \infty]$ is **measurable**, for $n = 1, 2, 3, \dots$, and

$$f(x) = \sum_{n=1}^{\infty} f_n(x) \quad (x \in X),$$

then

$$\int_X f \, d\mu = \sum_{n=1}^{\infty} \int_X f_n \, d\mu.$$

Proof. Let $g_n(x) = \sum_{k=1}^n f_k(x)$. Then $g_1 \leq g_2 \leq \dots \leq \infty$ and $f = \lim_{n \rightarrow \infty} g_n(x)$. Applying Monotone Convergence Theorem to $\{g_n\}$ and f , we have

$$\int_X f \, d\mu = \lim_{n \rightarrow \infty} \int_X \sum_{k=1}^n f_k(x) \, d\mu = \lim_{n \rightarrow \infty} \sum_{k=1}^n \int_X f_k(x) \, d\mu = \sum_{k=1}^{\infty} \int_X f_k \, d\mu$$

□

Corollary 2.4.7

If $a_{ij} \geq 0$ for i and $j = 1, 2, 3, \dots$, then

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij} = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}.$$

Proof. Let μ be counting measure on $\mathbb{Z}$. Let

$$f_i(j) = a_{ij}, \quad f(k) = \sum_{i=1}^{\infty} f_i(k) = \sum_{i=1}^{\infty} a_{ik}$$

By Monotone Convergence Theorem, for any measurable g ,

$$\int_{\mathbb{Z}} g \, d\mu = \int_{\mathbb{Z}} \lim_{n \rightarrow \infty} \chi(\{1, \dots, n\}) g \, d\mu = \lim_{n \rightarrow \infty} \int_{\mathbb{Z}} \chi(\{1, \dots, n\}) g \, d\mu = \sum_{k=-\infty}^{\infty} g(k)$$

Thus

$$\int_{\mathbb{Z}} f \, d\mu = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}$$

And we have

$$\sum_{i=1}^{\infty} \int_X f_i \, d\mu = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij}$$

The above corollary shows that

$$\int_{\mathbb{Z}} f \, d\mu = \sum_{i=1}^{\infty} \int_X f_i \, d\mu$$

That is

$$\sum_{i=1}^{\infty} \sum_{j=1}^{\infty} a_{ij} = \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} a_{ij}$$

□

Lemma 2.4.8 ► Fatou's Lemma

If $f_n : X \rightarrow [0, \infty]$ is **measurable**, for each positive integer n , then

$$\int_X \left(\liminf_{n \rightarrow \infty} f_n \right) d\mu \leq \liminf_{n \rightarrow \infty} \int_X f_n d\mu. \quad (1)$$

Note.

有了 Monotone Convergence Theorem 之后, 关键的不等式就剩下

$$\int_X \inf_{n \geq k} f_n d\mu \leq \inf_{n \geq k} \int_X f_n d\mu$$

它的含义是局部最小的积分小于等于积分的最小.

Proof. Let

$$g_k(x) = \inf_{n \geq k} f_n(x)$$

Then

$$\lim_{n \rightarrow \infty} \inf f_n = \lim_{n \rightarrow \infty} g_n =: g$$

where $\{g_k\}$, g are all measurable. It is obvious that $\{g_k\}$ is increasing. Thus

$$\lim_{n \rightarrow \infty} \int_X g_n d\mu = \int_X g d\mu = \int_X \left(\lim_{n \rightarrow \infty} \inf f_n \right) d\mu$$

Only need to show that

$$\int_X \inf_{n \geq k} f_n d\mu \leq \inf_{n \geq k} \int_X f_n d\mu$$

which is true since

$$\int_X \inf_{n \geq k} f_n d\mu \leq \int_X f_n d\mu, \text{ for all } n \geq k$$

□

Theorem 2.4.9

Suppose $f : X \rightarrow [0, \infty]$ is measurable, and

$$\varphi(E) = \int_E f d\mu \quad (E \in \mathfrak{M}).$$

Then φ is a measure on $\mathfrak{M}$, and

$$\int_X g d\varphi = \int_X gf d\mu$$

for every measurable g on X with range in $[0, \infty]$.

Note.

将 measurable function f 视为对 measure μ 的某种 countable 次算数加权操作. 那么得益于 μ 对 countable 操作的相容性, φ 也就成为一个测度. φ 的积分行为也验证了这种看法的合理性.

Proof. • Let $E_1, \dots, E_n, \dots$ be disjoint measurable sets such that $E = \bigcup_{n=1}^{\infty} E_n$. Then $\{\sum_{k=1}^n \chi_{E_k} f\}$ be increasing measurable function sequence such that $\lim_{n \rightarrow \infty} \sum_{k=1}^n \chi_{E_k} f = f$. The Monotone Convergence Theorem shows that

$$\begin{aligned} \sum_{n=1}^{\infty} \varphi(E_n) &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \int_{E_k} f d\mu = \lim_{n \rightarrow \infty} \int_E \sum_{k=1}^n \chi_{E_k} f d\mu \\ &= \int_E f d\mu = \varphi(E) \end{aligned}$$

Since $\varphi(\emptyset) = 0$. φ is a measure on X .

- To show the second results. See that it is true for $g = \chi_E$. Thus it is true for g is a simple function. And the general case follows from the Monotone Convergence Theorem.

□

2.5 Integration of Complex Functions

Definition 2.5.1 ▶ Lebegsgue Integrability

We define $L^1(\mu)$ to be the collection of all complex measurable functions f on X for which

$$\int_X |f| d\mu < \infty.$$

Note that the measurability of f implies that of $|f|$, hence the above integral is defined.

The members of $L^1(\mu)$ are called *Lebesgue integrable functions* (with respect to μ) or *summable functions*.

Definition 2.5.2 ► Complex Integration

If $f = u + iv$, where u and v are real measurable functions on X , and if $f \in L^1(\mu)$, we define

$$\int_E f d\mu = \int_E u^+ d\mu - \int_E u^- d\mu + i \int_E v^+ d\mu - i \int_E v^- d\mu \quad (1)$$

for every measurable set E .

Remark.

We have $u^+ \leq |u| < |f|$, etc., so that each of these four integrals is finite. Thus (1) defines the integral on the left as a complex number.

Definition 2.5.3 ► Real Integration

Occasionally it is desirable to define the integral of a measurable function f with range in $[-\infty, \infty]$ to be

$$\int_E f d\mu = \int_E f^+ d\mu - \int_E f^- d\mu, \quad (2)$$

provided that at least one of the integrals on the right of (2) is finite. The left side of (2) is then a number in $[-\infty, \infty]$.

Theorem 2.5.4

Suppose f and $g \in L^1(\mu)$ and α and β are complex numbers. Then $\alpha f + \beta g \in L^1(\mu)$, and

$$\int_X (\alpha f + \beta g) d\mu = \alpha \int_X f d\mu + \beta \int_X g d\mu$$

Proof sketch.

可积性来自于模长的三角不等式. 主要证明可加性. 将 $f + g = h$ 的正负部分到两边, 就都变成了非负函数. 积分即可.

Theorem 2.5.5

If $f \in L^1(\mu)$, then

$$\left| \int_X f d\mu \right| \leq \int_X |f| d\mu.$$

Note.

被缩放的部分实际上是 f 因改变了方向产生的损耗.

Proof. Let $z = \int_X f d\mu$, z is a complex number. There exists a complex number α ($\alpha = \frac{\bar{z}}{|z|}$), such that $|\alpha| = 1$, and

$$\left| \int_X f d\mu \right| = \alpha z = \alpha \int_X f d\mu = \int_X \alpha f d\mu$$

Suppose $\alpha f = u + iv$, since $\int_X \alpha f d\mu$ is a real number, $\int_X v d\mu = 0$. We have

$$\int_X \alpha f d\mu = \int_X u d\mu \leq \int_X |\alpha f| d\mu = \int_X |f| d\mu$$

□

Theorem 2.5.6 ▶ Lebesgue's Dominated Convergence Theorem

Suppose $\{f_n\}$ is a sequence of complex measurable functions on X such that

$$f(x) = \lim_{n \rightarrow \infty} f_n(x)$$

exists for every $x \in X$. If there is a function $g \in L^1(\mu)$ such that

$$|f_n(x)| \leq g(x) \quad (n = 1, 2, 3, \dots; x \in X),$$

then $f \in L^1(\mu)$,

$$\lim_{n \rightarrow \infty} \int_X |f_n - f| d\mu = 0,$$

and

$$\lim_{n \rightarrow \infty} \int_X f_n d\mu = \int_X f d\mu.$$

Proof sketch.

$\{f_n\}$ 不一定单调, 能将极限从积分外塞进积分里的工具, 目前我们只有 Fatou's Lemma, 但是它原始的样子跟我们需要的不等号是反向的. 自然的想法是填一个负号, 这又会导致函数失去非负性, 而控制函数 g 恰好能弥补这一点不足. 实现起来, 我们从 g 上扣去 f, f_n , 而非直接拿出 f, f_n 去做.

Proof. Since $|f| \leq g$ and f is measurable, $f \in L^1(\mu)$. Function $(2g - |f_n - f|)$ is a nonnegative measurable function. Applying Fatou's Lemma to it, we have

$$\int_X \liminf_{n \rightarrow \infty} (2g - |f_n - f|) d\mu \leq \liminf_{n \rightarrow \infty} \int_X (2g - |f_n - f|) d\mu$$

where

$$\liminf_{n \rightarrow \infty} (2g - |f_n - f|) = 2g - \limsup_{n \rightarrow \infty} |f_n - f| = 2g$$

$$\liminf_{n \rightarrow \infty} \int_X (2g - |f_n - f|) d\mu = 2 \int_X g d\mu - \limsup_{n \rightarrow \infty} \int_X |f_n - f| d\mu$$

The original inequality comes to

$$2 \int_X g d\mu \leq 2 \int_X g d\mu - \limsup_{n \rightarrow \infty} \int_X |f_n - f| d\mu$$

Since $\int_X g d\mu$ is finite, we can subtract $2 \int_X g d\mu$ both sides and get

$$\limsup_{n \rightarrow \infty} \int_X |f_n - f| d\mu \leq 0$$

Thus

$$\lim_{n \rightarrow \infty} \int_X |f_n - f| d\mu = 0$$

Finally, we have

$$\begin{aligned} \left| \lim_{n \rightarrow \infty} \int_X f_n \, d\mu - \int_X f \, d\mu \right| &= \left| \lim_{n \rightarrow \infty} \int_X (f_n - f) \, d\mu \right| \\ &\leq \lim_{n \rightarrow \infty} \int_X |f_n - f| \, d\mu \\ &= 0 \end{aligned}$$

which completes the proof. □

2.6 The Sets of Measure Zero

Definition 2.6.1

Let P be a property which a point x may or may not have. If μ is a measure on a σ -algebra $\mathfrak{M}$ and if $E \in \mathfrak{M}$, the statement “ P holds **almost everywhere on** E ” (abbreviated to “ P holds a.e. on E ”) means that there exists an $N \in \mathfrak{M}$ such that $\mu(N) = 0$, $N \subset E$, and P holds at every point of $E - N$.

Remark.

This concept of a.e. depends of course very strongly on the given measure, and we shall write “a.e. $[\mu]$ ” whenever clarity requires that the measure be indicated.

Example 2.6.2

For example, if f and g are measurable functions and if

$$\mu(\{x : f(x) \neq g(x)\}) = 0,$$

we say that $f = g$ a.e. $[\mu]$ on X , and we may write $f \sim g$.

Remark.

This is easily seen to be an equivalence relation. The transitivity ($f \sim g$ and $g \sim h$ implies $f \sim h$) is a consequence of the fact that the union of two sets of measure 0 has measure 0.

Proposition 2.6.3 ▶ Completion of a Measure

If $f \sim g$, then, for every $E \in \mathfrak{M}$,

$$\int_E f d\mu = \int_E g d\mu.$$

Theorem 2.6.4

Let $(X, \mathfrak{M}, \mu)$ be a measure space, let $\mathfrak{M}^*$ be the collection of all $E \subset X$ for which there exist sets A and $B \in \mathfrak{M}$ such that $A \subset E \subset B$ and $\mu(B - A) = 0$, and define $\mu(E) = \mu(A)$ in this situation. Then $\mathfrak{M}^*$ is a σ -algebra, and μ is a measure on $\mathfrak{M}^*$.

Remark.

This extended measure μ is called **complete**, since all subsets of sets of measure 0 are now measurable; the σ -algebra $\mathfrak{M}^*$ is called the **μ -completion** of $\mathfrak{M}$. The theorem says that every measure can be completed, so, whenever it is convenient, we may assume that any given measure is complete; this just gives us more measurable sets, hence more measurable functions.

Proof. We begin by checking that μ is well defined for every $E \in \mathfrak{M}^*$. If for $i = 1, 2$, $A_i \subseteq E \subseteq B_i$, where $A_i, B_i \in \mathfrak{M}$, $\mu(B_i - A_i) = 0$. We need to show that $\mu(A_1) = \mu(A_2)$. Since $A_1 \subseteq E \subseteq B_2$,

$$\mu(A_1) \leq \mu(B_2) = \mu(A_2) + \mu(B_2 - A_2) = \mu(A_2)$$

By symmetric,

$$\mu(A_2) \leq \mu(A_1)$$

Thus $\mu(A_1) = \mu(A_2)$, $\mu(E)$ is well defined. Next, let us verify that $\mathfrak{M}^*$ has the three defining properties of a σ -algebra.

1. It is clearly that $X \in \mathfrak{M}^*$.
2. If $A \subseteq E \subseteq B$, then $B^c \subseteq E^c \subseteq A^c$. And since $\mu(A^c - B^c) = \mu(A^c \cap B) = \mu(B - A) = 0$, $E^c \in \mathfrak{M}^*$.
3. Suppose $A_i \subseteq E_i \subseteq B_i$, $i = 1, 2, \dots$, where $A_i, B_i \in \mathfrak{M}$, $\mu(B_i - A_i) = 0$.

Let $E = \bigcup E_i, A = \bigcup A_i, B = \bigcup B_i$. Since

$$B - A = \bigcup_i (B_i - A) \subseteq \bigcup_i (B_i - A_i)$$

,

$$\mu(B - A) \leq \sum_i \mu(B_i - A_i) = 0$$

Then by $A \subseteq E \subseteq B$, it follows that $E \in \mathfrak{M}^*$.

Finally, we show the countable additive of μ . If the above sets E_i are disjoint, the same is true of the sets A_i , then

$$\mu(E) = \mu(A) = \sum_i \mu(A_i) = \sum_i \mu(E_i)$$

□

Definition 2.6.5 ▶ More Measurable Functions

We say a function f defined on a set $E \in \mathfrak{M}$ is **measurable on X** . If $\mu(E^c) = 0$ and if $f^{-1}(V) \cap E$ is a measurable for every open set V .

Remark.

- If we define $f(x) = 0$ for $x \in E^c$, we obtain a measurable function on X , in the old sense.
- If our measure happens to be complete, we can define f on E^c in a perfectly arbitrary manner, and we still get a measurable function. The integral of f over any set $A \in \mathfrak{M}$ is independent of the definition of f on E^c ; therefore this definition need not even be specified at all.

Theorem 2.6.6

Suppose $\{f_n\}$ is a sequence of complex measurable functions defined a.e. on X such that

$$\sum_{n=1}^{\infty} \int_X |f_n| d\mu < \infty. \quad (1)$$

Then the series

$$f(x) = \sum_{n=1}^{\infty} f_n(x) \quad (2)$$

converges for almost all x , $f \in L^1(\mu)$, and

$$\int_X f \, d\mu = \sum_{n=1}^{\infty} \int_X f_n \, d\mu. \quad (3)$$

Remark.

- 现在, 考察积分和求和是否可交换, 只需要检查模长积分的级数是否收敛.
- 即便 f_n 在 X 上处处被定义, 级数收敛的地方也只是 *almost everywhere*.

Note.

此时, 控制函数就是绝对级数本身, 它在一个满测集上将原级数控制住. 其特殊性间接给出了原级数的几乎处处收敛性质.

Proof. Let S_n be the set on which f_n is defined, so that $\mu(S_n^c) = 0$. Let $S = \bigcap S_n$. Put $\varphi(x) = \sum |f_n(x)|$, for $x \in S$. By Monotone Convergence Theorem

$$\int_S \varphi \, d\mu \leq \sum_{n=1}^{\infty} \int_S |f_n| \, d\mu < \infty$$

Let $E = \{x \in S : \varphi(x) < \infty\}$. Then it follows that $\mu(E^c) = 0$. We have

$$\sum_{n=1}^{\infty} |f_n(x)| < \infty, \quad \forall x \in E$$

Thus

$$f(x) = \sum_{n=1}^{\infty} f_n(x)$$

converges on E , which is a set with its complement measure zero. That is to say f converges for almost all x . And since $f \in L^1(\mu|_E)$, $\mu(E) = 0$, we have

$f \in L^1(\mu)$. Finally, by Monotone Convergence Theorem,

$$\int_X f \, d\mu = \int_E f \, d\mu = \sum_{n=1}^{\infty} \int_E f_n \, d\mu = \sum_{n=1}^{\infty} \int_X f_n \, d\mu$$

□

Theorem 2.6.7

1. Suppose $f : X \rightarrow [0, \infty]$ is measurable, $E \in \mathfrak{M}$, and $\int_E f \, d\mu = 0$. Then $f = 0$ a.e. on E .
2. Suppose $f \in L^1(\mu)$ and $\int_E f \, d\mu = 0$ for every $E \in \mathfrak{M}$. Then $f = 0$ a.e. on X .
3. Suppose $f \in L^1(\mu)$ and

$$\left| \int_X f \, d\mu \right| = \int_X |f| \, d\mu$$

Then there is a constant α such that $\alpha f = |f|$ a.e. on X .

Note.

3. 这个积分等式的意义是不存在积分项因方向改变产生的损耗. 结论就是说如果这种损耗不存在, 那么 f 几乎不改变方向.

Proof. 1. Let $A_n = \left\{ x \in E : f(x) \geq \frac{1}{n} \right\}$. Then

$$0 = \int_E f \, d\mu \geq \int_{A_n} f \, d\mu \geq \mu(A_n) \frac{1}{n}.$$

It follows that

$$\mu(A_n) = 0, \quad \forall n = 1, 2, \dots$$

Let $A = \bigcup_{n=1}^{\infty} A_n$. Then $\mu(A) = 0$, and $f \equiv 0$ on $E \setminus A$. Thus $f = 0$ a.e. on E .

2. Suppose $f = u + iv$. Let $E = \{x \in X : u(x) \geq 0\}$. Since $\operatorname{Re} f = u^+$ on E , then

$$\int_E u^+ \, d\mu = 0$$

It follows by 1. that $\mu^+ = 0$ a.e. on E . Since $\mu^+(X \setminus E) \equiv 0$, $\mu^+ = 0$ a.e.

on X . We conclude similarly that

$$u^- = v^+ = v^- = 0, \quad a.e.$$

3. Let $z = \int_X f d\mu$. Then z is a complex number. Take $\alpha = \bar{z}/|z|$. Then

$$\alpha z = |z| = \left| \int_X f d\mu \right|$$

Where

$$\alpha z = \alpha \int_X f d\mu = \int_X \alpha f d\mu$$

It follows that

$$\int_X \alpha f d\mu = \left| \int_X f d\mu \right| = \int_X |f| d\mu \implies \int_X (\alpha f - |f|) d\mu = 0$$

which shows that

$$\alpha f = |f|, \quad a.e. \text{ on } X$$

□

Theorem 2.6.8

Suppose $\mu(X) < \infty$, $f \in L^1(\mu)$, S is a closed set in the complex plane, and the averages

$$A_E(f) = \frac{1}{\mu(E)} \int_E f d\mu$$

lie in S for every $E \in \mathfrak{M}$ with $\mu(E) > 0$. Then $f(x) \in S$ for almost all $x \in X$.

Proof. Let Δ be circular disk, with centre α and radius r . And suppose that Δ lies in S^c . S^c can be written as the union of countably many such disks. Thus, we only need to show that $\mu(E) = 0$, where $E = f^{-1}(\Delta)$. If $\mu(E) > 0$, then

$$|A_E(f) - \alpha| = \frac{1}{\mu(E)} \left| \int_E (f - \alpha) d\mu \right| \leq \frac{1}{\mu(E)} \int_E |f - \alpha| d\mu \leq r$$

Which leads to a contradiction, since $A_E(f) \in S$. Hence $\mu(E) = 0$. □

Theorem 2.6.9

Let $\{E_k\}$ be a sequence of measurable sets in X , such that

$$\sum_{k=1}^{\infty} \mu(E_k) < \infty.$$

Then almost all $x \in X$ lie in at most finitely many of the sets E_k .

Proof. Let

$$g(x) = \sum_{k=1}^{\infty} \chi_{E_k}(x)$$

Then g is measurable. By Monotone Convergence Theorem,

$$\int_X g(x) \, d\mu = \sum_{k=1}^{\infty} \int_X \chi_{E_k}(x) \, d\mu = \sum_{k=1}^{\infty} \mu(E_k) < \infty$$

Thus $g \in L^1(\mu)$. There is a $E \in \mathfrak{M}$, $\mu(E^c) = 0$, such that $g^{-1}(\infty) \cap E = \emptyset$. If $x \in X$ lie in infinitely many of the sets E_k . Then $g(x) = \infty$, it follows that $x \in E^c$. Thus, points that lie in infinitely many of the sets E_k will always lie in E^c as well, whose measure is zero. $\square$

2.7 Exercises

Note.

关于积分的问题有两种常用标准技巧。一种是按取值划分集合。一种用可控的函数列逼近。

Problem 2.7.1

Does there exist an infinite σ -algebra which has only countably many members?

Proof.

$\square$

Problem 2.7.2

Prove that if f is a real function on a measurable space X such that $\{x : f(x) \geq r\}$ is measurable for every rational r , then f is measurable.

Proof. Take any real number a . We have

$$f^{-1}((a, \infty]) = \bigcup_{r \in \mathbb{Q}, r > a} f^{-1}([r, \infty])$$

which is a countably many union of measurable sets. Thus $f^{-1}((a, \infty])$ is measurable for all a . It follows that f is measurable. $\square$

Problem 2.7.3

Let $\{a_n\}$ and $\{b_n\}$ be sequences in $[-\infty, \infty]$, and prove the following assertions:

(a)

$$\limsup_{n \rightarrow \infty} (-a_n) = -\liminf_{n \rightarrow \infty} a_n.$$

(b)

$$\limsup_{n \rightarrow \infty} (a_n + b_n) \leq \limsup_{n \rightarrow \infty} a_n + \limsup_{n \rightarrow \infty} b_n$$

provided none of the sums is of the form $\infty - \infty$.

(c) If $a_n \leq b_n$ for all n , then

$$\liminf_{n \rightarrow \infty} a_n \leq \liminf_{n \rightarrow \infty} b_n.$$

Show by an example that strict inequality can hold in (b).

Proof. 1.

$$\limsup_{n \rightarrow \infty} (-a_n) = \lim_{n \rightarrow \infty} \sup_{k \geq n} \{-a_k\} = -\lim_{n \rightarrow \infty} \inf_{k \geq n} \{a_k\} = -\liminf_{n \rightarrow \infty} a_n$$

2.

$$\begin{aligned}
\limsup_{n \rightarrow \infty} (a_n + b_n) &= \lim_{n \rightarrow \infty} \sup_{k \geq n} \{a_k + b_k\} \\
&\leq \lim_{n \rightarrow \infty} \left(\sup_{k \geq n} \{a_k\} + \sup_{k \geq n} \{b_k\} \right) \\
&= \lim_{n \rightarrow \infty} \sup_{k \geq n} \{a_k\} + \lim_{n \rightarrow \infty} \sup_{k \geq n} \{b_k\} \\
&= \limsup_{n \rightarrow \infty} a_n + \limsup_{n \rightarrow \infty} b_n
\end{aligned}$$

Let $\{a_n\} = \{(-1)^n\}$, $\{b_n\} = \{(-1)^{n+1}\}$. Then the RHS of the inequality is $1 + 1 = 2$. The LHS is 0.

3. If $a_n \leq b_n$ for all n , then

$$\inf_{k \geq n} \{a_k\} \leq \inf_{k \geq n} \{b_k\}$$

Thus

$$\liminf_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \inf_{k \geq n} \{a_k\} \leq \lim_{n \rightarrow \infty} \inf_{k \geq n} \{b_k\} = \liminf_{n \rightarrow \infty} b_n$$

□

Problem 2.7.4

(a) Suppose $f : X \rightarrow [-\infty, \infty]$ and $g : X \rightarrow [-\infty, \infty]$ are measurable. Prove that the sets

$$\{x : f(x) < g(x)\}, \{x : f(x) = g(x)\}$$

are measurable.

(b) Prove that the set of points at which a sequence of measurable real-valued functions converges (to a finite limit) is measurable.

Proof. 1. Let

$$h = \begin{cases} g - f, & x \in f^{-1}((-\infty, \infty)) \\ 0, & f(x) = g(x) = -\infty \\ 0, & f(x) = g(x) = \infty \\ -1, & f(x) = \infty, g(x) < \infty \\ 1, & f(x) = -\infty, g(x) > -\infty \end{cases}$$

It is easy to see that h is measurable. Thus

$$\{x : f(x) < g(x)\} = h^{-1}((0, \infty])$$

is measurable. And

$$\{x : f(x) = g(x)\} = h^{-1}(0)$$

is measurable.

2. Let $\{f_n\}$ be a sequence of measurable real-valued functions. Take $x \in X$. $\{f_n(x)\}$ is converges. If and only if for any $k \in \mathbb{N}$, there exists $N \in \mathbb{N}$, such that for all $m, n \geq N$, $|f_m(x) - f_n(x)| \leq \frac{1}{k}$. That is to say

$$\{x : \{f_n(x)\} \text{ converges} \} = \bigcap_{k=1}^{\infty} \bigcup_{N=1}^{\infty} \bigcap_{m=N}^{\infty} \bigcap_{n=N}^{\infty} (|f_m - f_n|)^{-1}\left((-\infty, \frac{1}{k}]\right)$$

which is a measurable set.

□

Problem 2.7.5

Let X be an uncountable set, let $\mathfrak{M}$ be the collection of all sets $E \subset X$ such that either E or E^c is at most countable, and define $\mu(E) = 0$ in the first case, $\mu(E) = 1$ in the second. Prove that $\mathfrak{M}$ is a σ -algebra in X and that μ is a measure on $\mathfrak{M}$. Describe the corresponding measurable functions and their integrals.

Proof. We will first show $\mathfrak{M}$ is a σ -algebra step by step.

1. It is clear that if $E \in \mathfrak{M}$, E^c is as well.
2. Let $E_1, E_2, \dots$ be members of $\mathfrak{M}$. If all of E_n is countable, then $\bigcup_{n=1}^{\infty} E_n$ is at most countable. Otherwise, we may assume that E_1 is uncountable, then E_1^c is at most countable.

$$X \setminus \bigcup_{n=1}^{\infty} E_n = E_1^c \cap \bigcap_{n=2}^{\infty} E_n^c$$

is at most countable. Hence $\bigcup_{n=1}^{\infty} E_n \in \mathfrak{M}$.

3. Since $\emptyset$ is at most countable, $X \in \mathfrak{M}$.

The above shows that $\mathfrak{M}$ is a σ -algebra. To see μ is a measure, take $E_1, E_2, \dots \in \mathfrak{M}$ with any two of them have empty intersection. If all of E_n is countable, then $\bigcup_{n=1}^{\infty} E_n$ is as well. We have

$$\sum_{n=1}^{\infty} \mu(E_n) = 0 = \mu\left(\bigcup_{n=1}^{\infty} E_n\right)$$

If one of the sets E_n is uncountable, say E_1 . If $E_i, i \geq 2$ is uncountable.

$$X = (\emptyset)^c = (E_1 \cap E_2)^c = E_1^c \cup E_2^c$$

is at most countable, which leads to a contradiction. Thus $E_i, i \geq 2$ is at most countable. Hence

$$\sum_{n=1}^{\infty} \mu(E_n) = 1 + 0 + \dots + 0 = 1 = \mu\left(\bigcup_{n=1}^{\infty} E_n\right).$$

It follows that μ is a measure on $\mathfrak{M}$, since $\mu(\emptyset) = 0$.

To describe the measurable functions. If f is not *a.e.* constant, then there are two value a, b of f , such that $f^{-1}(a), f^{-1}(b)$ is uncountable. Then

$$f^{-1}(b) \subseteq (f^{-1}(a))^c$$

is at most countable, which leads to a contradiction. Thus f is *a.e.* constant. Take any measurable function f . Suppose that $E = f^{-1}(c), \mu(E) \neq 0$. Then $\mu(E^c) = 0$. It follows that f is integrable and

$$\int_X f \, d\mu = \int_E f \, d\mu = c\mu(E) = c$$

□

Problem 2.7.6

Suppose $f_n : X \rightarrow [0, \infty]$ is measurable for $n = 1, 2, 3, \dots$, $f_1 \geq f_2 \geq f_3 \geq \dots \geq 0$, $f_n(x) \rightarrow f(x)$ as $n \rightarrow \infty$, for every $x \in X$, and $f_1 \in L^1(\mu)$. Prove that then

$$\lim_{n \rightarrow \infty} \int_X f_n \, d\mu = \int_X f \, d\mu$$

and show that this conclusion does not follow if the condition " $f_1 \in L^1(\mu)$ " is omitted.

Proof. Let $g_n = f_1 - f_n$, then

$$0 = g_1 \leq g_2 \leq \cdots, \quad g_n(x) \rightarrow f_1(x) - f(x) \text{ as } n \rightarrow \infty$$

Applying the Monotone Convergence Theorem to $\{g_n\}$, we have

$$\int_X f_1 d\mu - \lim_{n \rightarrow \infty} \int_X f_n d\mu = \lim_{n \rightarrow \infty} \int_X g_n d\mu = \int_X (f_1 - f) d\mu = \int_X f_1 d\mu - \int_X f d\mu$$

Since $\int_X f_1 d\mu < \infty$, subtracting $\int_X f_1 d\mu$ both sides, we have

$$\lim_{n \rightarrow \infty} \int_X f_n d\mu = \int_X f d\mu.$$

Let $X = \mathbb{R}$ with borel measure, and let $f_n = \chi_{\mathbb{R} \setminus [0, n]} \frac{1}{n}$. Then $f_n(x) \rightarrow f(x) = 0$, $\int_X f d\mu = 0$. But

$$\int_X f_n d\mu = \infty, \quad \forall n$$

Thus the condition " $f_1 \in L^1(\mu)$ " can not be omitted. □

Problem 2.7.7

Put $f_n = \chi_E$ if n is odd, $f_n = 1 - \chi_E$ if n is even. What is the relevance of this example to Fatou's lemma?

Proof.

$$\liminf_{n \rightarrow \infty} f_n(x) \equiv 0, \quad \int_X \liminf_{n \rightarrow \infty} f_n d\mu = 0$$

$$\int_X f_n d\mu = \begin{cases} \mu(E), & n \text{ is odd} \\ \mu(X \setminus E), & n \text{ is even} \end{cases}$$

$$\liminf_{n \rightarrow \infty} \int_X f_n d\mu = \min \{ \mu(E), \mu(X \setminus E) \}$$

□

Problem 2.7.8

Suppose μ is a positive measure on X , $f : X \rightarrow [0, \infty]$ is measurable, $\int_X f d\mu = c$, where $0 < c < \infty$, and α is a constant. Prove that

$$\lim_{n \rightarrow \infty} \int_X n \log[1 + (f/n)^\alpha] d\mu = \begin{cases} \infty & \text{if } 0 < \alpha < 1, \\ c & \text{if } \alpha = 1, \\ 0 & \text{if } 1 < \alpha < \infty. \end{cases}$$

Hint: If $\alpha \geq 1$, the integrands are dominated by αf . If $\alpha < 1$, Fatou's lemma can be applied.

Proof. By differentiating $\frac{1}{y} \log(1 + ky)$. We know $n \log\left(1 + \frac{f}{n}\right)$ is increasing as n is increasing. By Monotone Convergence Theorem,

$$\lim_{n \rightarrow \infty} \int_X n \log\left[1 + \left(\frac{f}{n}\right)\right] d\mu = \int_X \lim_{n \rightarrow \infty} n \log\left[1 + \frac{f}{n}\right] d\mu = \int_X f d\mu = c$$

. If $\alpha > 1$, then note that

$$0 \leq n \log\left[1 + \left(\frac{f}{n}\right)^\alpha\right] \leq \alpha f, \quad \int_X \alpha f d\mu = \alpha c < \infty$$

The Dominated Convergence Theorem gives that

$$\lim_{n \rightarrow \infty} \int_X n \log\left[1 + \left(\frac{f}{n}\right)^\alpha\right] d\mu = \int_X \lim_{n \rightarrow \infty} n \log\left[1 + \left(\frac{f}{n}\right)^\alpha\right] d\mu = 0$$

If $\alpha < 1$, the Fatou's Lemma gives that

$$\lim_{n \rightarrow \infty} \int_X n \log\left[1 + \left(\frac{f}{n}\right)^\alpha\right] d\mu \geq \int_X \lim_{n \rightarrow \infty} n \log\left[1 + \left(\frac{f}{n}\right)^\alpha\right] d\mu = \int_X \infty d\mu = \infty$$

□

Problem 2.7.9

Suppose $\mu(X) < \infty$, $\{f_n\}$ is a sequence of bounded complex measurable

functions on X , and $f_n \rightarrow f$ uniformly on X . Prove that

$$\lim_{n \rightarrow \infty} \int_X f_n d\mu = \int_X f d\mu,$$

and show that the hypothesis " $\mu(X) < \infty$ " cannot be omitted.

Proof. Take any $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} \int_X |f_n - f| d\mu \leq \mu(X) \varepsilon$$

Let $\varepsilon \rightarrow 0$, we have

$$\lim_{n \rightarrow \infty} \int_X |f_n - f| d\mu = 0$$

Thus

$$\left| \lim_{n \rightarrow \infty} \int_X f_n d\mu - \int_X f d\mu \right| \leq \lim_{n \rightarrow \infty} \int_X |f_n - f| d\mu = 0$$

Let $f_n = \chi_{[0,n]} \frac{1}{n}$. Then $f = 0$,

$$\lim_{n \rightarrow \infty} \int_X f_n d\mu = 1$$

$$\int_X f d\mu = 0$$

Thus $\mu(X) < \infty$ cannot be omitted. □

Problem 2.7.10

Let $\{E_k\}$ be a sequence of measurable sets in X , such that

$$\sum_{k=1}^{\infty} \mu(E_k) < \infty.$$

Suppose A is the set of all x which lie in infinitely many E_k . Show that

$$A = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} E_k$$

Then almost all $x \in X$ lie in at most finitely many of the sets E_k .

Proof. $x \in A^c$ iff there exists $k \in \mathbb{N}$, such that for all $n \geq k$, $x \in E_k^c$. That is

$$A^c = \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} E_k^c$$

Thus

$$A = \bigcap_{k=1}^{\infty} \bigcup_{n=k}^{\infty} E_k$$

$$\mu(A) = \lim_{k \rightarrow \infty} \mu\left(\bigcup_{n=k}^{\infty} E_k\right) \leq \lim_{k \rightarrow \infty} \sum_{n=k}^{\infty} \mu(E_k) = \left(\sum_{k=1}^{\infty} \mu(E_k) - \lim_{k \rightarrow \infty} \sum_{m=1}^k \mu(E_k)\right) = 0$$

□

Problem 2.7.11

Suppose $f \in L^1(\mu)$. Prove that to each $\epsilon > 0$ there exists a $\delta > 0$ such that $\int_E |f| d\mu < \epsilon$ whenever $\mu(E) < \delta$.

Proof. Let

$$f_n := \min\{|f|, n\}$$

Then

$$0 \leq f_1 \leq f_2 \leq \dots, \quad f_n \rightarrow |f| \text{ as } n \rightarrow \infty$$

$$\int_E |f| d\mu \leq \lim_{n \rightarrow \infty} \int_X f_n d\mu$$

There exists $N \in \mathbb{N}$, such that

$$\int_E |f| d\mu \leq \int_E f_N d\mu + \frac{\epsilon}{2} \leq \mu(E)N + \frac{\epsilon}{2} \leq \epsilon$$

whenever $\mu(E) < \delta := \frac{\varepsilon}{2N}$

□

Problem 2.7.12

If $f \geq 0$, then

$$\int_E \infty \cdot f \, d\mu = \infty \cdot \int_E f \, d\mu.$$

Proof. Let $E_1 = f^{-1}(0) \cap E$, $E_2 = E \setminus E_1$.

1. If $\mu(E_2) > 0$ Then

$$\int_E \infty \cdot f \, d\mu \geq \int_{E_2} \infty \cdot f \, d\mu = \infty$$

$$\int_E f \, d\mu \geq \int_{E_2} f \, d\mu > 0, \quad \infty \cdot \int_E f \, d\mu = \infty$$

2. If $\mu(E_2) = 0$, then

$$\int_E \infty \cdot f \, d\mu = \int_{E_1} \infty \cdot 0 \, d\mu + \int_{E_2} \infty \cdot f \, d\mu = 0$$

$$\int_E f \, d\mu = \int_{E_1} 0 \, d\mu + \int_{E_2} f \, d\mu = 0, \quad \infty \cdot \int_E f \, d\mu = 0$$

□

Measure Theory

3.1 Outer Measure

Definition 3.1.1 ► Outer Measure

A function φ defined for every subset A of an arbitrary set X is called an **outer measure** on X if the following conditions are satisfied:

- (i) $\varphi(\emptyset) = 0$,
- (ii) $0 \leq \varphi(A) \leq \infty$ whenever $A \subset X$,
- (iii) $\varphi(A_1) \leq \varphi(A_2)$ whenever $A_1 \subset A_2$,
- (iv) $\varphi(\bigcup_{i=1}^{\infty} A_i) \leq \sum_{i=1}^{\infty} \varphi(A_i)$ for every countable collection of sets $\{A_i\}$ in X .

Example 3.1.2 ► Outer Measure

- (i) In an arbitrary set X , define $\varphi(A) = 1$ if A is nonempty and $\varphi(\emptyset) = 0$.
- (ii) Let $\varphi(A)$ be the number (possibly infinite) of points in A .
- (iii) Let $\varphi(A) = \begin{cases} 0 & \text{if card } A \leq \aleph_0, \\ 1 & \text{if card } A > \aleph_0. \end{cases}$
- (iv) If X is a metric space, fix $\varepsilon > 0$. Let $\varphi(A)$ be the smallest number of balls of radius ε that cover A .
- (v) Select a fixed x_0 in an arbitrary set X , and let

$$\varphi(A) = \begin{cases} 0 & \text{if } x_0 \notin A, \\ 1 & \text{if } x_0 \in A; \end{cases}$$

φ is called the **Dirac measure** concentrated at x_0 .

Definition 3.1.3

Let φ be an outer measure on a set X . A set $E \subset X$ is called **φ -measurable** if

$$\varphi(A) = \varphi(A \cap E) + \varphi(A - E)$$

for every set $A \subset X$. In view of property (iv) above, observe that φ -measurability requires only

$$\varphi(A) \geq \varphi(A \cap E) + \varphi(A - E).$$

Lemma 3.1.4

A set $E \subset X$ is φ -measurable if and only if

$$\varphi(P \cup Q) = \varphi(P) + \varphi(Q)$$

for all sets P and Q such that $P \subset E$ and $Q \subset E^c$.

Proof. For Necessity, if E is φ -measurable, then for all such P, Q ,

$$\varphi(P \cup Q) = \varphi((P \cup Q) \cap E) + \varphi((P \cup Q) - E) = \varphi(P) + \varphi(Q)$$

For sufficiency, let $P = A \cap E$, $Q = A \cap E^c = A \setminus E$. Then

$$\varphi(A) = \varphi(P \cup Q) = \varphi(P) + \varphi(Q) = \varphi(A \cap E) + \varphi(A \setminus E)$$

□

Theorem 3.1.5

Suppose φ is an outer measure on an arbitrary set X . Then the following four statements hold:

- (i) E is φ -measurable whenever $\varphi(E) = 0$.
- (ii) $\emptyset$ and X are φ -measurable.
- (iii) $E_1 - E_2$ is φ -measurable whenever E_1 and E_2 are φ -measurable.
- (iv) If $\{E_i\}$ is a **countable collection of disjoint φ -measurable sets**, then $\bigcup_{i=1}^{\infty} E_i$ is φ -measurable and

$$\varphi\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} \varphi(E_i).$$

More generally, if $A \subset X$ is an arbitrary set, then

$$\varphi(A) = \sum_{i=1}^{\infty} \varphi(A \cap E_i) + \varphi(A \cap \tilde{S}),$$

where

$$S = \bigcup_{i=1}^{\infty} E_i.$$

Proof. 1. Take any $A \subseteq X$. Since $A \cap E \subseteq E$, $\varphi(A \cap E) = 0$. Thus

$$\varphi(A \cap E) + \varphi(A \cap E^c) = \varphi(A \cap E^c) \leq \varphi(A)$$

2. Follows immediately from lemma 3.1.4.

3. Take any P, Q such that $P \subseteq (E_1 \setminus E_2) = E_1 \cap E_2^c$, $Q \subseteq (E_1 \setminus E_2)^c = E_1^c \cup E_2$.

$$\varphi(Q) = \varphi(Q \cap E_2) + \varphi(Q - E_2)$$

Note that $Q - E_2 \subseteq E_1^c$, $Q \cap E_2 \subseteq E_2$. Since $P \subseteq E_1$, the measurability of E_1 implies that

$$\varphi(P) + \varphi(Q - E_2) = \varphi(P \cup (Q - E_2))$$

Note that $P \cup (Q - E_2) \subseteq E_2^c$. Since $Q \cap E_2 \subseteq E_2$, we have

$$\varphi(P \cup (Q - E_2)) + \varphi(Q \cap E_2) = \varphi(P \cup Q)$$

Finally, we have

$$\begin{aligned} \varphi(P) + \varphi(Q) &= \varphi(P) + \varphi(Q \cap E_2) + \varphi(Q - E_2) \\ &= \varphi(P \cup (Q - E_2)) + \varphi(Q \cap E_2) \\ &= \varphi(P \cup Q) \end{aligned}$$

4. Let $S_k = \bigcup_{i=1}^k E_i$. We proceed by finite induction and first note that the result is obviously true for $k = 1$. For $k > 1$ assume that S_k is φ -measurable and that

$$\varphi(A) \geq \sum_{i=1}^k \varphi(A \cap E_i) + \varphi(A \cap S_k^c), \quad \forall A \subseteq X$$

Then

$$\begin{aligned}\varphi(A) &= \varphi(A \cap E_{k+1}) + \varphi(A \cap E_{k+1}^c) \\ &= \varphi(A \cap E_{k+1}) + \varphi(A \cap E_{k+1}^c \cap S_k^c) + \varphi(A \cap E_{k+1}^c \cap S_k) \\ &= \varphi(A \cap E_{k+1}) + \varphi(A \cap S_{k+1}^c) + \varphi(A \cap S_k)\end{aligned}$$

Replaced A by $A \cap S_k$, we have

$$\varphi(A \cap S_k) \geq \sum_{i=1}^k \varphi(A \cap S_k \cap E_i) + \varphi(A \cap S_k \cap S_k^c) = \sum_{i=1}^k \varphi(A \cap E_i)$$

Thus

$$\varphi(A) \geq \sum_{i=1}^{k+1} \varphi(A \cap E_{k+1}) + \varphi(A \cap S_{k+1}^c)$$

Then the subadditivity gives that

$$\varphi(A) \geq \varphi(A \cap S_{k+1}) + \varphi(A \cap S_{k+1}^c)$$

this, in turn, implies that S_{k+1} is φ -measurable. Since we now know for every set $A \subseteq X$ and for all positive integers k

$$\varphi(A) \geq \sum_{i=1}^k \varphi(A \cap E_i) + \varphi(A \cap S_k^c)$$

We have

$$\varphi(A) \geq \sum_{i=1}^{\infty} \varphi(A \cap E_i) + \varphi(A \cap S^c) \geq \varphi(A \cap S) + \varphi(A \cap S^c)$$

where $S = \bigcup_{i=1}^{\infty} E_k$. Thus S is measurable. The subadditivity gives that

$$\varphi(A) \leq \sum_{i=1}^{\infty} \varphi(A \cap E_i) + \varphi(A \cap S^c)$$

Finally, we have

$$\varphi(A) = \sum_{i=1}^{\infty} \varphi(A \cap E_i) + \varphi(A \cap S^c)$$

Replace A by S , we can get

$$\varphi(S) = \sum_{i=1}^{\infty} \varphi(E_i)$$

□

Index

Definitions

1.3.1	Limit Inferior and Limit Superior	3
1.3.3	Limit of a Set Sequence	3
2.1.1		6
2.1.5	Borel	8
2.2.1		9
2.3.1	Measure and Measure Space	11
2.4.1	Integration of Positive Functions	13
2.5.1	Lebesgue Integrability	20
2.5.2	Complex Integration	21
2.5.3	Real Integration	21
2.6.1		24
2.6.5	More Measurable Functions	26
3.1.1	Outer Measure	40
3.1.3		40

Examples

2.3.3		13
2.6.2		24
3.1.2	Outer Measure	40

Problems

2.7.1		30
2.7.2		31
2.7.3		31
2.7.4		32
2.7.5		33

2.7.6		34
2.7.7		35
2.7.8		36
2.7.9		36
2.7.10		37
2.7.11		38
2.7.12		39

Propositions

1.3.2		3
1.3.4		4
1.3.5		5
2.4.3		14
2.6.3	Completion of a Measure	25

Corollarys

2.1.3		7
2.1.8		9
2.4.2		14
2.4.5		16
2.4.6		17
2.4.7		17

Lemmas

1.2.1		2
2.1.2		6
2.4.8	Fatou's Lemma	18
3.1.4		41

Theorems

2.1.4	8
2.1.6	8
2.1.7	8
2.2.2	10
2.3.2	11
2.4.4	Lebesgue's Monotone Convergence Theorem 15
2.4.9	19

2.5.4	21
2.5.5	22
2.5.6	Lebesgue's Dominated Convergence Theorem 22
2.6.4	25
2.6.6	26
2.6.7	28
2.6.8	29
2.6.9	30
3.1.5	41