

Contents

Index	2
1 Basics	4
1.1 单位分解	4
1.2 向量场和微分算子	4
1.3 积分	8
1.4 映射与子流形	9
1.5 Frobenius 定理	14
2 Basic Concepts of Lie Algebra	16
2.1 Definitions and first examples	16
2.2 Lie algebras of derivations	17
2.3 Ideals and homomorphism	17
2.3.1 Ideals	17
2.3.2 Homomorphisms and representations	17
3 Solvable Lie Algebras	19
3.1 Solvable Lie Algebras	19
4 Subalgebra of the General Linear Algebra	22
4.1 Nilpotent Maps	22
4.2 Weights	23
4.3 Invariance Lemma	23
5 Engel' s Theorem and Lie' s Theorem	24
5.1 Engel's Theorem	24
5.2 Lie's Theorem	27
6 Representation of $\mathfrak{sl}(2, \mathbb{C})$	28
6.1 The Modules V_d	28
6.2 Classifying the Irreducible $\mathfrak{sl}(2, \mathbb{C})$ -Modules	28
6.3 Weyl's Theorem	29
7 Cartan Criteria	30
7.1 Jordan Decomposition	30
7.2 Testing for Solvability	31

7.3	Killing Form	34
7.3.1	Testing for Semisimplicity	35
7.4	Derivations of Semisimple Lie Algebras	37
7.5	Abstract Jordan Decomposition	38
8	The Root Space Decomposition	40
8.1	Preliminary Results	40
8.2	Cartan Subalgebras	43
8.3	Definition of the Root Space Decomposition	43
8.4	Root Strings and Eigenvalues	46
9	Low Dimension Lie Algebras	48
9.1	Dimension 3	48
9.1.1	Lie Algebras where $L' = L$	48

Index

Basics

1.1 单位分解

Theorem 1.1.1 ▶ 单位分解的存在性

设 M 是一个流形, $\mathcal{U} = \{U_\alpha : \alpha \in A\}$ 是 M 的一个开覆盖, 则

1. 存在从属于 $\mathcal{U}$ 的单位分解 $\Phi = \{\varphi_i : i \in I\}$ 且 I 是至多可数的, $\text{supp } \varphi_i$ 是紧的.
2. 存在从属于 $\mathcal{U}$ 的具有相同指标集的单位分解 $\Phi = \{\varphi_\alpha : \alpha \in A\}$, 且至多可数多个 $\varphi_\alpha \neq 0$.
3. 如果 $\mathcal{U}$ 局部有限, 且 $\overline{U_\alpha}$ 是紧的. 则存在从属于 $\mathcal{U}$ 的具有相同指标集的单位分解 $\Phi = \{\phi_\alpha : \alpha \in A\}$ 且 $\text{supp } \phi_\alpha$ 是紧的.^a

^a由于紧的闭子集是紧的, 所以第三条可以有第二条简单地导出.

Corollary 1.1.2

设 M 是一个流形, $D \subseteq M$ 是闭集, $U \subseteq M$ 是 D 的一个开邻域. 则存在 $f \in C^\infty(M)$, 使得 $f|_D \equiv 1$, 且 $\text{supp } f \subseteq U$.

Proof. 考虑 M 的一个特殊的开覆盖, $\{U, M \setminus D\}$. 对它应用上面定理的第二种情况即可. $\square$

1.2 向量场和微分算子

Definition 1.2.1

1. 设 M 是一个流形, $p \in M$. 若 $f, g \in C^\infty(M)$ 满足, 存在 p 的开邻域 U , 使得 $f|_U = g|_U$, 则称 f 和 g 在 p 附近等价.
2. 称 $C^\infty(M) / \sim$ 为 p 附近的函数芽, 记作 $C_p^\infty(U)$.
3. 对于 $f \in C_p^\infty(M)$, 它的芽记作 $[f] \in C_p^\infty(M)$

在欧式空间上, 光滑函数有展开式

$$f(x) = f(x_0) + \sum_{i=1}^m \frac{\partial f}{\partial x^i} (x - x_0)^i + \sum_{i,j=1}^m \frac{1}{2} \frac{\partial^2 f}{\partial x^i \partial x^j} (x_0) (x - x_0)^i (x - x_0)^j + O(\|x - x_0\|^2)$$

$$D = C(x) + \sum_{i=1}^m a_i(x) \frac{\partial}{\partial x^i} + \sum_{i,j=1}^m \frac{1}{2} b_{ij}(x) \frac{\partial^2}{\partial x^i \partial x^j}$$

Definition 1.2.2

令 $m_p = \{f \in C_p^\infty(M) : f(p) = 0\}$, 则 m_p 是 $C_p^\infty(M)$ 的一个极大理想. 特别地, m_p^k 也是 $C_p^\infty(M)$ 的理想.

则 f 的展开式的第二项和第三项分别视作 m_p 和 m_p^2 中的元素.

Definition 1.2.3 ▶ Differential Expression of Order r

1. 一个 r 阶微分表达式, 是指一个线性函数 $D : C_p^\infty(M) \rightarrow \mathbb{R}$ 满足 $D(m_p^{r+1}) = 0$.
2. 全体 p 处的 r 阶微分表达式记作 $T_p^{(r)}(M)$
3. 定义 $T_p^{(\infty)}(M) = \bigcup_{r \geq 0} T_p^{(r)}(M)$

在这个定义下, 当 $r = 2$ 时,

$$\begin{aligned} D(f(x)) &= D\left(f(x_0) + \sum \frac{\partial f}{\partial x^i} (x - x_0)^i + \sum \frac{1}{2} \frac{\partial^2 f}{\partial x^i \partial x^j} (x - x_0)^i (x - x_0)^j\right) \\ &= D(f(x_0)) + \sum_i \frac{\partial f}{\partial x^i} (x_0) D((x - x_0)^i) + \sum_{i,j} \frac{1}{2} \frac{\partial^2 f}{\partial x^i \partial x^j} (x_0) D((x - x_0)^i (x - x_0)^j) \\ &=: cf(x_0) + \sum_i a_i \frac{\partial f}{\partial x^i} (x_0) + \sum_{i,j} \frac{1}{2} b_{ij} \frac{\partial^2 f}{\partial x^i \partial x^j} (x_0) \end{aligned}$$

其中 $c, a_i, b_{i,j}$ 均为常数.

Definition 1.2.4 ▶ 切向量

1. 对于一个 1 阶微分表达式 X , 若它满足

$$X(f \cdot g) = X(f) \cdot g(x_0) + f(x_0)X(g)$$

则称 X 为 p 处的一个切向量.

具体地,

$$X(f) = \sum_{i=1}^m a_i \frac{\partial f}{\partial x^i}(x_0)$$

2. 记 $T_p(M)$ 为 p 处全体切向量构成的线性空间. 则 $T_p(M) \simeq \mathbb{R}^m$. $\partial_i := \frac{\partial}{\partial x^i}$ 构成 $T_p(M)$ 的基.

让 $p \in U$ 在 U 上可变 $\iff x \in V$ 跑起来. 则 $X = \sum a^i(x) \frac{\partial}{\partial x^i}$, $a^i \in C^\infty(U)$.

Definition 1.2.5 ▶ 多重指标

- 一个多重指标 α 是指 $\alpha = (\alpha_1, \dots, \alpha_m)$, $\alpha_i \in \mathbb{N}$. 定义 α 的长度为 $|\alpha| = \alpha_1 + \dots + \alpha_m$
- 定义记号

$$\partial^\alpha := \left(\frac{\partial}{\partial x^1} \right)^{\alpha_1} \cdots \left(\frac{\partial}{\partial x^m} \right)^{\alpha_m}$$

Proposition 1.2.6

$T_p^{(r)}(M)$ 的一组基为 $\{\partial^\alpha : |\alpha| \leq r\}$

Definition 1.2.7

一个线性映射 $\mathcal{D} : C^\infty(M) \rightarrow C^\infty(M)$ 是一个阶数至多为 r 的微分算子, 如果对于流形 M 上的任意一个局部坐标图 $(U, x^1, \dots, x^m)$, 算子在 U 上的限制 $\mathcal{D}|_U$ 都可以表示为:

$$\mathcal{D}|_U = \sum_{|\alpha| \leq r} a_\alpha(x) \partial^\alpha$$

其中:

- $\alpha = (\alpha_1, \dots, \alpha_m)$ 是多重指标.
- $\partial^\alpha = \frac{\partial^{|\alpha|}}{(\partial x^1)^{\alpha_1} \cdots (\partial x^m)^{\alpha_m}}$ 是偏微分.

- 每一个系数 $a_\alpha(x)$ 都是定义在 U 上的光滑函数, 即 $a_\alpha \in C^\infty(U)$.

Definition 1.2.8

- 开集 U 上的切向量的全体记作 $\mathfrak{X}(U)$.
- 开集 U 上的微分算子的全体记作 $D(U)$

Remark.

$D(U)$ 上按算子的复合, 可以定义出自然的乘法. 它是结合但不交换的.

Definition 1.2.9

$\mathfrak{X}(U)$ 上有 Lie 括号

$$\begin{aligned} [\cdot, \cdot] : \mathfrak{X}(U) \times \mathfrak{X}(U) &\rightarrow \mathfrak{X}(U) \\ [X, Y] &:= XY - YX \end{aligned}$$

Proposition 1.2.10

- $C^\infty(U)$ 是交换结合的 $\mathbb{R}$ -代数.
- $\mathfrak{X}(U)$ 和 $D(U)$ 都是 $C^\infty(U)$ 模

Proposition 1.2.11

若 D_1 是 r_1 阶的微分算子, D_2 是 r_2 阶的微分算子, 则

- $D_1 \cdot D_2$ 的阶数 $\leq r_1 + r_2$.
- $D_1 D_2 - D_2 D_1$ 的阶数 $\leq r_1 + r_2 - 1$

Theorem 1.2.12

设 M 是一个流形, $\dim M = m$. $X_1, \dots, X_m \in \mathfrak{X}(M)$ 满足对于任意的 $p \in M$. $X_1(p), X_2(p), \dots, X_m(p)$ 构成 $T_p(M)$ 的一组基^a. 记 $X^\alpha = X_1^{\alpha_1} \cdots X_m^{\alpha_m}$. 则

- $\forall p \in M, \{X_{(p)}^\alpha : |\alpha| \leq r\}$ 构成 $T_p^{(r)}(M)$ 的基.
- 等价地说, $\{X^\alpha : \alpha = (\alpha_1, \dots, \alpha_m)\}$ 构成 $D(M)$ 的作为 $C^\infty(M)$ 模的基.
- 此时, D 可以写作

$$D = \sum a_\alpha X^\alpha, \quad a_\alpha \in C^\infty(M)$$

^a即 M 是一个可平行化的流形

1.3 积分

设 M 是流形, $\dim M = m$, ω 是 M 上的一个 m -形式. $f \in C^\infty(M)$.

1. 取 M 的一个图册 $\mathcal{U} = \{(U_\alpha, \varphi_\alpha) : \alpha \in A\}$.
2. 取从属于 $\mathcal{U}$ 的一个紧支单位分解. $\Phi = \{\varphi_i : i \in I\}$
3. 则 $f = \sum_{i \in I} f \cdot \varphi_i$. 记 $f_i = f \varphi_i$
4. 设 $\text{supp } \varphi_i \subseteq U_\alpha$.
5. 设 V_α 中的坐标为 $(x^1, \dots, x^n)$, $f_i : U_\alpha \rightarrow \mathbb{R}$. 定义 $\tilde{f}_i = f_i \circ \varphi_\alpha^{-1} : V_\alpha \rightarrow \mathbb{R}$.
6. 设 ω 在 U_α 上有表达式 $\omega = \omega_\alpha(x) dx^1 \wedge \dots \wedge dx^m$.

Definition 1.3.1 ▶ 对密度的积分

定义

$$J_i := \int_{U_\alpha} \tilde{f}_i(x) |\omega_\alpha(x)| dx^1 \dots dx^n$$

定义

$$J := \int_M f |\omega| := \sum_{i \in I} J_i$$

称为是对密度的积分.

Remark.

需要证明 J 与紧支单位分解的选取无关.

Definition 1.3.2 ▶ 可定向性

对于流形 M . 如果存在一个图册 $\{U_\alpha : \alpha \in A\}$ 使得任意两个坐标卡之间的转移函数的 Jacobi 行列式都是正的. 则称 M 是可定向的.

- 称这样的图册为定向图册
- 若 $\mathcal{U}_1$ 和 $\mathcal{U}_2$ 是两个定向图册, 满足 $\mathcal{U}_1 \cup \mathcal{U}_2$ 还是定向图册, 则称 $\mathcal{U}_1$ 和 $\mathcal{U}_2$ 是定向相容的.

Definition 1.3.3 ▶ 对形式的积分

若 M 还是可定向的, 可以定义

$$J_i = \int_{U_\alpha} \tilde{f}_i(x) \omega_\alpha(x) dx^1 \cdots dx^n$$

定义

$$J := \int_M f \omega := \sum_{i \in I} J_i$$

Remark.

可以证明, J 不依赖于与 $\mathcal{U}$ 定向相同的图册的选取.

Definition 1.3.4

M 是可定向的, 当且仅当 M 上存在一个处处非零的 m -形式.

Theorem 1.3.5

设 M 是可定向的带边流形, ω 是一个紧支的 $(m-1)$ 形式, 则

$$\int_M d\omega = \int_{\partial M} \omega$$

1.4 映射与子流形

Lemma 1.4.1

设 M, N 是 C^∞ 流形, $\varphi : M \rightarrow N$ 连续. 若对于任意的 N 的开集 V , 和 $g \in C^\infty(V)$. $g \circ \varphi \in C^\infty(\varphi^{-1}(V))$. 则 φ 是 C^∞ 的.

Lemma 1.4.2

对于 $y \in N$ 附近的两个 C^∞ 函数 g_1, g_2 , 若 $g_1 \sim_y g_2$ (函数芽). 则 $g_1 \circ \varphi \sim_x g_2 \circ \varphi$.

Definition 1.4.3 ▶ 拉回映射

M, N, φ 同前. 则 φ 诱导出线性映射

$$\begin{aligned}\varphi^* : C_y^\infty(N) &\rightarrow C_x^\infty(M) \\ g &\mapsto g \circ \varphi\end{aligned}$$

称为是关于 φ 的拉回映射

Definition 1.4.4 ▶ 切映射

M, N, φ 同前. 则 φ 诱导出对偶空间 (微分算子空间) 之间的映射

$$(\mathrm{d}\varphi)_* : C_x^\infty(M)^* \rightarrow C_{\varphi(x)}^\infty(N)^*$$

通过做限制得到映射

1.

$$(\mathrm{d}\varphi)_x : T_x M \rightarrow T_{\varphi(x)} N$$

2.

$$(\mathrm{d}\varphi)_x^{(\infty)} : C_x^\infty(M)^* \rightarrow C_{\varphi(x)}^\infty(N)^*$$

Remark.

这里蕴含了一些事实

1. 如果 X 是切向量, 则 $(\mathrm{d}\varphi)_x(X)$ 也是.
2. $\varphi^{-1*}(m_y) \subseteq m_x$, 从而 $\varphi^{-1*}(m_y^k) \subseteq m_x^k$.
3. 若 D 是一个 r 阶微分表达式, $(\mathrm{d}\varphi)_x(D)$ 也是 r 阶微分表达式.

Definition 1.4.5

M, N, φ 同前. 若 X 是 M 上的向量场. 若 $X \in \mathfrak{X}(M), Y \in \mathfrak{X}(N)$ 满足

$$\forall_{x \in M} (\mathrm{d}\varphi)_x(X(x)) = Y(\varphi(x))$$

则称 X 和 Y 是 φ -相关的.

Lemma 1.4.6

设 D 是 M 上的微分算子. $D : C^\infty(M) \rightarrow C^\infty(M)$. 若 $\tilde{D} \in D(C^\infty(N))$. 则 $\tilde{D} : C^\infty(N) \rightarrow C^\infty(N)$

Definition 1.4.7

$\varphi : M \rightarrow N$ 诱导出映射

$$\varphi^* : \Omega^*(N) \rightarrow \Omega^*(M)$$

Definition 1.4.8

设 M, N, φ 同前. $\dim M = m, \dim N = n. r_x := \text{rank}(\text{d}\varphi)_x$.

1. 若对于任意的 $x \in M, r_x = n$. 则称 φ 是一个淹没 (Submersion).
2. 若对于任意的 $x \in M, r_x = m$. 则称 φ 是一个浸入 (immersion).

Proposition 1.4.9

1. 若 φ 是一个淹没, 则 $m \geq n$. 且存在局部坐标系, 使得

$$\tilde{\varphi}(x^1, \dots, x^m) = (x^1, \dots, x^n)$$

2. 若 φ 是一个浸入, 则 $m \leq n$. 且存在局部坐标, 使得

$$\tilde{\varphi}(x^1, \dots, x^m) = (x^1, \dots, x^n, 0, \dots, 0)$$

Definition 1.4.10

设 M, N 是拓扑流形. $\varphi : M \rightarrow N$ 是连续映射.

1. 若对于任意的 $x \in M$. 记 $y = \varphi(x)$. 存在 x, y 附近的坐标卡 $(U, \varphi_U : U \rightarrow \tilde{U}), (V, \varphi_V : V \rightarrow \tilde{V})$. 使得

$$\tilde{\varphi} = \varphi_V \circ \varphi \circ \varphi_U^{-1} : \tilde{U} \rightarrow \tilde{V}$$

是一个从 $\mathbb{R}^m$ 到 $\mathbb{R}^n$ 的投影. 则称 φ 是一个拓扑淹没.

2. 可以类似地定义出拓扑浸入.

Theorem 1.4.11

设 M 是 C^∞ 流形. N 是拓扑流形. 若 $\varphi : M \rightarrow N$ 是拓扑的淹没且满, 则存在唯一的 C^∞ 结构, 使得 φ 是一个 C^∞ 淹没.

Definition 1.4.12

设 $\varphi : M \rightarrow N$ 是一个浸入.

1. 若 φ 是单的, 则称 φ 是一个嵌入.

2. 若 φ 是嵌入, 且 $\varphi_M : M \rightarrow \varphi(M)$ 是同胚. 则称 φ 是一个正则嵌入.

Theorem 1.4.13 ▶ 正则嵌入的水平集刻画

设 $\varphi : M \rightarrow N$ 是正则嵌入. 则对于 $y \in \varphi(M)$. 存在 y 在 N 中的开邻域 V , 以及 V 上的 C^∞ 映射 $F : V \rightarrow \mathbb{R}^{n-m}$. 使得 $\varphi(M) \cap V = \{z \in V : F(z) = 0\}$

Theorem 1.4.14 ▶ 正则嵌入的参数刻画

对于 $x \in M, y = \varphi(x) \in \varphi(M)$. 存在 x 在 M 中的开邻域 U, y 在 N 中的开邻域 V . 使得 $\varphi(U) = \varphi(M) \cap V$.

Theorem 1.4.15

设 P 是任一 C^∞ 流形. $\varphi : M \rightarrow N$ 是正则嵌入. $u : P \rightarrow M$ 是任一映射. 则

$$u \text{ 是 } C^\infty \text{ 的} \iff \varphi \circ u : P \rightarrow N \text{ 是 } C^\infty \text{ 的.}$$

Definition 1.4.16

若 $\varphi : M \rightarrow N$ 是嵌入且满足

$$u \text{ 是 } C^\infty \text{ 的} \iff \varphi \circ u : P \rightarrow N \text{ 是 } C^\infty \text{ 的.}$$

则称 φ 是拟正则嵌入.

Definition 1.4.17

设 N 是一个 C^∞ 流形. $M \subseteq N$. 若 $i : M \rightarrow N$ 是嵌入. 则称 M 是 N 的子流形. 若 i 还是正则或拟正则的. 则称 M 是 N 的正则或拟正则子流形.

Theorem 1.4.18 ▶ 正则值

设 M, N 是 C^∞ 流形. $\varphi : M \rightarrow N$ 是 C^∞ 映射. 对于 $y \in N$. 记 $M' = \varphi^{-1}(y)$. 若对任一的 $x \in M'$. $\text{rank}(\text{d}\varphi)_x = n$. 则 M' 是 M 的一个正则子流形, 维数为 $m - n$. 这样的 y 叫 φ 的一个正则值.

Note.

- $m - n$ 衡量的是 M 上有多少维数的信息, 在通过 φ 映射的过程中被压缩掉了.
- y 的正则性, 体现在它的“来源信息”在 M 上是光滑的.
- 正则值 y 给出的正则子流形 M' , 就是 M 上那些在映射 φ 下被压缩成 y 的信息构成的子流形.

Example 1.4.19

1. $\text{Mat}(n, \mathbb{R}) \simeq \mathbb{R}^{n^2}$ 是 n^2 维流形.
2. $\text{GL}(n, \mathbb{R}) = \{M \in \text{Mat}(n, \mathbb{R}) : \det M \neq 0\}$ 作为 $\text{Mat}(n, \mathbb{R})$ 的一个开子集也是一个 n^2 维流形.
3. $\text{SL}(n, \mathbb{R}) = \{M \in \text{Mat}(n, \mathbb{R}) : \det M = 1\}$

$$\begin{aligned} \det : \text{Mat}(n, \mathbb{R}) &\rightarrow \mathbb{R} \\ M &\mapsto \det M \end{aligned}$$

$\text{SL}(n, \mathbb{R}) = \det^{-1}(1)$. 取一个 $M \in \text{Mat}(n, \mathbb{R})$, $\det M = 1$. 要证明 $(d(\det))_M : \mathbb{R}^{n^2} \rightarrow \mathbb{R}$. 按分量求偏导, 得到

$$\frac{\partial}{\partial a_{ij}} (\det M) = A_{ij} (\text{代数余子式})$$

由于 $\det M = 1$. 至少有一个 $A_{ij} \neq 0$. 从而 $(d(\det))_M$ 是满的.

4. 定义 $P : \text{Mat}(n, \mathbb{R}) \rightarrow \text{SMat}(n, \mathbb{R})$ (对称矩阵) $\simeq \mathbb{R}^{\frac{n(n+1)}{2}}$.

$$P(A) := A^T A$$

则 $O(n, \mathbb{R}) = P^{-1}(0)$. 对于 $A \in O(n, \mathbb{R})$.

$$(dP)_A : \text{Mat}(n, \mathbb{R}) \rightarrow \text{SMat}(n, \mathbb{R})$$

与上例不同, 这里用分量的形式去计算 Jacobi 是看不清楚的, 不过由于 P 本身足够简单, 我们可以用微分最原始的含义来观察. 考虑

$$P(A + \varepsilon B) - P(A) = \varepsilon (dP)_A(B) + O(\dots)$$

带入得到

$$(A^T + \varepsilon B^T)(A + \varepsilon B) - A^T A = \varepsilon (B^T A + A^T B) + O(\varepsilon)$$

于是 $(dP)_A(B) = B^T A + A^T B$. 只需要证明这个关于 B 的映射是满的. 为此, 解方程

$$A^T B + B^T A = C$$

取 $B = \frac{1}{2}AC$, 利用 A, C 对称, 带入得到

$$\frac{1}{2}A^T AC + \frac{1}{2}C^T A^T A = \frac{1}{2}(C + C^T) = C$$

故 $O(n, \mathbb{R})$ 是 $\text{Mat}(n, \mathbb{R})$ 的一个 $\frac{n(n-1)}{2}$ 维正则子流形. 注意到对于 $A \in O(n, \mathbb{R})$, $\det A = 1$ 或 $\det A = -1$, 可以窥见事实上 $O(n, \mathbb{R})$ 有两个连通分支.

1.5 Frobenius 定理

Definition 1.5.1 $\triangleright C^\infty$ 分布

在光滑流形 M 上, 一个 k -维 C^∞ 分布 (smooth distribution) 是一个映射 $\Delta : M \rightarrow \bigcup_{p \in M} \text{Gr}(k, T_p M)$, 这里

$$\text{Gr}(k, V) = \{W \subseteq V \mid W \text{ 是 } k\text{-维子空间}\}$$

其中:

- 对于每个 $p \in M$, Δ_p 是切空间 $T_p M$ 的一个 k -维子空间.
- Δ 是光滑的, 即它对应于切丛 TM 的一个光滑子向量丛: 局部存在光滑向量场 $X_1, \dots, X_k$ 生成 $\Delta_p = \text{span}\{X_1(p), \dots, X_k(p)\}$.

Definition 1.5.2

设 $\mathcal{L}$ 是 N 上的一个分布.

- 设 $\mathcal{L}$ 是一个 m 维分布. M 是 N 的一个 m 维子流形. 若对于 $x \in M$, $T_x M = L_x$, 则称 M 是 $\mathcal{L}$ 的一个积分子流形.
- 若对于任意的 $x \in N$. 都存在一个过 x 的 $\mathcal{L}$ 的积分子流形. 则称 $\mathcal{L}$ 是可积的.
- 若 $[\mathcal{L}, \mathcal{L}] \subseteq \mathcal{L}$, 则称 $\mathcal{L}$ 是对合的 (involutive).

Remark.

由于 Lie bracket 是封闭的, 从而可积一定是对合的.

Theorem 1.5.3 ▶ 局部 Frobenius

设 $\mathcal{L}$ 是 N 上的一个分布. 若 $\mathcal{L}$ 是对合的, 则 $\mathcal{L}$ 是可积的.

Theorem 1.5.4 ▶ 整体 Frobenius

设 N 是一个 n 维 C^∞ 流形. $\mathcal{L}$ 是 N 上的一个 m 维对合分布. 则过 N 上任一点 p , 存在唯一的 $\mathcal{L}$ 的极大积分子流形 M_p . $\mathcal{L}$ 的任何积分子流形都是 N 的拟正则子流形, 并且是某一个极大积分子流形的开子流形.

Proof sketch.

对 $p \in N$,

$$M_p = \{q \in N : \text{存在一条分片光滑的 } \mathcal{L} \text{ 的积分曲线, 连接 } p \text{ 和 } q\}$$

参考 Warner 的 GTM94

Basic Concepts of Lie Algebra

2.1 Definitions and first examples

Definition 2.1.1

A vector space L over a field $\mathbb{F}$, with an operation $L \times L \rightarrow L$, denoted $(x, y) \mapsto [xy]$ and called the **bracket** or **commutator** of x and y , is called a **Lie algebra** over $\mathbb{F}$ if the following axioms are satisfied:

- (L1) The bracket operation is bilinear.
- (L2) $[xx] = 0$ for all $x \in L$.
- (L3) $[x[yz]] + [y[zx]] + [z[xy]] = 0 \quad (x, y, z \in L)$.

Remark.

- (L1) and (L2), applied to $[x + y, x + y]$ imply anticommutativity :
(L2') $[xy] = [-yx]$.
- Conversely, if $\text{char } \mathbb{F} \neq 2$, it is clear that (L2') will imply (L2).

Definition 2.1.2

We define a **Lie subalgebra** of L to be a vector subspace $K \subseteq L$ such that

$$[x, y] \in K \quad \text{for all } x, y \in K.$$

Example 2.1.3

Suppose that V is a finite-dimensional vector space over F . Write $\mathfrak{gl}(V)$ for the set of all linear maps from V to V . This is again a vector space over F , and it becomes a Lie algebra, known as the **general linear algebra**, if we define the Lie bracket $[-, -]$ by

$$[x, y] := x \circ y - y \circ x \quad \text{for } x, y \in \mathfrak{gl}(V),$$

where $\circ$ denotes the composition of maps.

2.2 Lie algebras of derivations

Definition 2.2.1

- By an ***F*-algebra** (not necessarily associative) we simply mean a vector space $\mathfrak{A}$ over F endowed with a bilinear operation $\mathfrak{A} \times \mathfrak{A} \rightarrow \mathfrak{A}$, usually denoted by juxtaposition (unless $\mathfrak{A}$ is a Lie algebra, in which case we always use the bracket).
- By a ***derivation*** of $\mathfrak{A}$ we mean a linear map $\delta : \mathfrak{A} \rightarrow \mathfrak{A}$ satisfying the familiar product rule $\delta(ab) = a\delta(b) + \delta(a)b$ for each $a, b \in \mathfrak{A}$.

2.3 Ideals and homomorphism

2.3.1 Ideals

Definition 2.3.1

A subspace I of a Lie algebra L is called an ***ideal*** of L if $x \in L, y \in I$ together imply $[xy] \in I$.

Remark.

由于 $[x, y] = -[y, x]$, 理想不分左右.

2.3.2 Homomorphisms and representations

Definition 2.3.2

- A linear transformation $\phi : L \rightarrow L'$ (L, L' Lie algebras over F) is called a ***homomorphism*** if $\phi([xy]) = [\phi(x)\phi(y)]$, for all $x, y \in L$.
- ϕ is called a ***monomorphism*** if $\text{Ker}\phi = 0$, an ***epimorphism*** if $\text{Im}\phi = L'$, an ***isomorphism*** if it is both mono- and epi-.

Definition 2.3.3

If L_1 and L_2 are Lie algebras over a field F , then we say that a map $\varphi : L_1 \rightarrow L_2$ is a ***homomorphism*** if φ is a linear map and

$$\varphi([x, y]) = [\varphi(x), \varphi(y)] \quad \text{for all } x, y \in L_1.$$

Notice that in this equation the first Lie bracket is taken in L_1 and the

second Lie bracket is taken in L_2 . We say that φ is an *isomorphism* if φ is also bijective.

Example 2.3.4

An extremely important homomorphism is the *adjoint homomorphism*. If L is a Lie algebra, we define

$$\text{ad} : L \rightarrow \mathfrak{gl}(L)$$

by $(\text{ad } x)(y) := [x, y]$ for $x, y \in L$. It follows from the bilinearity of the Lie bracket that the map $\text{ad } x$ is linear for each $x \in L$. For the same reason, the map $x \mapsto \text{ad } x$ is itself linear. So to show that ad is a homomorphism, all we need to check is that

$$\text{ad}([x, y]) = \text{ad } x \circ \text{ad } y - \text{ad } y \circ \text{ad } x \quad \text{for all } x, y \in L;$$

this turns out to be equivalent to the Jacobi identity. The kernel of ad is the centre of L .

Proof. We have from Jacobi identity that

$$\begin{aligned} \text{ad}([x, y])(z) &= [[x, y], z] \\ &= [x, [y, z]] + [y, [z, x]] \\ &= [x, \text{ad } y(z)] - [y, \text{ad } x(z)] \\ &= \text{ad } x \circ \text{ad } y(z) - \text{ad } y \circ \text{ad } x(z) \end{aligned}$$

□

Solvable Lie Algebras

3.1 Solvable Lie Algebras

Lemma 3.1.1

Suppose that I is an ideal of L . Then L/I is abelian if and only if I contains the derived algebra L' .

Proof. L/I is abelian, if and only if for each $x, y \in L$,

$$[x + I, y + I] = [x, y] + I = 0 + I \iff [x, y] \in I$$

That is L/I is abelian, if and only if $L' = [L, L] \subseteq I$. □

Definition 3.1.2

This lemma tells us that the derived algebra L' is the smallest ideal of L with an abelian quotient. By the same argument, the derived algebra L' itself has a smallest ideal whose quotient is abelian, namely the derived algebra of L' , which we denote $L^{(2)}$, and so on. We define the **derived series** of L to be the series with terms

$$L^{(1)} = L' \quad \text{and} \quad L^{(k)} = [L^{(k-1)}, L^{(k-1)}] \text{ for } k \geq 2.$$

Then $L \supseteq L^{(1)} \supseteq L^{(2)} \supseteq \dots$. As the product of ideals is an ideal, $L^{(k)}$ is an ideal of L (and not just an ideal of $L^{(k-1)}$).

Definition 3.1.3

The Lie algebra L is said to be **solvable** if for some $m \geq 1$ we have $L^{(m)} = 0$.

Lemma 3.1.4

If L is a Lie algebra with ideals

$$L = I_0 \supseteq I_1 \supseteq \dots \supseteq I_{m-1} \supseteq I_m = 0$$

such that I_{k-1}/I_k is abelian for $1 \leq k \leq m$, then L is solvable.

Note.

- 我们把这样的理想链视作过滤掉非交换性的多层滤网, 每次都把可交换的元 (相当于粗颗粒) 保留, 将不能交换的程度 (细颗粒) 过滤下去.
- 导出列实现对不可交换性的精准过滤和采集.
- 普通的理想链是可能更粗的滤网, 它能承载所有上一层级的非交换性, 但是可能混入可交换性的细颗粒.
- 这个引理相当于是说, 如果我们通过可能更不精细的手法将所有的非交换性过滤干净, 那么原有的精确过滤也一定能过滤干净.

Proof. We shall inductively shows that $L^{(k)}$ is contained in I_k , then by taking $k = m$, we get $L^{(m)} = 0$.

Since $L/I_1 = I_0/I_1$ is abelian, $I_1 \supseteq L'$. From inductive hypothesis, we know that $L^{(k-1)} \subseteq I_{k-1}$, since I_{k-1}/I_k is abelian, that is

$$[I_{k-1}/I_k, I_{k-1}/I_k] = 0$$

which is equivalent to

$$[I_{k-1}, I_{k-1}] \subseteq I_k$$

Since $L^{(k-1)} \subseteq I_{k-1}$, we have

$$L^{(k)} = [L^{(k-1)}]$$

□

Proposition 3.1.5

Suppose that $\varphi : L_1 \rightarrow L_2$ is a surjective homomorphism of Lie algebras. Show that

$$\varphi(L_1^{(k)}) = (L_2)^{(k)}.$$

Proof. For $k = 0$,

$$\varphi(L_1) = L_2$$

Inductively,

$$\varphi(L_1^{(k)}) = \varphi([L_1^{(k-1)}, L_1^{(k-1)}]) = [\varphi(L_1^{(k-1)}), \varphi(L_1^{(k-1)})] = [L_2^{(k-1)}, L_2^{(k-1)}] = L_2^{(k)}$$

□

Lemma 3.1.6

Let L be a Lie algebra.

- (a) If L is solvable, then every subalgebra and every homomorphic image of L are solvable.
- (b) Suppose that L has an ideal I such that I and L/I are solvable. Then L is solvable.
- (c) If I and J are solvable ideals of L , then $I + J$ is a solvable ideal of L .

(a) Omitted.

Note.

- (b) 商去 I 是可解的, 当且仅当各层级的非交换性可以被 I 给兜住.

From the above proposition, we have $(L/I)^{(k)} = (L^{(k)} + I)/I$. If L/I is solvable, then $L^{(m)} \subseteq I$ for some m . We have for some k ,

$$(L^{(m)})^{(k)} \subseteq I^{(k)} = 0$$

Alternatively, it follows from (a) that $L^{(m)}$ is solvable.

$$L^{(m+k)} = (L^{(m)})^{(k)} = 0$$

- (c) $(I + J)/I \simeq J/I \cap J$ is solvable by (a). And since I is solvable, $I + J$ is solvable by (b).

Subalgebra of the General Linear Algebra

4.1 Nilpotent Maps

Lemma 4.1.1

Let $x \in L$. If the linear map $x : V \rightarrow V$ is nilpotent, then $\text{ad } x : L \rightarrow L$ is also nilpotent.

Proof. Take any $y \in L$, and expand

$$(\text{ad } x)^m(y) = [x, [x, \dots [x, y] \dots]]$$

which is of the form

$$x^i y x^j, \quad i + j = m$$

Suppose that $x^r = 0$, taking $m = 2r$, we have $(\text{ad } x)^m(y) = 0$ for all $y \in L$. $\square$

Lemma 4.1.2

Suppose that A is an ideal of a Lie subalgebra L of $\mathfrak{gl}(V)$. Let

$$W = \{v \in V : a(v) = 0 \text{ for all } a \in A\}.$$

Then W is an L -invariant subspace of V .

Remark.

我们熟知, 如果每个 a 和 y 可交换, 则 W 一定是 L 不变的. 因此 W 成为 L 不变子空间, 只在于是否在 $[a, y]$ 下保持, 这也是为什么我们要假设 A 是一个理想.

Proof. Take $w \in W$ and $y \in L$, we must show that $a(yw) = 0$ for all a . But $ay = ya + [a, y]$, we have

$$a(yw) = y(aw) + [a, y](w)$$

And $y(aw) = 0$ for $aw = 0$; $[a, y](w) = 0$ for $[a, y] \in A$. □

4.2 Weights

Definition 4.2.1

A **weight** for a Lie subalgebra A of $\mathfrak{gl}(V)$ is a linear map $\lambda : A \rightarrow F$ such that

$$V_\lambda := \{v \in V : a(v) = \lambda(a)v \text{ for all } a \in A\}$$

is a non-zero subspace of V .

4.3 Invariance Lemma

Lemma 4.3.1

Suppose that A is an ideal of a Lie subalgebra L of $\mathfrak{gl}(V)$. Let

$$W = \{v \in V : a(v) = 0 \text{ for all } a \in A\}.$$

Then W is an L -invariant subspace of V .

Lemma 4.3.2 ► Invariance Lemma

Assume that F has characteristic zero. Let L be a Lie subalgebra of $\mathfrak{gl}(V)$ and let A be an ideal of L . Let $\lambda : A \rightarrow F$ be a weight of A . The associated weight space

$$V_\lambda = \{v \in V : av = \lambda(a)v \text{ for all } a \in A\}$$

is an L -invariant subspace of V .

Engel's Theorem and Lie's Theorem

5.1 Engel's Theorem

Lemma 5.1.1

Suppose that L is a Lie subalgebra of $\mathfrak{gl}(V)$, where V is non-zero, such that every element of L is a nilpotent linear transformation. Then there is some non-zero $v \in V$ such that $xv = 0$ for all $x \in L$.

Proof sketch.

对维数归纳

对于任意 L 的真子代数 A , 模 A 的伴随:

$$\begin{aligned}\varphi : A &\rightarrow \mathfrak{gl}(L/A) \\ a &\mapsto (x + A \mapsto [a, x] + A)\end{aligned}$$

我们看到 $\varphi(A)$ 的公共 0-特征向量, 就是关于 A 代数封闭的向量.

根据归纳假设, $\varphi(A)$ 有非零公共 0-特征向量 y , 从而 $y \notin A$ 且 y 是关于 A 代数封闭的. 这就意味着 A 可以成为一个更大的代数 $\tilde{A} = A \oplus \text{span}\{y\}$ 的理想. 如果我们取 A 是极大的, 那么 $\tilde{A} = L$. 从而 A 成为 L 的一个余维数为 1 的理想.

一个引理是, L 的理想 A 的零子空间 W 是 L 不变的子空间. 这就意味着当我们寻找 $L = A \oplus \text{span}\{y\}$ 的公共零特征向量时, 可以从 A 的零子空间 W 中, 寻找 $\text{span}\{y\}$ 的公共零特征向量, 这是可行的, 因为 $\text{span}\{y\}$ 也是一组幂零的线性变换.

Proof. We induct on the dimension of L . Let A be a maximal subalgebra of L . Define a map

$$\begin{aligned}\varphi : A &\rightarrow \mathfrak{gl}(L/A) \\ a &\mapsto (x + A \mapsto [a, x] + A)\end{aligned}$$

It is straightforward to verify that φ is a Lie algebra homomorphism. Thus $\varphi(A)$ is a Lie algebra. For each $a \in A$, since a is nilpotent, $\text{ad } a$ is as well, hence $\varphi(a)$ is nilpotent. We know from inductive hypothesis that there exists nonzero $y + A \in L/A$ such that $0 = \varphi(a)(y + A)$, $\forall a \in A$. That is $y \notin A$ and

$[A, y] \subseteq A$. By defining $\tilde{A} := A \oplus \text{span}\{y\}$, we know A is an ideal of $\tilde{A}$. Since A is maximal, there must be $\tilde{A} = L$, A is an ideal of L with codimension 1.

A lemma gives that the space W

$$W = \{v \in V : a(v) = 0, \forall a \in A\}$$

is L -invariant. Specifically, $y(W) \subseteq W$. $y|_W$ is nilpotent, there exist nonzero $v \in W$ such that $y(v) = y|_W(v) = 0$. Then for each element of L , we can write it as $a + ky$ for some $k \in F$. Then

$$(a + ky)(v) = a(v) + ky(v) = 0$$

which completes the proof. □

Theorem 5.1.2 ▶ Engel's Theorem

Let V be a vector space. Suppose that L is a Lie subalgebra of $\mathfrak{gl}(V)$ such that every element of L is a nilpotent linear transformation of V . Then there is a basis of V in which every element of L is represented by a strictly upper triangular matrix.

Proof. We induct on the dimension of V . By Lemma 5.1.1, there exists $w \in V$ such that $x(w) = 0, \forall x \in L$. Let $W = \text{span}\{w\}$. Then every x induces a linear transformation $\bar{x} : V/W \rightarrow V/W$, $\bar{x}(v + W) = x(v) + W$. It is easily to check that the map

$$L \rightarrow \mathfrak{gl}(V/W), \quad x \mapsto \bar{x}$$

is a Lie homomorphism.

The image of L under this homomorphism is a subalgebra of $\mathfrak{gl}(V/W)$, which satisfies the hypothesis of Engel's Theorem. Moreover $\dim(V/W) = \dim V - 1$. We know from inductive hypothesis that there is a basis of V/W , say $\{v_i + W\}_{2 \leq i \leq n}$ in which every element of L is represented by a strictly upper triangular matrix. That is equivalent to for each $x \in L$,

$$\bar{x}(v_j + W) \in \text{span}\{v_2 + W, \dots, v_{j-1} + W\}, \quad \forall 3 \leq j \leq n,$$

and

$$\bar{x}(v_2 + W) = 0 + W$$

By taking $v_1 = w$, it follows that

$$x(v_1) = 0, \quad x(v_2) \in \text{span}\{v_1\}$$

$$x(v_j) \in \text{span}\{v_1, v_2, \dots, v_{j-1}\}, \quad \forall 3 \leq j \leq n$$

That is every element of L is represented by a strictly upper triangular matrix under the basis $\{v_1, v_2, \dots, v_n\}$. $\square$

Definition 5.1.3

Recall that a Lie algebra is said to be nilpotent if for some $m \geq 1$ we have $L^m = 0$ or, equivalently, if for all $x_0, x_1, \dots, x_m \in V$ we have

$$[x_0, [x_1, \dots, [x_{m-1}, x_m] \dots]] = 0.$$

Theorem 5.1.4 ► Engel's Theorem, second version

A Lie algebra L is nilpotent if and only if for all $x \in L$ the linear map $\text{ad } x : L \rightarrow L$ is nilpotent.

Proof. The ‘only if’ is straightforward. For the ‘if’ part, if all $\text{ad } x$ is nilpotent, the algebra

$$\text{ad}(L) = \{\text{ad } x : x \in L\}$$

is a subalgebra of $\mathfrak{gl}(L)$ satisfying the hypothesis of Engel's Theorem. it follows from Engel's Theorem that there exists a basis of L say $\{e_1, \dots, e_n\}$, such that

$$[L, e_i] \subseteq \text{span}\{e_1, \dots, e_{i-1}\}$$

Denote L_j by $\text{span}\{e_1, \dots, e_j\}$, then

$$[L, L_j] \subseteq L_{j-1}, j \geq 2, \quad [L, L_1] = 0$$

We have

$$L^0 = L = L_n$$

$$L^1 = [L, L] \subseteq L_{n-1}$$

Inductively, we have

$$L^k = [L, L^{k-1}] \subseteq [L, L_{n-k+1}] \subseteq L_{n-k}$$

Finally, we have

$$L^n = 0$$

L is nilpotent. □

5.2 Lie's Theorem

Proposition 5.2.1

Let V be a non-zero complex vector space. Suppose that L is a solvable Lie subalgebra of $\mathfrak{gl}(V)$. Then there is some non-zero $v \in V$ which is a simultaneous eigenvector for all $x \in L$

Proof sketch.

- 一个李代数 L 的子代数 A 如果包含了 L' , 则 A 是 L 的一个理想.
- 如果 L 是可解的, 则 L' 是真子空间, 这意味着我们可以构造 L 的余一维的理想.
- 归纳假设给出 A 的一个特征向量, 从而对应的 weight 子空间非空.
- 理想的 Weight 子空间是 L -不变的, 特别地是 v -不变的.
- 由于空间是 $\mathbb{C}$ 上的, 在 Weight 子空间上存在特征向量 v , 它就是公共的特征向量.

Theorem 5.2.2 ▶ Lie's Theorem

Let V be an n -dimensional complex vector space and let L be a solvable Lie subalgebra of $\mathfrak{gl}(V)$. Then there is a basis of V in which every element of L is represented by an upper triangular matrix.

Proof sketch.

证明完全类似于 Engel 定理的证明.

Representation of $\mathfrak{sl}(2, \mathbb{C})$

6.1 The Modules V_d

6.2 Classifying the Irreducible $\mathfrak{sl}(2, \mathbb{C})$ -Modules

Lemma 6.2.1

Suppose that V is an $\mathfrak{sl}(2, \mathbb{C})$ -module and $v \in V$ is an eigenvector of h with eigenvalue λ .

- Either $e \cdot v = 0$ or $e \cdot v$ is an eigenvector of h with eigenvalue $\lambda + 2$.
- Either $f \cdot v = 0$ or $f \cdot v$ is an eigenvector of h with eigenvalue $\lambda - 2$.

Lemma 6.2.2

Let V be a finite-dimensional $\mathfrak{sl}(2, \mathbb{C})$ -module. Then V contains eigenvector w for h such that $e \cdot w = 0$.

Theorem 6.2.3

If V is finite-dimensional irreducible $\mathfrak{sl}(2, \mathbb{C})$ -module, then V is isomorphic to one of the V_d .

Corollary 6.2.4

If V is a finite-dimensional representation of $\mathfrak{sl}(2, \mathbb{C})$ and $w \in V$ is an h -eigenvector such that $e \cdot w = 0$, then $h \cdot w = dw$ for some non-negative integer d and the submodule of V generated by w is isomorphic to V_d .

6.3 Weyl's Theorem

Theorem 6.3.1

Let L be a complex semisimple Lie algebra. Every finite-dimensional representation of L is completely reducible.

Cartan Criteria

7.1 Jordan Decomposition

Lemma 7.1.1

Let x be a linear transformation of the complex vector space V . Suppose that x has Jordan decomposition $x = d + n$, where d is diagonalisable, n is nilpotent, and d and n commute.

- (a) There is a polynomial $p(X) \in \mathbb{C}[X]$ such that $p(x) = d$.
- (b) Fix a basis of V in which d is diagonal. Let $\bar{d}$ be the linear map whose matrix with respect to this basis is the complex conjugate of the matrix of d . There is a polynomial $q(X) \in \mathbb{C}[X]$ such that $q(x) = \bar{d}$.

Exercise 7.1.2

Let V be a vector space, and suppose that $x \in \mathfrak{gl}(V)$ has Jordan decomposition $d + n$. Show that $\text{ad } x : \mathfrak{gl}(V) \rightarrow \mathfrak{gl}(V)$ has Jordan decomposition $\text{ad } d + \text{ad } n$.

Proof. Only need to show that $\text{ad } d$ is semisimple and $\text{ad } n$ is nilpotent, then the proposition follows from the linearity of ad and the uniqueness of Jordan decomposition.

- Suppose that $e_1, \dots, e_n$ is the basis in which d is represented diagonal, say $\text{diag}(\lambda_1, \dots, \lambda_n)$. Let $E_{ij} \in \mathfrak{gl}(V)$ be the maps that mapping e_j to e_i , that is $E_{ij}(e_k) = \delta_j^k e_i$, $k = 1, \dots, n$. Then

$$\begin{aligned}
 \text{ad}(d)(E_{ij})(e_k) &= [d, E_{ij}](e_k) \\
 &= (dE_{ij} - E_{ij}d)(e_k) \\
 &= d(E_{ij}e_k) - E_{ij}(de_k) \\
 &= d(\delta_j^k e_i) - E_{ij}(\lambda_k e_k) \\
 &= \delta_j^k \lambda_i e_i - \lambda_k \delta_j^k e_i \\
 &= (\lambda_i - \lambda_k) \delta_j^k e_i = (\lambda_i - \lambda_j) E_{ij}
 \end{aligned}$$

Which implies that E_{ij} is an eigenvector for $\text{ad}(d)$ for each $i, j = 1, \dots, n$.
 $\text{ad } n$ is diagonal under the basis $E_{ij}, i, j = 1, \dots, n$

- Since n is nilpotent, it follows that $\text{ad } n$ is nilpotent.
- It follows from Jacobi identity that

$$(\text{ad } d) \circ (\text{ad } n)(y) = [d, [n, y]] = -[y, [d, n]] - [n, [y, d]]$$

Since d, n commutes, $[d, n] = 0$. Thus $\text{ad } n$ and $\text{ad } d$ commute.

□

7.2 Testing for Solvability

Exercise 7.2.1

Let V be a complex vector space, L be a Lie subalgebra of $\mathfrak{gl}(V)$. Suppose that L is solvable. Use Lie's Theorem to show that there is a basis of V in which every element of L' is represented by a strictly upper triangular matrix. Conclude that $\text{tr } xy = 0$ for all $x \in L$ and $y \in L'$.

Remark.

$\mathbb{C}$ can be replaced by $\mathbb{F}$ such that $\text{char } \mathbb{F} = 0, \bar{\mathbb{F}} = \mathbb{F}$.

Proof. From Lie's Theorem, there is a basis of V in which every element of L is upper triangular, say $\{e_1, \dots, e_n\}$. Take $z \in L', z = [x, y], x, y \in L$. Then

$$z(e_1) = [x, y](e_1) = x(y(e_1)) - y(x(e_1)) = 0$$

Since e_1 is simultaneously an eigenvector of x and y . Let $L_k = \text{span}\{e_1, \dots, e_k\}$, then

$$z(e_k) = x(y(e_k)) - y(x(e_k))$$

Suppose,

$$y(e_k) = \sum_{j=1}^k y_j e_j, \quad x(e_k) = \sum_{j=1}^k x_j e_j$$

then

$$\begin{aligned} x(y(e_k)) + \text{span}\{e_1, \dots, e_{k-1}\} &= x(y_k e_k) + \text{span}\{e_1, \dots, e_{k-1}\} \\ &= x_k y_k e_k + \text{span}\{e_1, \dots, e_{k-1}\} \end{aligned}$$

Similarly,

$$y(x(e_k)) + \text{span}\{e_1, \dots, e_{k-1}\} = x_i y_i e_k + \text{span}\{e_1, \dots, e_{k-1}\}$$

Thus

$$z(e_k) \in \text{span}\{e_1, \dots, e_{k-1}\}$$

It follows that z is nilpotent. Thus every element of L' is represented by a strictly upper triangular matrix.

The matrix of x and y under the basis $\{e_1, \dots, e_n\}$ are upper triangular matrix and strictly upper triangular matrix, respectively, say A, B . Then

$$\text{tr}xy = \text{tr}(AB) = \sum_{k=1}^n a_{ik}b_{kj} = 0$$

Since for each $i > k$, $a_{ik} = 0$, and for each $i \leq k$, $b_{kj} \leq 0$. □

Proposition 7.2.2

Let V be a complex vector space and let L be a Lie subalgebra of $\mathfrak{gl}(V)$. If $\text{tr}xy = 0$ for all $x, y \in L$, then L is solvable.

Proof. We shall show that every element of L' is nilpotent, then it follows from Engel's Theorem that L' is nilpotent, thus L is solvable.

Take any element $x \in L'$. Suppose that x has a Jordan decomposition $x = d + n$. We choose a basis in which d is diagonal $d = \text{diag}(\lambda_1, \dots, \lambda_n)$ and n is strictly upper triangular. We only need to show that

$$\begin{aligned} \sum_{i=1}^n \lambda_i \bar{\lambda}_i &= 0 \\ \text{tr}(\bar{d}x) &= \text{tr}(\bar{d}d + \bar{d}n) = \text{tr}(\bar{d}d + \bar{d}n) \\ &= \text{tr}(\bar{d}d) + 0 \\ &= \sum_{i=1}^n \lambda_i \bar{\lambda}_i \end{aligned}$$

Since $x \in L'$, we write $x = [y, z]$, $y, z \in L'$.

$$\begin{aligned}\operatorname{tr}(\bar{d}x) &= \operatorname{tr}(\bar{d}[y, z]) \\ &= \operatorname{tr}(\bar{d}yz - \bar{d}zy) = \operatorname{tr}(\bar{d}yz - y\bar{d}z) \\ &= \operatorname{tr}([\bar{d}, y]z)\end{aligned}$$

Note.

这里我们折腾 $\operatorname{ad} \bar{d}$ 而非 $\bar{d}$ 的意义在于, 多项式对不变性的保持需要我们对结合代数封闭, 而李代数不一定保持结合代数的封闭性.

Thus to show $d = 0$, we only need to show that $[\bar{d}, y] \in L$, that is $\operatorname{ad}_{\bar{d}}(y) \in L$. $\operatorname{ad} x$ has Jordan decomposition, $\operatorname{ad} x = \operatorname{ad} d + \operatorname{ad} n$. Then from Lemma 7.1.1, there exists $q(X) \in \mathbb{C}[X]$ such that

$$q(\operatorname{ad} x) = \operatorname{ad} \bar{d}$$

Since L is $\operatorname{ad} x$ -invariant, so dose $q(\operatorname{ad} x)$ and $\operatorname{ad} \bar{d}$, which completes the proof. $\square$

Theorem 7.2.3

Let L be a complex Lie algebra. Then L is solvable if and only if $\operatorname{tr}(\operatorname{ad} x \circ \operatorname{ad} y) = 0$ for all $x \in L$ and all $y \in L'$.

Note.

由于 L 的 solvability 等价于任意阶导出子代数 $L^{(k)}$ 的 solvability, 因此命题中的

$$x \in L, \quad y \in L'$$

对于任意固定的 $k \geq 1, l \geq 0$ 可以替换为

$$x \in L^{(k)}, \quad y \in L^{(l)},$$

Proof. Suppose L is solvable. Then $\operatorname{ad}(L) \subseteq \mathfrak{gl}(L)$ is solvable. Then it follows from exercise 7.2.1 that $\operatorname{tr}(\operatorname{ad} x \circ \operatorname{ad} y) = 0$ for all $x \in L, y \in L'$.

Conversly, it follows from proposition 7.2.2 that $\operatorname{ad}(L')$ is solvable, then so dose L' and L . $\square$

7.3 Killing Form

Definition 7.3.1

Let L be a complex Lie algebra. The Killing form on L is the symmetric bilinear form defined by

$$\kappa(x, y) := \text{tr}(\text{ad } x \circ \text{ad } y), \quad \text{for } x, y \in L$$

Proposition 7.3.2

$$\kappa([x, y], z) = \kappa(x, [y, z]).$$

Theorem 7.3.3 ► Cartan's First Criterion

The complex Lie algebra L is solvable if and only if $\kappa(x, y) = 0$ for all $x \in L$ and $y \in L$.

Proof. Just a restatement of theorem 7.2.3 . □

Lemma 7.3.4

Suppose that L is a Lie algebra and I is an ideal of L . We write κ for the Killing form on L and κ_I for the Killing form on I , considered as a Lie algebra in its own right. If $x, y \in I$, then $\kappa_I(x, y) = \kappa(x, y)$.

Proof. We expand a basis of I to a basis of L . Since

$$\text{ad } x(L) \subseteq [x, L] \subseteq I$$

Then $\text{ad } x$ is represented of the form

$$\begin{pmatrix} A_x & B_x \\ 0 & 0 \end{pmatrix}$$

$\text{ad } y$ is represented of the form

$$\begin{pmatrix} A_y & B_y \\ 0 & 0 \end{pmatrix}$$

We have

$$\begin{aligned}
 \kappa(x, y) &= \text{tr} \left(\begin{pmatrix} A_x & B_x \\ O & O \end{pmatrix} \begin{pmatrix} A_y & B_y \\ O & O \end{pmatrix} \right) \\
 &= \text{tr} \left(\begin{pmatrix} A_x A_y & A_x B_y \\ O & O \end{pmatrix} \right) \\
 &= \text{tr}(A_x A_y) \\
 &= \kappa_I(x, y)
 \end{aligned}$$

□

7.3.1 Testing for Semisimplicity

Definition 7.3.5 ► perpendicular space

- Let β be a symmetric bilinear form on a finite-dimensional complex vector space V . If S is a subset of V , we define the **perpendicular space** to S by

$$S^\perp = \{x \in V : \beta(x, s) = 0 \text{ for all } s \in S\}.$$

- We say that β is **non-degenerate** if $V^\perp = 0$; that is, there is no non-zero vector $v \in V$ such that $\beta(v, x) = 0$ for all $x \in V$.

Proposition 7.3.6

If β is non-degenerate and W is a vector subspace of V , then

$$\dim W + \dim W^\perp = \dim V.$$

Proof. Define a linear map $\varphi : V \rightarrow V^*$,

$$\varphi(v) \mapsto (u \mapsto \beta(u, v))$$

Then

$$\ker \varphi = \{v \in V : \beta(u, v) = 0, \forall u \in V\} = V^\perp = 0$$

Since V is finite-dimensional, we have $\dim V = \dim V^*$. Then φ is an isomorphism. Define $W^0 \subseteq V^*$:

$$W^0 = \{\lambda \in V^* : \lambda(w) = 0, \forall w \in W\}$$

We have

$$\varphi(W^\perp) = W^0, \quad \dim W^\perp = \dim W^0$$

Let $i : W \hookrightarrow V$ the inclusion map, for each $\lambda^* \in V^*$, we define a map $i^* : V^* \rightarrow W^*$

$$i^*(\lambda)(w) = \lambda(i(w))$$

Then $\text{Im } i^* = W^*$, $\ker i^* = W^0$, we have

$$\dim W^0 + \dim W^* = \dim V^*$$

Then it follows that

$$\dim W^\top + \dim W^* = \dim V^* \implies \dim W^\perp + \dim W = \dim V$$

Since W, V have finite dimension. □

Exercise 7.3.7

Suppose I is an ideal of L . Shows that $I^\perp$ is an ideal of L .

Proof.

$$I^\perp = \{x \in L : \kappa(i, x) = 0, \forall i \in I\}$$

Take $x \in I^\perp, y \in L$, note that

$$[y, x] \in I^\perp \iff \kappa([x, y], i) = 0, \quad \forall i \in I$$

Since we have

$$\kappa(x, [y, i]) = 0, \forall y \in L, i \in I$$

Then

$$\kappa([x, y], i) = \text{tr}([\text{ad}_x, \text{ad}_y] \circ \text{ad}_i) = \text{tr}(\text{ad}_x \circ [\text{ad}_y \circ \text{ad}_i]) = \kappa(x, [y, i]) = 0$$

For all $i \in I$. That is $[x, y] \in I^\perp$. $[L, I^\perp] \subseteq I^\perp$. □

Corollary 7.3.8

If L is semisimple, then κ is non-degenerate.

Proof. Other wise, $L^\perp$ is an nonzero ideal of L . Since $(L^\perp)' \subseteq L$, we have

$$\kappa(L^\perp, (L^\perp)') \subseteq \kappa(L^\perp, L) = 0$$

Then it follows from the Cartan First Criteria that $L^\top$ is solvable. Then $L^\perp$ is a nonzero solvable ideal of L , which is a contradiction to the Semisimplicity of L . $\square$

Theorem 7.3.9 ► Cartan's Second Criterion

The complex Lie algebra L is semisimple if and only if the Killing form κ of L is non-degenerate.

7.4 Derivations of Semisimple Lie Algebras

Proposition 7.4.1

If L is a finite-dimensional complex semisimple Lie algebra, then $\text{ad } L = \text{Der } L$.

Proof. For each $x \in L$,

$$\begin{aligned} \text{ad } x [y, z] &= [x, [y, z]] \\ &= -[z, [x, y]] - [y, [z, x]] \\ &= [\text{ad } x (y), z] + [y, \text{ad } x (z)] \end{aligned}$$

$\text{ad } x$ is a derivation. ad is a Lie algebra homomorphism from L to $\text{Der } L$. For each $\delta \in \text{Der } L$,

$$\begin{aligned} [\delta, \text{ad } x]y &= \delta [x, y] - \text{ad } x (\delta y) \\ &= [\delta x, y] + [x, \delta y] - [x, \delta y] \\ &= \text{ad } (\delta x) (y) \end{aligned}$$

Thus $[\delta, \text{ad } x] = \text{ad } (\delta x)$, the image of ad is an ideal of $\text{Der } L$.

Now we bring in our assumption that L is complex and semisimple. First, note that $\text{ad} : L \rightarrow \text{Der } L$ is one-to-one, as $\ker \text{ad} = Z(L) = 0$, so the Lie algebra $M := \text{ad } L$ is isomorphic to L , therefore it is semisimple as well. We have

$$\dim M + \dim M^\perp = \dim \text{Der } L$$

so our aim is to show that $\dim M^\perp = 0$. As M is an ideal of $\text{Der } L$, the Killing form κ_M of M is the restriction of the Killing form on $\text{Der } L$. By Cartan's Second Criterion, κ_M is non-degenerate, so $M^\perp \cap M = 0$, and hence $[M^\top, M] = 0$. Thus if $\delta \in M^\perp$, and $\text{ad } x \in M$, then $[\delta, \text{ad } x] = 0$ for $M, M^\perp$ are ideals. But

we saw above that

$$[\delta, \text{ad } x] = \text{ad } (\delta x)$$

so, for all $x \in L$, we have $\delta(x) = 0$; in other words, $\delta = 0$. □

7.5 Abstract Jordan Decomposition

Proposition 7.5.1

Let L be a complex Lie algebra. Suppose that δ is a derivation of L with Jordan decomposition $\delta = \sigma + \nu$, where σ is diagonalisable and ν is nilpotent. Then σ and ν are also derivations of L .

Proof. For $\lambda \in \mathbb{C}$, let

$$L_\lambda = \{x \in L : (\delta - \lambda) x = 0\}$$

L decomposes as a direct sum of generalised eigenspaces

$$L = \bigoplus_\lambda L_\lambda$$

As σ acts diagonalisably, then λ -eigenspace of σ is L_λ . Take $x \in L_\lambda$, $y \in L_\mu$, then

$$\begin{aligned} (\delta - (\lambda + \mu) 1_L) [x, y] &= [\delta x, y] + [x, \delta y] - [\lambda x, y] - [x, \mu y] \\ &= [(\delta - \lambda 1_L) x, y] + [x, (\delta - \mu 1_L) y] \end{aligned}$$

$(\delta - \lambda 1_L)^{m+n} [x, y]$ is the linear combination of the form $[(\delta - \lambda 1_L)^i x, (\delta - \mu 1_L)^j y]$, which is zero. Thus $[x, y] \in L_{\lambda+\mu}$, so

$$\sigma([x, y]) = (\lambda + \mu) [x, y]$$

which is the same as

$$[\sigma(x), y] + [x, \sigma(y)] = [\lambda x, y] + [x, \mu y]$$

Thus σ is a derivation, and so is $\delta = \sigma + \nu$. □

Theorem 7.5.2

Let L be a complex semisimple Lie algebra. Each $x \in L$ can be written uniquely as $x = d + n$, where $d, n \in L$ are such that $\text{ad } d$ is diagonalisable, $\text{ad } n$ is nilpotent, and $[d, n] = 0$. Furthermore, if $y \in L$ commutes

with x , then $[d, y] = 0$ and $[n, y] = 0$.

Proof. Let $\text{ad } x = \sigma + \nu$, where $\sigma \in \mathfrak{gl}(L)$ is diagonalisable, $\nu \in \mathfrak{gl}(L)$ is nilpotent, and $[\sigma, \nu] = 0$. We know that σ and ν are derivations of the semisimple Lie algebra L . And we know that $\text{ad } L = \text{Der } L$, so there exists $d, n \in L$ such that $\text{ad } d = \sigma$ and $\text{ad } n = \nu$. As ad is injective and

$$\text{ad } x = \sigma + \nu = \text{ad } (d + n)$$

we get that $x = d + n$. Moreover, $\text{ad } [d, n] = [\text{ad } d, \text{ad } n] = 0$. The uniqueness of d and n follows from the uniqueness of the Jordan decomposition of $\text{ad } x$. Suppose that $y \in L$ and that $(\text{ad } x)y = 0$. Since σ and ν may be expressed as polynomials in $\text{ad } x$. Let

$$\nu = c_0 1_L + c_1 \text{ad } x + \cdots + c_r (\text{ad } x)^r$$

Applying ν to y we see that $\nu(y) = c_0 y$. But ν is nilpotent and $\nu(x) = c_0 x$, so $c_0 = 0$. Thus $\nu(y) = 0$, and so $\sigma(y) = (\text{ad } x - \nu)y = 0$ also. $\square$

Definition 7.5.3

Let L be a semisimple Lie algebra and let $\theta : L \rightarrow \mathfrak{gl}(V)$ be a representation of L . Suppose that $x \in L$ has abstract Jordan decomposition $x = d + n$. Then the Jordan decomposition of $\theta(x) \in \mathfrak{gl}(V)$ is $\theta(x) = \theta(d) + \theta(n)$.

The Root Space Decomposition

8.1 Preliminary Results

Proposition 8.1.1

Suppose that L is a complex semisimple Lie algebra containing an abelian subalgebra H consisting of semisimple elements.

For each $\alpha \in H^*$, let

$$L_\alpha := \{x \in L : [h, x] = \alpha(h)x, \quad \forall h \in H\}$$

Let Φ denote the set of non-zero $\alpha \in L^*$ for which L_α is non-zero. We can write the decomposition of L into weight spaces for H as

$$L = L_0 \oplus \bigoplus_{\alpha \in \Phi} L_\alpha. \quad (\star)$$

Since L is finite-dimensional, this implies that Φ is finite.

Lemma 8.1.2

Suppose $\alpha, \beta \in H^*$, then

1. $[L_\alpha, L_\beta] = L_{\alpha+\beta}$
2. If $\alpha + \beta \neq 0$, then $\kappa(L_\alpha, L_\beta) = 0$.
3. The restriction of κ to L_0 is non-degenerate.

Proof. 1. Take $x \in L_\alpha, y \in L_\beta$, then

$$\begin{aligned} [h, [x, y]] &= -[y, [h, x]] - [x, [y, h]] \\ &= -[y, \alpha x] - [x, -\beta y] \\ &= (\alpha + \beta)[x, y] \end{aligned}$$

2. If $\alpha + \beta \neq 0$, then $\alpha + \beta \neq 0$. Take $x \in L_\alpha, y \in L_\beta$

$$\begin{aligned}\alpha(h)\kappa(x, y) &= \kappa(\alpha(h)x, y) \\ &= \kappa([h, x], y) \\ &= -\kappa([x, h], y) \\ &= -\kappa(x, [h, y]) \\ &= -\kappa(x, \beta(h)y) \\ &= -\beta(h)\kappa(x, y)\end{aligned}$$

Thus

$$(\alpha(h) + \beta(h))\kappa(x, y) = 0$$

Then it follows that $\kappa(x, y) = 0$ since $\alpha(h) + \beta(h) \neq 0$

3. Suppose that $z \in L_0, \kappa(z, x_0) = 0$ for all $x_0 \in L_0$. From 2., for each $\alpha \neq 0$, we have $\kappa(z, x_\alpha) = 0$ for all $x_\alpha \in L_\alpha$. Then for each element of x , we write

$$x = x_0 + \sum_{\alpha \in \Phi} x_\alpha$$

there is

$$\kappa(z, x) = \kappa(z, x_0) + \sum_{\alpha \in \Phi} \kappa(z, x_\alpha) = 0$$

Since L is semisimple, κ is non-degenerate (Cartan second Criterion). It follows that $z = 0$.

□

Exercise 8.1.3

Show that if $x \in L_\alpha$, where $\alpha \neq 0$, then ad_x is nilpotent.

Proof. Suppose that $\Phi = \{\alpha_0 := \alpha, \alpha_1, \dots, \alpha_k\}$, then for each $j = 0, 1, \dots, k$, since $\alpha_j \neq 0$, there exists $m_j \in \mathbb{Z}_{>0}$ such that $\alpha_j + m_j\alpha$ is different from any element of $\Phi \cup \{0\}$. Then

$$(\text{ad}_x)^{m_j}(x_{\alpha_j}) = 0, \quad \forall j = 0, 1, \dots, k, \quad \forall x_{\alpha_j} \in L_{\alpha_j}$$

Furthermore,

$$(\text{ad}_x)^{m_0+1}(x_0) = (\text{ad}_x)^{m_0}([x, x_0]) = 0, \quad \forall x_0 \in L_0$$

since $[x, x_0] \in L_\alpha$. Set $N = \max\{m_0 + 1, m_1, \dots, m_k\}$. Then

$$(\text{ad}_x)^N(L) = (\text{ad}_x)^N(L_0 + \bigoplus_{\alpha_j \in \Phi} L_{\alpha_j}) = 0$$

ad_x is nilpotent. □

Remark.

If H is small, then the decomposition (★) is likely to be rather coarse, with few non-zero weight spaces other than L_0 . Furthermore, if H is properly contained in L_0 , then we get little information about how the elements in L_0 that are not in H act on L . This is illustrated by the following exercise.

Exercise 8.1.4

Let $L = \mathfrak{sl}(n, \mathbb{C})$, where $n \geq 2$, and let $H = \text{Span}\{h\}$, where $h = e_{11} - e_{22}$. Find $L_0 = C_L(H)$, and determine the direct sum decomposition (★) with respect to H .

Solution. Take $x \in \mathfrak{sl}(n, \mathbb{C})$, suppose $x = \sum_{i \neq j, \text{ and } i, j \neq 1, 2} x^{ij} x_{ij} + x^{12} e_{12} + x^{21} e_{21} + y^k \sum_{k=2}^n (e_{11} - e_{kk})$. For each $1 \leq i \neq j \leq n$ If $i, j \neq 1, 2$, then

$$[e_{11}, e_{ij}] = e_{11} e_{ij} - e_{ij} e_{11} = 0$$

$$[e_{22}, e_{ij}] = e_{22} e_{ij} - e_{ij} e_{22} = 0$$

$$[h, x^{12} e_{12}] = x^{12} [h, e_{12}] = x^{12} (e_{11} e_{12} - e_{12} e_{11} - e_{22} e_{12} + e_{12} e_{22}) = 2x^{12} e_{12}$$

$$[h, x^{21} e_{21}] = x^{21} [h, e_{21}] = x^{21} (e_{11} e_{21} - e_{21} e_{11} - e_{22} e_{21} + e_{21} e_{22}) = -2x^{21} e_{21}$$

$$[h, e_{11} - e_{kk}] = [e_{22}, e_{kk}] = 0$$

Then $x \in L_0 \iff x^{12} = x^{21} = 0$.

$$L_0 = \text{span}(\{e_{ij}, i \neq j, \text{ and } i, j \neq 1, 2\} \cup \{e_{12} - e_{21}\} \cup \{e_{11} - e_{kk} : k = 2, 3, \dots, n\})$$

◇

8.2 Cartan Subalgebras

Definition 8.2.1

A Lie subalgebra H of a Lie algebra L is said to be a **Cartan subalgebra** (or CSA) if H is abelian and every element $h \in H$ is semisimple, and moreover H is maximal with these properties.

Lemma 8.2.2

Let H be a Cartan subalgebra of L . Suppose that $h \in H$ is such that the dimension of $C_L(h)$ is minimal. Then every $s \in H$ is central in $C_L(h)$, and so $C_L(h) \subseteq C_L(s)$. Hence $C_L(h) = C_L(H)$.

Theorem 8.2.3

If H is a Cartan subalgebra of L and $h \in H$ is such that $C_L(h) = C_L(H)$, then $C_L(h) = H$. Hence H is self-centralising.

8.3 Definition of the Root Space Decomposition

Definition 8.3.1

Let H be a Cartan subalgebra of our semisimple Lie algebra L . As $H = C_L(H)$, the direct sum decomposition of L into weight spaces for H considered in §10.1 may be written as

$$L = H \oplus \bigoplus_{\alpha \in \Phi} L_{\alpha},$$

where Φ is the set of $\alpha \in H^*$ such that $\alpha \neq 0$ and $L_{\alpha} \neq 0$. Since L is finite-dimensional, Φ is finite.

- If $\alpha \in \Phi$, then we say that α is a **root** of L and L_{α} is the associated **root space**.
- The direct sum decomposition above is the **root space decomposition**.

Remark.

It should be noted that the roots and root spaces depend on the choice of Cartan subalgebra H .

Lemma 8.3.2

Let $x, y : V \rightarrow V$ be linear maps from a complex vector space V to itself. Suppose that x and y both commute with $[x, y]$. Then $[x, y]$ is a nilpotent map.

Proof. Let $S = \text{span}\{x, y\}$, then

$$S' = \text{span}\{[x, y]\}$$

commutes with all element of S , that is $[S', S] = 0$, it follows that S is nilpotent, thus it is solvable. From Lie's theorem, there is a basis of V in which x, y is represented upper triangular. Let $L_i = \text{span}\{e_1, \dots, e_i\}$, then L_i is x -invariant and y -invariant, $i = 1, \dots, n$. There is $x^{ii}, y^{ii}, i = 1, \dots, n$ such that

$$x(e_i) + L_{i-1} = x^{ii}e_i, \quad y(e_i) + L_{i-1} = y^{ii}e_i.$$

Then

$$[x, y](e_i) + L_{i-1} = x^{ii}y^{ii}e_i - y^{ii}x^{ii}e_i + L_{i-1} = 0 + L_{i-1}$$

Thus $[x, y](e_i) \in L_{i-1}$. Thus $[x, y]$ is nilpotent. $\square$

Lemma 8.3.3

Suppose that $\alpha \in \Phi$ and that x is a non-zero element in L_α . Then $-\alpha$ is a root and there exists $y \in L_{-\alpha}$ such that $\text{Span}\{x, y, [x, y]\}$ is a Lie subalgebra of L isomorphic to $\mathfrak{sl}(2, \mathbb{C})$.

Proof. κ is non-degenerate, there exists $w = y_0 + \sum_{\beta \in \Phi} y_\beta$ such that $\kappa(x, w) \neq 0$. But for each $\beta \neq \alpha, \kappa(x, y_\beta) = 0$, there must be $-\alpha \in \Phi, \kappa(x, y_{-\alpha}) = \kappa(x, w) \neq 0$. We take $y = y_{-\alpha}$. Since $\alpha \neq 0$, there exists $t \in H$ such that $\alpha(t) \neq 0$. Then

$$\kappa(t, [x, y]) = \kappa([t, x], y) = \alpha(t) \kappa(x, y) \neq 0$$

thus $[x, y] \neq 0$.

Let $S := \text{span}\{x, y, [x, y]\}$, $[x, y]$ lies in $L_0 = H$. As x and y are simultaneous eigenvectors for all elements of $\text{ad } H$, and so in particular for $\text{ad}[x, y]$, this shows that S is a Lie subalgebra of L . It remains to show that S is isomorphic to $\mathfrak{sl}(2, \mathbb{C})$.

Let $h := [x, y] \in S$. We claim that $\alpha(h) \neq 0$. If not, then $[h, x] = \alpha(h)x = 0$;

similarly $[x, y] = -\alpha(h)x = 0$, so $\text{ad } h : L \rightarrow L$ commutes with $\text{ad } x$ and $\text{ad } y$, thus is nilpotent. The other hand, because H is a Cartan subalgebra, h is semisimple. But the only element of L that is both semisimple and nilpotent is 0, so $h = 0$, a contradiction.

Thus S is a 3-dimensional complex Lie algebra with $S' = S$. S is isomorphic to $\mathfrak{sl}(2, \mathbb{C})$. $\square$

Exercise 8.3.4

Show that for each $\alpha \in \Phi$, S_α has a basis $\{e_\alpha, f_\alpha, h_\alpha\}$ such that

1. $e_\alpha \in L_\alpha, f_\alpha \in L_{-\alpha}, h_\alpha \in H$ and $\alpha(h_\alpha) = 2$.
2. The map $\theta : S_\alpha \rightarrow \mathfrak{sl}(2, \mathbb{C})$, defined by $\theta(e_\alpha) = e, \theta(f_\alpha) = f, \theta(h_\alpha) = h$ is a Lie algebra isomorphism.

Proof. Let $x \in L_\alpha$. Since κ is non-degenerate, there exists $w = y_0 + \sum_{\alpha \in \Phi} y_\alpha$ such that $\kappa(x, w) \neq 0$, then

$$\kappa(x, w) = \kappa\left(x, y_0 + \sum_{\alpha \in \Phi} y_\alpha\right) = \kappa(x, y_{-\alpha}) \neq 0$$

Denote $y = y_{-\alpha}$.

$[x, y] \in L_{\alpha+(-\alpha)} = L_0 = H$. Take $h = [x, y]$. $h(x) = \alpha(h)x$

$$\kappa(h(x), y) = \alpha(h)\kappa(x, y) \neq 0$$

Then $\alpha(h) \neq 0$. Take $e_\alpha = x, f_\alpha = \frac{2}{\alpha(h)}y, h_\alpha = [e_\alpha, f_\alpha]$. Then

$$h_\alpha(e_\alpha) = \frac{2}{\alpha(h)}h(x) = 2x = 2e_\alpha$$

Let $S_\alpha = \text{span}\{e_\alpha, f_\alpha, h_\alpha\}$. Then $S'_\alpha = S_\alpha$. S is isomorphic to $\mathfrak{sl}(2, \mathbb{C})$.

$$[e_\alpha, f_\alpha] = h_\alpha, \quad [h_\alpha, e_\alpha] = 2e_\alpha, \quad [h_\alpha, f_\alpha] = 2f_\alpha$$

$\square$

8.4 Root Strings and Eigenvalues

Lemma 8.4.1

Given $h \in H$, let θ_h denote the map $\theta_h \in H^*$ defined by

$$\theta_h(k) = \kappa(h, k), \quad \forall k \in H$$

Then it is an isomorphism since $\kappa|_H$ is non-degenerate. In particular, for $\alpha \in H^*$, there is a unique element $t_\alpha \in H$ such that

$$\kappa(t_\alpha, k) = \alpha(k), \quad \forall k \in H$$

Lemma 8.4.2

Let $\alpha \in \Phi$. If $x \in L_\alpha$, $y \in L_{-\alpha}$, then $[x, y] = \kappa(x, y)t_\alpha$. In particular $h_\alpha = [e_\alpha, f_\alpha] \in \text{span}\{t_\alpha\}$.

Proof. For $h \in H$, we have

$$\kappa(h, [x, y]) = \alpha(h)\kappa(x, y) = \kappa(t_\alpha, h)\kappa(x, y) = \kappa(h, \kappa(x, y)t_\alpha)$$

This shows that $[x, y] - \kappa(x, y)t_\alpha \in H^\perp = 0$. Thus $[x, y] = \kappa(x, y)t_\alpha$ □

Definition 8.4.3

Let α be a root. We may regard L as an $\mathfrak{sl}(\alpha)$ -module via restriction of the adjoint representation. Thus if $a \in \mathfrak{sl}(\alpha)$ and $y \in L$, then the action is given by

$$a \cdot y = (\text{ad } a)(y) = [a, y]$$

Remark.

- The $\mathfrak{sl}(\alpha)$ -submodules of L are precisely the vector subspaces M of L such that $[s, m] \in M$ for all $s \in \mathfrak{sl}(\alpha)$ and $m \in M$.

Lemma 8.4.4

If M is a $\mathfrak{sl}(\alpha)$ -submodule of L , then the eigenvalues of h_α acting on M are integers.

Proof. Since M is finite-dimensional, by Weyl's Theorem, M may be decomposed into a direct sum of irreducible $\mathfrak{sl}(\alpha)$ -modules; for irreducible $\mathfrak{sl}(2, \mathbb{C})$ -

modules, the eigenvalues of h acting on M are integers. The lemma follows from the isomorphism between $\mathfrak{sl}(\alpha)$ and $\mathfrak{sl}(2, \mathbb{C})$. $\square$

Definition 8.4.5

If $\beta \in \Phi$ or $\beta = 0$, let

$$M := \bigoplus_c L_{\beta+c\alpha}$$

where the sum is over all $c \in \mathbb{C}$ such that $\beta + c\alpha \in \Phi$. M is an $\mathfrak{sl}(\alpha)$ -submodule of L . This module is said to be the α -*root string through* $m\beta$.

Proof. For each $c_1, c_2 \in \mathbb{C}$. If $\beta \neq 0$, then $[L_{\beta+c_1\alpha}, L_{\beta+c_2\alpha}] \in L_{\beta+(\beta^{-1}\alpha+c_1+c_2)\alpha}$. If $\beta = 0$, then $[L_{\beta+c_1\alpha}, L_{\beta+c_2\alpha}] \in L_{\beta+(c_1+c_2)\alpha}$. $\square$

Proposition 8.4.6

Let $\alpha \in \Phi$. The root spaces $L_{\pm\alpha}$ are 1-dimensional. Moreover, the only multiples of α which lie in Φ are $\pm\alpha$.

Low Dimension Lie Algebras

9.1 Dimension 3

9.1.1 Lie Algebras where $L' = L$

Theorem 9.1.1

Suppose that L is a complex Lie algebra of dimension 3 such that $L = L'$. Then L is isomorphic to $\mathfrak{sl}(2, \mathbb{C})$.

Proof. 1. Let $x \in L$ be non-zero. We claim that $\text{ad } x$ has rank 2. Extend x to a basis of L say $\{x, y, z\}$. Then L' is spanned by

$$\{[x, y], [x, z], [y, z]\}$$

But $L' = L$, this set must be linearly independent, and hence the image of $\text{ad } x$ has a basis $\{[x, y], [x, z]\}$ of size 2, as required.

2. We claim that there is some $h \in L$ such that $\text{ad } h$ has an eigenvector with a non-zero eigenvalue. Choose any non-zero $x \in L$. If $\text{ad } x$ has a non-zero eigenvalue, then we take $h = x$. If $\text{ad } x$ has no non-zero eigenvalues, then, as it has rank 2, its Jordan canonical form must be

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

This matrix indicates there is a basis of L extending $\{x\}$, say $\{x, y, z\}$ such that $[x, y] = x$ and $[x, z] = y$. So $\text{ad } y$ has x as an eigenvector with eigenvalue -1 , and we may take $h = y$.

3. By the previous step, we may find $h, x \in L$ such that $[h, x] = \alpha x \neq 0$. Since $h \in L$ and $L = L'$, we know $\text{ad } h$ has trace zero. It follows that $\text{ad } h$ must have three distinct eigenvalues $\alpha, 0, -\alpha$. If y is an eigenvector for $\text{ad } h$ with eigenvalue $-\alpha$, then $\{h, x, y\}$ is a basis of L . In this basis, $\text{ad } h$ is represented by a diagonal matrix.

4. To determine $[x, y]$. Note that

$$[h, [x, y]] = [[h, x], y] + [x, [h, y]] = \alpha[x, y] + (-\alpha)[x, y] = 0$$

5. We now make two applications of step 1. Firstly, $\ker \operatorname{ad} h = \operatorname{span} \{h\}$, so $[x, y] = \lambda h$. Secondly, $\lambda \neq 0$ otherwise $\ker \operatorname{ad} x$ is 2-dimensional. By replacing x with $\lambda^{-1}x$ we may assume that $\lambda = 1$.

□