Riemannian Geometry

for Dummies

Colin Roberts

Colorado State University

Section 1

Introduction

Riemannian geometry is the study of a ***smooth manifold*** M along with a ***Riemannian metric*** g.

The point of Riemmannian geometry is to generalize the differentiable and metric structure of $\mathbb{R}^n$.

We generalize to spaces that have interesting topology and geometry.

This will require us to rethink some notions we found “easy" in $\mathbb{R}^n$.

But we will gain a very general framework for working with differentiable objects.

Section 2

Motivation

Why study this in the first place?

Example: Partial differential equations (PDEs) on spaces that are not flat.

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;
- Electrical Impedence Tomography (EIT);

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;
- Electrical Impedence Tomography (EIT);
- General relativity.

Example: Optimization in interesting spaces.

Example: Optimization in interesting spaces.

- Grassmannians;

Example: Optimization in interesting spaces.

- Grassmannians;
- Flags.

Section 3

Preliminaries

add more math text before/after pics so that people see some notation. More examples.

Subsection 1

Smooth Manifolds

<u>Our To-Do List:</u>

- Start with a topological space M;

Our To-Do List:

- Start with a topological space M;
- Look at open sets U that cover M;

Our To-Do List:

- Start with a topological space M;
- Look at open sets U that cover M;
- Construct local coordinates φ;

Our To-Do List:

- Start with a topological space M;
- Look at open sets U that cover M;
- Construct local coordinates φ;
- Show coordinate transition functions are smooth.

Define the sphere as the set of points in $\mathbb{R}^3$... then say we'll mostly use this as an example so keep it in mind


M


M
U_α
U_β
$U_\alpha \cap U_\beta$



M
φ_α
$\mathbb{R}^m$
$\varphi_{\alpha\beta} = \varphi_\alpha^{-1} \circ \varphi_\beta$
$\mathbb{R}^m$
φ_β

Subsection 2

Vector Fields



M
$\gamma(0) = p$
γ


T_pM
M
p
$\frac{d\gamma}{dt}\Big|_{t=0}$
γ



$d\varphi(x)$
T_pM
$d\varphi(x)e_2$
$T_x\mathbb{R}^m$
x_2
e_2
M
$\varphi(x) = p$
$d\varphi(x)e_1$
e_1
x
x_1
φ


T_qM
T_rM
q
r
M
T_pM
p


T_rM
T_pM
T_qM
M
$\rightarrow \bigcupdot_{p\in M} T_pM \rightarrow$
TM



X_q
X_r
M
X_p
TM
X
r
q
p

Subsection 3

Example

Example of charts and maps for sphere as well as a vector field. Do spherical coordinates


z
M
p
ϕ
y
θ
x
$\varphi(\theta, \phi) = (\sin\theta\cos\phi, \sin\theta\sin\phi, \cos\theta)$
ϕ
π
θ
2π



z
θ = const.
ϕ = const.
$d\varphi(\theta, \phi)$
p
$\hat{\theta}$
$\hat{\phi}$
y
x
ϕ
θ

h

Section 4

Riemannian Geometry

Our To-Do List:

- Build an inner product on the tangent space T_pM;

Our To-Do List:

- Build an inner product on the tangent space $T_p M$;
- Have the inner product vary smoothly as we vary the point p;

<u>Our To-Do List:</u>

- Build an inner product on the tangent space T_pM;
- Have the inner product vary smoothly as we vary the point p;
- Define this as our Riemannian metric tensor field g;

Our To-Do List:

- Build an inner product on the tangent space T_pM;
- Have the inner product vary smoothly as we vary the point p;
- Define this as our Riemannian metric tensor field g;
- Extract geometrical and analytical qualities of the underlying manifold M.

Subsection 1

Riemannian Metric

$$g_{ij}(x) = \varphi^*(x)e_i \cdot \varphi^*(x)e_k = (\partial_i\varphi) \cdot (\partial_j\varphi) = \sum_{k=1}^{m} \frac{\partial\varphi_k}{\partial x_i}\frac{\partial\varphi_k}{\partial x_j}$$

connection covariant derivative interpretation, covariant derivative in spherical coordinates,second covariant matrix and covariant laplacian from hessian? Compatability with riemannian metric picture. Show a geodesic in the coordinates between cities or something?

From minimization of length/energy. Both are good to mention. Geodesic equation

$$\nabla_{\dot{\gamma}}\dot{\gamma} = 0$$

equivalent to

$$\ddot{x}^l + \dot{x}^j \dot{x}^k \Gamma^l_{jk} = 0$$

which is saying that the only "acceleration" of the curve comes from the geometry it lies on. When flat space, $\Gamma^l_{jk} = 0$ and we have $\ddot{x} = 0$.

Section 5

Applications

navier stokes on the Earth. Riemannian metric as a conductivity?

Section 6

Conclusions