

Clifford Analysis and a Noncommutative Gelfand Representation

Colin Roberts

Overview

- 1 Introduction
- 2 Clifford analysis
- 3 Gelfand theory
- 4 Future work
- 5 Conclusions

Section 1

Introduction

Motivating problems

Motivating problems

- *Electrical Impedance Tomography (EIT)* asks whether one can determine the conductivity of a medium from the voltage-to-current map.

Motivating problems

- *Electrical Impedance Tomography (EIT)* asks whether one can determine the conductivity of a medium from the voltage-to-current map.
- The *Calderón problem* replaces the medium with a manifold M , conductivity with g , and replaces the voltage-to-current map with the Dirichlet-to-Neumann operator Λ .

Other questions

Other questions

- What topological information can we retrieve from functions on a manifold M ?

Other questions

- What topological information can we retrieve from functions on a manifold M ?
- Do these functions also contain geometric information such as metric data?

Other questions

- What topological information can we retrieve from functions on a manifold M ?
- Do these functions also contain geometric information such as metric data?
- How much can we learn about M if our data is supported only on the boundary?

Subsection 1

Preliminaries

- *Clifford algebra* originated in 1878 with William Kingdon Clifford's work that extends Hermann Grassmann's *exterior algebra*.

- *Clifford algebra* originated in 1878 with William Kingdon Clifford's work that extends Hermann Grassmann's *exterior algebra*.
- *Clifford analysis* arrived in the 1980's due to Hestenes, Sobczyk, Sommen, Brackx, and Delenghe in order to enrich Élie Cartan's *differential forms*. See: [Hestenes, Sobczyk: 1984] and [Doran, Lasenby: 2003].

Clifford algebras

Clifford algebras

Let V be a vector space over a field \mathbb{F} with quadratic form Q .

Clifford algebras

Let V be a vector space over a field \mathbb{F} with quadratic form Q .

- Define the tensor algebra

$$\mathcal{T}(V) := \bigoplus_{j=0}^{\infty} V^{\otimes j} = \mathbb{F} \oplus (V \otimes V) \oplus (V \otimes V \otimes V) \oplus \dots.$$

Clifford algebras

Let V be a vector space over a field \mathbb{F} with quadratic form Q .

- Define the tensor algebra

$$\mathcal{T}(V) := \bigoplus_{j=0}^{\infty} V^{\otimes j} = \mathbb{F} \oplus (V \otimes V) \oplus (V \otimes V \otimes V) \oplus \dots.$$

- Form the *Clifford algebra* via a quotient

$$Cl(V, Q) := \mathcal{T}(V) / \langle \mathbf{v} \otimes \mathbf{v} - Q(\mathbf{v}) \rangle.$$

Geometric and exterior algebras

Geometric and exterior algebras

- Given a (pseudo) inner product g , we set $Q(\cdot) = g(\cdot, \cdot)$ and define a *geometric algebra*

$$\mathcal{G} := Cl(V, g).$$

Geometric and exterior algebras

- Given a (pseudo) inner product g , we set $Q(\cdot) = g(\cdot, \cdot)$ and define a *geometric algebra*

$$\mathcal{G} := Cl(V, g).$$

- The *exterior algebra* is given by

$$\bigwedge(V) := Cl(V, 0).$$

Algebraic structure

Algebraic structure

Multiplication in \mathcal{G} is seen by looking at how \otimes acts in the quotient.

Algebraic structure

Multiplication in \mathcal{G} is seen by looking at how \otimes acts in the quotient.

- Given $\mathbf{u}, \mathbf{v} \in \mathcal{G}$ we can take the product

$$\mathbf{u}\mathbf{v} = \underbrace{\mathbf{u} \cdot \mathbf{v}}_{\text{scalar}} + \underbrace{\mathbf{u} \wedge \mathbf{v}}_{\text{bivector}} .$$

Algebraic structure

Multiplication in \mathcal{G} is seen by looking at how \otimes acts in the quotient.

- Given $\mathbf{u}, \mathbf{v} \in \mathcal{G}$ we can take the product

$$\mathbf{u}\mathbf{v} = \underbrace{\mathbf{u} \cdot \mathbf{v}}_{\text{scalar}} + \underbrace{\mathbf{u} \wedge \mathbf{v}}_{\text{bivector}}.$$

- The scalar part is symmetric: $\mathbf{u} \cdot \mathbf{v} = g(\mathbf{u}, \mathbf{v})$.

Algebraic structure

Multiplication in \mathcal{G} is seen by looking at how \otimes acts in the quotient.

- Given $\mathbf{u}, \mathbf{v} \in \mathcal{G}$ we can take the product

$$\mathbf{u}\mathbf{v} = \underbrace{\mathbf{u} \cdot \mathbf{v}}_{\text{scalar}} + \underbrace{\mathbf{u} \wedge \mathbf{v}}_{\text{bivector}}.$$

- The scalar part is symmetric: $\mathbf{u} \cdot \mathbf{v} = g(\mathbf{u}, \mathbf{v})$.
- The bivector part is antisymmetric: $\mathbf{u} \wedge \mathbf{v} = -\mathbf{v} \wedge \mathbf{u}$.

Multivectors

Multivectors

- \mathcal{G} is graded and of dimension 2^n .

Multivectors

- \mathcal{G} is graded and of dimension 2^n .
 - There are $\binom{n}{r}$ elements in the space of grade- r elements, \mathcal{G}^r , called *r -vectors*.

Multivectors

- \mathcal{G} is graded and of dimension 2^n .
 - There are $\binom{n}{r}$ elements in the space of grade- r elements, \mathcal{G}^r , called *r -vectors*.
 - Those that are exterior products of r independent vectors are *r -blades*.
E.g., $\mathbf{A}_r = \mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_r$.

Multivectors

- \mathcal{G} is graded and of dimension 2^n .
 - There are $\binom{n}{r}$ elements in the space of grade- r elements, \mathcal{G}^r , called *r -vectors*.
 - Those that are exterior products of r independent vectors are *r -blades*.
E.g., $\mathbf{A}_r = \mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_r$.
 - Elements of the even grade subalgebra, \mathcal{G}^+ , are called *spinors*.

Multivectors

- \mathcal{G} is graded and of dimension 2^n .
 - There are $\binom{n}{r}$ elements in the space of grade- r elements, \mathcal{G}^r , called *r -vectors*.
 - Those that are exterior products of r independent vectors are *r -blades*.
E.g., $\mathbf{A}_r = \mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_r$.
 - Elements of the even grade subalgebra, \mathcal{G}^+ , are called *spinors*.
- The most general elements are *multivectors* and are given by

$$A = \sum_{r=0}^n \langle A \rangle_r,$$

where $\langle A \rangle_r \in \mathcal{G}^r$ extracts the grade r part of A . So $\mathcal{G} = \bigoplus_{r=0}^n \mathcal{G}^r$.

Algebraic Structure

Algebraic Structure

- Extend the multiplication from vectors to multivectors.

Algebraic Structure

- Extend the multiplication from vectors to multivectors.
- On homogeneous elements,

$$A_r B_s = \langle A_r B_s \rangle_{|r-s|} + \langle A_r B_s \rangle_{|r-s|+2} + \cdots + \langle A_r B_s \rangle_{r+s}$$

Algebraic Structure

- Extend the multiplication from vectors to multivectors.
- On homogeneous elements,

$$A_r B_s = \langle A_r B_s \rangle_{|r-s|} + \langle A_r B_s \rangle_{|r-s|+2} + \cdots + \langle A_r B_s \rangle_{r+s}$$

- The most important products are

$$A_r \cdot B_s := \langle A_r B_s \rangle_{|r-s|}$$

$$A_r \wedge B_s := \langle A_r B_s \rangle_{r+s}$$

$$A_r \rfloor B_s := \langle A_r B_s \rangle_{s-r}$$

$$A_r \lrcorner B_s := \langle A_r B_s \rangle_{r-s}$$

Reciprocals and reverses

Reciprocals and reverses

- Given any vector basis \mathbf{v}_i , define the *reciprocal vectors* by $\mathbf{v}^i \cdot \mathbf{v}_j = \delta_j^i$.

Reciprocals and reverses

- Given any vector basis \mathbf{v}_i , define the *reciprocal vectors* by $\mathbf{v}^i \cdot \mathbf{v}_j = \delta_j^i$.
- The *reverse* of a multivector is extended linearly from the action on r -blades by

$$\mathbf{A}_r^\dagger = (\mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_r)^\dagger = \mathbf{v}_r \wedge \cdots \wedge \mathbf{v}_1.$$

Inner product and norm

Inner product and norm

- Define the *multivector inner product* by

$$(A, B) := \langle A^\dagger B \rangle$$

which is bilinear, symmetric, and positive definite if g is positive definite.

Inner product and norm

- Define the *multivector inner product* by

$$(A, B) := \langle A^\dagger B \rangle$$

which is bilinear, symmetric, and positive definite if g is positive definite.

- Define the *multivector norm* by

$$|A| := \sqrt{(A, A)}.$$

Adjoint

Note the reverse acts as an adjoint by

$$(CA, B) = (A, C^\dagger B)$$

$$(AC, B) = (A, BC^\dagger).$$

Pseudoscalars

Pseudoscalars

- *Pseudoscalars* are the grade- n elements. For example, the volume element

$$\boldsymbol{\mu} = \mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_n.$$

Pseudoscalars

- *Pseudoscalars* are the grade- n elements. For example, the volume element

$$\boldsymbol{\mu} = \mathbf{v}_1 \wedge \cdots \wedge \mathbf{v}_n.$$

- We define the *unit pseudoscalar* by

$$\boldsymbol{I} := \frac{1}{|\boldsymbol{\mu}|} \boldsymbol{\mu}.$$

Blades and subspaces

Blades and subspaces

- If g is positive definite all blades are invertible [Chisholm: 2012].

Blades and subspaces

- If g is positive definite all blades are invertible [Chisholm: 2012].
- If $|\mathbf{A}_r| = 1$, then \mathbf{A}_r is a *unit blade*.

Blades and subspaces

- If g is positive definite all blades are invertible [Chisholm: 2012].
- If $|\mathbf{A}_r| = 1$, then \mathbf{A}_r is a *unit blade*.
- Unit r -blades correspond to r -dimensional subspaces so they correspond to points in $\text{Gr}(r, n)$.

Duality

Duality

- Given any multivector A , we can take its *dual*

$$A^\perp := A\boldsymbol{I}^{-1}.$$

Duality

- Given any multivector A , we can take its *dual*

$$A^\perp := A\mathbf{I}^{-1}.$$

- Note $A_r^\perp \in \mathcal{G}^{n-r}$, like the Hodge star \star .

Projection and rejection

Projection and rejection

- The *projection* of B into a subspace \mathbf{A}_r by

$$P_{\mathbf{A}_r}(B) := B \rfloor \mathbf{A}_r \mathbf{A}_r^{-1}$$

Projection and rejection

- The *projection* of B into a subspace \mathbf{A}_r by

$$P_{\mathbf{A}_r}(B) := B \rfloor \mathbf{A}_r \mathbf{A}_r^{-1}$$

- The *rejection* by

$$R_{\mathbf{A}_r}(B) := B \wedge \mathbf{A}_r \mathbf{A}_r^{-1}.$$

Projection and rejection

- The *projection* of B into a subspace \mathbf{A}_r by

$$P_{\mathbf{A}_r}(B) := B \rfloor \mathbf{A}_r \mathbf{A}_r^{-1}$$

- The *rejection* by

$$R_{\mathbf{A}_r}(B) := B \wedge \mathbf{A}_r \mathbf{A}_r^{-1}.$$

- Both are grade preserving.

Examples

Examples

- Define $\mathcal{G}_{p,q}$ by letting $\mathbf{e}_i^2 = -1$ for $i = 1, \dots, p$ and $\mathbf{e}_i^2 = +1$ otherwise.

Examples

- Define $\mathcal{G}_{p,q}$ by letting $\mathbf{e}_i^2 = -1$ for $i = 1, \dots, p$ and $\mathbf{e}_i^2 = +1$ otherwise.
- Claim: \mathbb{H} arises naturally as the even subalgebra $\mathcal{G}_3^+ := \mathcal{G}_{0,3}^+$.

Examples

- Define $\mathcal{G}_{p,q}$ by letting $\mathbf{e}_i^2 = -1$ for $i = 1, \dots, p$ and $\mathbf{e}_i^2 = +1$ otherwise.
- Claim: \mathbb{H} arises naturally as the even subalgebra $\mathcal{G}_3^+ := \mathcal{G}_{0,3}^+$.
- Claim: \mathbb{C} arises naturally as the even subalgebra $\mathcal{G}_2^+ := \mathcal{G}_{0,2}^+$.

Examples

- Define $\mathcal{G}_{p,q}$ by letting $\mathbf{e}_i^2 = -1$ for $i = 1, \dots, p$ and $\mathbf{e}_i^2 = +1$ otherwise.
- **Claim:** \mathbb{H} arises naturally as the even subalgebra $\mathcal{G}_3^+ := \mathcal{G}_{0,3}^+$.
- **Claim:** \mathbb{C} arises naturally as the even subalgebra $\mathcal{G}_2^+ := \mathcal{G}_{0,2}^+$.
 - Take the standard basis $\mathbf{e}_1, \mathbf{e}_2$, and define $\mathbf{B}_{12} = \mathbf{e}_1 \mathbf{e}_2$ and note $\mathbf{B}_{12}^2 = -1$.
Thus,

$$(u_1 + v_1 \mathbf{B}_{12})(u_2 + v_2 \mathbf{B}_{12}) = u_1 u_2 - v_1 v_2 + (u_1 v_2 + u_2 v_1) \mathbf{B}_{12}.$$

Examples

- Define $\mathcal{G}_{p,q}$ by letting $\mathbf{e}_i^2 = -1$ for $i = 1, \dots, p$ and $\mathbf{e}_i^2 = +1$ otherwise.
- **Claim:** \mathbb{H} arises naturally as the even subalgebra $\mathcal{G}_3^+ := \mathcal{G}_{0,3}^+$.
- **Claim:** \mathbb{C} arises naturally as the even subalgebra $\mathcal{G}_2^+ := \mathcal{G}_{0,2}^+$.
 - Take the standard basis $\mathbf{e}_1, \mathbf{e}_2$, and define $\mathbf{B}_{12} = \mathbf{e}_1 \mathbf{e}_2$ and note $\mathbf{B}_{12}^2 = -1$.
Thus,

$$(u_1 + v_1 \mathbf{B}_{12})(u_2 + v_2 \mathbf{B}_{12}) = u_1 u_2 - v_1 v_2 + (u_1 v_2 + u_2 v_1) \mathbf{B}_{12}.$$

- Right multiplication by \mathbf{B}_{12} rotates counter-clockwise by $\pi/2$.

Section 2

Clifford analysis

Multivector Fields

Multivector Fields

- Let M be a smooth, compact, connected, and oriented n -dimensional Riemannian manifold with metric g .

Multivector Fields

- Let M be a smooth, compact, connected, and oriented n -dimensional Riemannian manifold with metric g .
- Idea: Form the Clifford algebras on tangent spaces.

Multivector Fields

- Let M be a smooth, compact, connected, and oriented n -dimensional Riemannian manifold with metric g .
- Idea: Form the Clifford algebras on tangent spaces.
 - Each $Cl(T_p M, g_p)$ is a *geometric tangent space* which we glue together to form

$$Cl(TM, g) := \bigsqcup_{p \in M} Cl(T_p M, g_p).$$

Multivector Fields

- Let M be a smooth, compact, connected, and oriented n -dimensional Riemannian manifold with metric g .
- Idea: Form the Clifford algebras on tangent spaces.
 - Each $Cl(T_p M, g_p)$ is a *geometric tangent space* which we glue together to form

$$Cl(TM, g) := \bigsqcup_{p \in M} Cl(T_p M, g_p).$$

- The space of *(smooth) multivector fields* is

$$\mathcal{G}(M) := \{C^\infty\text{-smooth sections of } Cl(TM, g)\}.$$

Multivector Fields

- Let M be a smooth, compact, connected, and oriented n -dimensional Riemannian manifold with metric g .
- Idea: Form the Clifford algebras on tangent spaces.
 - Each $Cl(T_p M, g_p)$ is a *geometric tangent space* which we glue together to form

$$Cl(TM, g) := \bigsqcup_{p \in M} Cl(T_p M, g_p).$$

- The space of *(smooth) multivector fields* is

$$\mathcal{G}(M) := \{C^\infty\text{-smooth sections of } Cl(TM, g)\}.$$

- Retain the same naming scheme as before.

Multivector derivative

Multivector derivative

On M , take the unique Levi-Civita connection ∇ and covariant derivative $\nabla_{\mathbf{u}}$.

Multivector derivative

On M , take the unique Levi-Civita connection ∇ and covariant derivative $\nabla_{\mathbf{u}}$.

- $\nabla_{\mathbf{u}}$ is extended to multivectors and is grade preserving [Schindler: 2018],

$$\nabla_{\mathbf{u}} A_r = \langle \nabla_{\mathbf{u}} A_r \rangle_r.$$

Multivector derivative

On M , take the unique Levi-Civita connection ∇ and covariant derivative $\nabla_{\mathbf{u}}$.

- $\nabla_{\mathbf{u}}$ is extended to multivectors and is grade preserving [Schindler: 2018],

$$\nabla_{\mathbf{u}} A_r = \langle \nabla_{\mathbf{u}} A_r \rangle_r.$$

- $\nabla_{\mathbf{u}}$ is compatible with dot and wedge since

$$\nabla_{\mathbf{u}}(A \cdot B) = (\nabla_{\mathbf{u}} A) \cdot B + A \cdot (\nabla_{\mathbf{u}} B)$$

$$\nabla_{\mathbf{u}}(A \wedge B) = (\nabla_{\mathbf{u}} A) \wedge B + A \wedge (\nabla_{\mathbf{u}} B).$$

Gradient

Gradient

- Define the *gradient* (or *Dirac operator*) locally by

$$\nabla = \sum_{i=1}^n \mathbf{v}^i \nabla_{\mathbf{v}_i}$$

Gradient

- Define the *gradient* (or *Dirac operator*) locally by

$$\nabla = \sum_{i=1}^n \mathbf{v}^i \nabla_{\mathbf{v}_i}$$

- ∇ acts as a vector in $\mathcal{G}(M)$ and obeys the Leibniz rule

$$\nabla(AB) = \dot{\nabla} \dot{A} B + \dot{\nabla} A \dot{B}.$$

Gradient

- Define the *gradient* (or *Dirac operator*) locally by

$$\nabla = \sum_{i=1}^n \mathbf{v}^i \nabla_{\mathbf{v}_i}$$

- ∇ acts as a vector in $\mathcal{G}(M)$ and obeys the Leibniz rule

$$\nabla(AB) = \dot{\nabla} \dot{A} B + \dot{\nabla} A \dot{B}.$$

- Note $\nabla^2 = \Delta$, the Laplace-Beltrami operator.

Example

Example

- In $\mathcal{G}_3(\mathbb{R}^3)$, $\nabla_{\mathbf{e}_i}$ is the partial derivative.

Example

- In $\mathcal{G}_3(\mathbb{R}^3)$, $\nabla_{\mathbf{e}_i}$ is the partial derivative.
- Take a vector field \mathbf{v} , then

$$\nabla \mathbf{v} = \underbrace{\nabla \cdot \mathbf{v}}_{\text{divergence}} + \underbrace{\nabla \wedge \mathbf{v}}_{\text{curl}}.$$

Example

- In $\mathcal{G}_3(\mathbb{R}^3)$, $\nabla_{\mathbf{e}_i}$ is the partial derivative.
- Take a vector field \mathbf{v} , then

$$\nabla \mathbf{v} = \underbrace{\nabla \cdot \mathbf{v}}_{\text{divergence}} + \underbrace{\nabla \wedge \mathbf{v}}_{\text{curl}}.$$

- Specifically,

$$\text{curl}(\mathbf{v}) = (\nabla \wedge \mathbf{v})^\perp$$

Differential forms

Differential forms

- Define the *r-dimensional directed measure*

$$dX_r := \mathbf{v}_{j_1} \wedge \cdots \wedge \mathbf{v}_{j_r} dx^{j_1} \cdots dx^{j_r}$$

where $1 \leq j_1 < \cdots < j_r \leq n$ and summation is implied.

Differential forms

- Define the *r-dimensional directed measure*

$$dX_r := \mathbf{v}_{j_1} \wedge \cdots \wedge \mathbf{v}_{j_r} dx^{j_1} \cdots dx^{j_r}$$

where $1 \leq j_1 < \cdots < j_r \leq n$ and summation is implied.

- Define an *r*-form a_r by

$$\alpha_r = A_r \cdot dX_r^\dagger$$

where $A_r = \frac{1}{r!} a_{i_1 \dots i_r} \mathbf{v}^{i_1} \wedge \cdots \wedge \mathbf{v}^{i_r}$.

Differential forms

- Define the *r-dimensional directed measure*

$$dX_r := \mathbf{v}_{j_1} \wedge \cdots \wedge \mathbf{v}_{j_r} dx^{j_1} \cdots dx^{j_r}$$

where $1 \leq j_1 < \cdots < j_r \leq n$ and summation is implied.

- Define an *r*-form a_r by

$$\alpha_r = A_r \cdot dX_r^\dagger$$

where $A_r = \frac{1}{r!} a_{i_1 \dots i_r} \mathbf{v}^{i_1} \wedge \cdots \wedge \mathbf{v}^{i_r}$.

- Refer to A_r the *multivector equivalent* of α_r .

Exterior algebra and calculus

Exterior algebra and calculus

- Given r -forms a_r , b_r , and an s -form c_s , we have

$$a_r + b_r = (A_r + B_r) \cdot dX_r^\dagger, \quad a_r \wedge c_s = (A_r \wedge C_s) \cdot dX_{r+s}^\dagger.$$

Exterior algebra and calculus

- Given r -forms a_r , b_r , and an s -form c_s , we have

$$a_r + b_r = (A_r + B_r) \cdot dX_r^\dagger, \quad a_r \wedge c_s = (A_r \wedge C_s) \cdot dX_{r+s}^\dagger.$$

- The exterior derivative on multivector equivalents is

$$da_r = (\nabla \wedge A_r) \cdot dX_{r+1}^\dagger$$

Exterior algebra and calculus

- Given r -forms a_r , b_r , and an s -form c_s , we have

$$a_r + b_r = (A_r + B_r) \cdot dX_r^\dagger, \quad a_r \wedge c_s = (A_r \wedge C_s) \cdot dX_{r+s}^\dagger.$$

- The exterior derivative on multivector equivalents is

$$da_r = (\nabla \wedge A_r) \cdot dX_{r+1}^\dagger$$

- The Hodge star on multivector equivalents is

$$\star a_r = (I^{-1} A_r)^\dagger \cdot dX_{n-r}^\dagger$$

Volume form

Volume form

- The *volume form* on M is given in local coordinates by

$$\mu = \sqrt{|g|} dx^1 \cdots dx^n = \mathbf{I}^{-1} \cdot dX_n$$

Volume form

- The *volume form* on M is given in local coordinates by

$$\mu = \sqrt{|g|} dx^1 \cdots dx^n = \mathbf{I}^{-1} \cdot dX_n$$

- We integrate scalar fields A_0 on M by

$$\int_M A_0^\perp \cdot dX_n = \int_M A_0 \mu.$$

Multivector field inner product

Multivector field inner product

- We define the L^2 -inner product on multivector fields by

$$\langle\langle A, B \rangle\rangle := \frac{1}{\text{vol}(M)} \int_M (A, B) \mu$$

Multivector field inner product

- We define the L^2 -inner product on multivector fields by

$$\ll A, B \gg := \frac{1}{\text{vol}(M)} \int_M (A, B) \mu$$

- This realizes the r -form inner product

$$\int_M \alpha_r \wedge \star \beta_r = \int_M \langle A_r^\dagger B_r \rangle \mu = \text{vol}(M) \ll A, B \gg$$

Multivector field inner product

- We define the L^2 -inner product on multivector fields by

$$\ll A, B \gg := \frac{1}{\text{vol}(M)} \int_M (A, B) \mu$$

- This realizes the r -form inner product

$$\int_M \alpha_r \wedge \star \beta_r = \int_M \langle A_r^\dagger B_r \rangle \mu = \text{vol}(M) \ll A, B \gg$$

- $\ll A_r, B_s \gg = 0$ when $r \neq s$ so the L^2 -direct sum agrees with the grade based direct sum.

Boundary

Boundary

- On ∂M , the boundary pseudoscalar $\boldsymbol{I}_{\partial}$ induces the boundary normal

$$\boldsymbol{\nu} = \boldsymbol{I}_{\partial}^{\perp}.$$

Boundary

- On ∂M , the boundary pseudoscalar \mathbf{I}_∂ induces the boundary normal

$$\boldsymbol{\nu} = \mathbf{I}_\partial^\perp.$$

- The boundary volume form is

$$\mu_\partial := \mathbf{I}_\partial^{-1} \cdot dX_{n-1}$$

and we define

$$\ll A, B \gg_\partial := \frac{1}{\text{vol}(M)} \int_{\partial M} (A, B) \mu_\partial.$$

Multivector valued integrals

We can define a multivector valued integral on an oriented submanifold R by taking $A \in \mathcal{G}(M)$ and computing

$$\int_R A \mathbf{I}_R \mu_R.$$

Theorem (Hestenes, Sobczyk, 1984)

Let $A, B \in \mathcal{G}(M)$, then

$$\begin{aligned}\int_M \dot{A} \dot{\nabla} \mathbf{I} \mu &= \int_{\partial M} A \mathbf{I} \partial \mu \partial \\ \int_M \mathbf{I} \nabla B \mu &= \int_{\partial M} \mathbf{I} \partial B \mu \partial \\ \int_M \dot{A} \dot{\nabla} \mathbf{I} B \mu &= (-1)^n \int_M A \mathbf{I} \nabla B \mu + \int_{\partial M} A \mathbf{I} \partial B \mu \partial.\end{aligned}$$

Theorem

We have the Green's formula for the gradient

$$\langle\langle A, \mathbf{I} \nabla B \rangle\rangle = (-1)^n \langle\langle \nabla A, \mathbf{I} B \rangle\rangle + \langle\langle A, \mathbf{I} \partial B \rangle\rangle_{\partial} .$$

Special fields

Special fields

- Define the *monogenic fields*

$$\mathcal{M}(M) := \{A \in \mathcal{G}(M) \mid \nabla A = 0\}.$$

Special fields

- Define the *monogenic fields*

$$\mathcal{M}(M) := \{A \in \mathcal{G}(M) \mid \nabla A = 0\}.$$

- Let $f = u + v\mathbf{B} \in \mathcal{G}_2^+(\mathbb{R}^2)$ then $\nabla f = 0$ yields the Cauchy-Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y},$$

$$\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Special fields

- Define the *monogenic fields*

$$\mathcal{M}(M) := \{A \in \mathcal{G}(M) \mid \nabla A = 0\}.$$

- Let $f = u + v\mathbf{B} \in \mathcal{G}_2^+(\mathbb{R}^2)$ then $\nabla f = 0$ yields the Cauchy-Riemann equations

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \qquad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

- Define the *gradients*

$$\nabla \mathcal{G}(M) := \{\nabla A \mid A \in \mathcal{G}(M) \text{ and } A|_{\partial M} = 0\}.$$

Cauchy kernel

Cauchy kernel

- For the remainder, take M imbedded in \mathbb{R}^n with $n \geq 2$.

Cauchy kernel

- For the remainder, take M imbedded in \mathbb{R}^n with $n \geq 2$.
- Define the vector field

$$E(x) := \frac{1}{S_n} \frac{x}{|x|^n}$$

where S_n is the surface area of the unit ball.

Cauchy kernel

- For the remainder, take M imbedded in \mathbb{R}^n with $n \geq 2$.
- Define the vector field

$$E(x) := \frac{1}{S_n} \frac{x}{|x|^n}$$

where S_n is the surface area of the unit ball.

- Note,

$$\nabla E(x) = -\dot{E}(x)\dot{\nabla} = \delta(x).$$

Cauchy kernel

- For the remainder, take M imbedded in \mathbb{R}^n with $n \geq 2$.
- Define the vector field

$$E(x) := \frac{1}{S_n} \frac{x}{|x|^n}$$

where S_n is the surface area of the unit ball.

- Note,

$$\nabla E(x) = -\dot{E}(x)\dot{\nabla} = \delta(x).$$

- Define the *Cauchy kernel* by $G(x, x') := E(x' - x)$.

Cauchy integral

Cauchy integral

- Let $A \in \mathcal{M}(M)$, then we have the *Cauchy integral formula*

$$A(x) = (-1)^n \mathbf{I}^{-1} \int_{\partial M} G(x, x') \mathbf{I}_{\partial}(x') A(x') \mu_{\partial}(x').$$

Cauchy integral

- Let $A \in \mathcal{M}(M)$, then we have the *Cauchy integral formula*

$$A(x) = (-1)^n \mathbf{I}^{-1} \int_{\partial M} G(x, x') \mathbf{I}_{\partial}(x') A(x') \mu_{\partial}(x').$$

- This uniquely determines a monogenic field from boundary values.

Lemma

Let $A \in \mathcal{M}(M)$ be such that $A|_{\partial M} = 0$. Then $A = 0$ on all of M .

Lemma

Fix a multivector field $A \in \mathcal{G}(M)$. If

$$\ll A, B \gg = 0$$

for all $B \in \mathcal{G}(M)$ with $B|_{\partial M} = 0$, then $A = 0$.

Theorem (Clifford-Hodge-Morrey Decomposition)

The space of multivector fields $\mathcal{G}(M)$ has the L^2 -orthogonal decomposition

$$\mathcal{G}(M) = \mathcal{M}(M) \oplus \mathbf{I}\nabla\mathcal{G}(M).$$

Proof.

Proof.

■ *Orthogonality:* Let $A \in \mathcal{M}(M)$ and $\mathbf{I}\nabla B \in \mathbf{I}\nabla\mathcal{G}(M)$ and note

$$\ll A, \mathbf{I}\nabla B \gg = (-1)^n \ll \nabla A, \mathbf{I}B \gg + \ll A, \mathbf{I}\partial B \gg = 0,$$

by the multivector Green's formula.

- Let $C \in \mathcal{G}(M)$ be in the orthogonal complement to $\mathbf{I}\nabla\mathcal{G}(M)$.

- Let $C \in \mathcal{G}(M)$ be in the orthogonal complement to $\mathbf{I}\nabla\mathcal{G}(M)$.
- Use the Cauchy integral formula, construct a monogenic field \tilde{C} from $C|_{\partial M}$ and note $C = \tilde{C} + C_0$ where $C_0|_{\partial M} = 0$.

- Let $C \in \mathcal{G}(M)$ be in the orthogonal complement to $\mathbf{I}\nabla\mathcal{G}(M)$.
- Use the Cauchy integral formula, construct a monogenic field \tilde{C} from $C|_{\partial M}$ and note $C = \tilde{C} + C_0$ where $C_0|_{\partial M} = 0$.
- Note,

$$0 = \langle\langle C, \mathbf{I}\nabla B \rangle\rangle = \langle\langle \nabla C_0, \mathbf{I}B \rangle\rangle .$$

- Let $C \in \mathcal{G}(M)$ be in the orthogonal complement to $\mathbf{I}\nabla\mathcal{G}(M)$.
- Use the Cauchy integral formula, construct a monogenic field \tilde{C} from $C|_{\partial M}$ and note $C = \tilde{C} + C_0$ where $C_0|_{\partial M} = 0$.
- Note,

$$0 = \langle\langle C, \mathbf{I}\nabla B \rangle\rangle = \langle\langle \nabla C_0, \mathbf{I}B \rangle\rangle .$$

- By the previous lemmas, it must be that $C_0 = 0$. Hence the orthogonal complement to $\mathbf{I}\nabla\mathcal{G}(M)$ is $\mathcal{M}(M)$.

Comparing to Hodge-Morrey

Comparing to Hodge-Morrey

- The Hodge-Morrey decomposition reads

$$\Omega^r(M) = \underbrace{\mathcal{E}_D^r(M)}_{\text{Im}(\nabla \wedge)} \oplus \underbrace{\mathcal{C}_N^r(M)}_{\text{Im}(\nabla \lrcorner)} \oplus \underbrace{\mathcal{H}^r(M)}_{\text{Ker}(\nabla)}.$$

via [Schwarz: 1995].

Comparing to Hodge-Morrey

- The Hodge-Morrey decomposition reads

$$\Omega^r(M) = \underbrace{\mathcal{E}_D^r(M)}_{\text{Im}(\nabla \wedge)} \oplus \underbrace{\mathcal{C}_N^r(M)}_{\text{Im}(\nabla \lrcorner)} \oplus \underbrace{\mathcal{H}^r(M)}_{\text{Ker}(\nabla)}.$$

via [Schwarz: 1995].

- Whereas the Clifford-Hodge-Morrey decomposition ignores specific grades

$$\mathcal{G}(M) = \mathcal{M}(M) \oplus \mathbf{I} \nabla \mathcal{G}(M).$$

Section 3

Gelfand theory

Open questions

Open questions

- In [Belishev: 2003], we see an algebraic proof for the 2-dimensional Calderón problem.

Open questions

- In [Belishev: 2003], we see an algebraic proof for the 2-dimensional Calderón problem.
- In [Belishev, Vakulenko: 2017], we see a proof for a noncommutative Gelfand representation using quaternion fields for a ball \mathbb{B} in \mathbb{R}^3 .

Open questions

- In [Belishev: 2003], we see an algebraic proof for the 2-dimensional Calderón problem.
- In [Belishev, Vakulenko: 2017], we see a proof for a noncommutative Gelfand representation using quaternion fields for a ball \mathbb{B} in \mathbb{R}^3 .
- Belishev and Vakulenko ask whether this is true in higher dimensions.

Open questions

- In [Belishev: 2003], we see an algebraic proof for the 2-dimensional Calderón problem.
- In [Belishev, Vakulenko: 2017], we see a proof for a noncommutative Gelfand representation using quaternion fields for a ball \mathbb{B} in \mathbb{R}^3 .
- Belishev and Vakulenko ask whether this is true in higher dimensions.
- We prove an analogous result for an arbitrary \mathbb{B} in \mathbb{R}^n .

Open questions

- In [Belishev: 2003], we see an algebraic proof for the 2-dimensional Calderón problem.
- In [Belishev, Vakulenko: 2017], we see a proof for a noncommutative Gelfand representation using quaternion fields for a ball \mathbb{B} in \mathbb{R}^3 .
- Belishev and Vakulenko ask whether this is true in higher dimensions.
- We prove an analogous result for an arbitrary \mathbb{B} in \mathbb{R}^n .
- This approach can hopefully be used to prove the analogous result for any smooth orientable Riemannian manifold M .

2-dimensional BC method

The boundary control (BC) method is implemented in [\[Belishev: 2003\]](#) in the following manner.

2-dimensional BC method

The boundary control (BC) method is implemented in [Belishev: 2003] in the following manner.

- Determine the algebra $\mathcal{A}(M)$ of holomorphic functions on M from continuous function algebra on the boundary $\mathcal{A}(\partial M)$ using Λ .

2-dimensional BC method

The boundary control (BC) method is implemented in [Belishev: 2003] in the following manner.

- Determine the algebra $\mathcal{A}(M)$ of holomorphic functions on M from continuous function algebra on the boundary $\mathcal{A}(\partial M)$ using Λ .
- The classical Gelfand representation shows $\mathcal{A}(M)$ is homeomorphic to M via the weak- $*$ topology.

2-dimensional BC method

The boundary control (BC) method is implemented in [Belishev: 2003] in the following manner.

- Determine the algebra $\mathcal{A}(M)$ of holomorphic functions on M from continuous function algebra on the boundary $\mathcal{A}(\partial M)$ using Λ .
- The classical Gelfand representation shows $\mathcal{A}(M)$ is homeomorphic to M via the weak- $*$ topology.
- Functions in $\mathcal{A}(M)$ determine the complex structure on M .

2-dimensional BC method

The boundary control (BC) method is implemented in [Belishev: 2003] in the following manner.

- Determine the algebra $\mathcal{A}(M)$ of holomorphic functions on M from continuous function algebra on the boundary $\mathcal{A}(\partial M)$ using Λ .
- The classical Gelfand representation shows $\mathcal{A}(M)$ is homeomorphic to M via the weak- $*$ topology.
- Functions in $\mathcal{A}(M)$ determine the complex structure on M .
- Thus, we can find a g that is conformal with the complex structure.

Subsurface spinor fields

Subsurface spinor fields

- Let $\mathbf{B} \in \mathcal{G}(M)$ be a constant unit 2-blade, then $f_+ \in \mathcal{G}^+(M)$ satisfying

$$f_+ = P_{\mathbf{B}} \circ f_+ \circ P_{\mathbf{B}}$$

is a *subsurface spinor field*. Let $\mathcal{G}_{\mathbf{B}}^+(M)$ denote the space such fields.

Subsurface spinor fields

- Let $\mathbf{B} \in \mathcal{G}(M)$ be a constant unit 2-blade, then $f_+ \in \mathcal{G}^+(M)$ satisfying

$$f_+ = P_{\mathbf{B}} \circ f_+ \circ P_{\mathbf{B}}$$

is a *subsurface spinor field*. Let $\mathcal{G}_B^+(M)$ denote the space such fields.

- The space of monogenic subsurface spinors

$$\mathcal{A}_B(M) = \{f_+ \in \mathcal{G}_B^+(M) \mid \nabla f_+ = 0\}$$

is a commutative unital Banach algebra.

Functionals

Functionals

- Define the *spinor dual* $\mathcal{M}^*(M)$ as the continuous right \mathcal{G}_n^+ -module homomorphisms

$$\mathcal{M}^*(M) := \{l: \mathcal{M}^+(M) \rightarrow \mathcal{G}_n^+ \mid l(fs+g) = l(f)s+l(g), \forall f, g \in \mathcal{M}(M), s \in \mathcal{G}_n^+\}$$

and refer to the elements as *spin functionals*.

Functionals

- Define the *spinor dual* $\mathcal{M}^*(M)$ as the continuous right \mathcal{G}_n^+ -module homomorphisms

$$\mathcal{M}^*(M) := \{l: \mathcal{M}^+(M) \rightarrow \mathcal{G}_n^+ \mid l(fs+g) = l(f)s + l(g), \forall f, g \in \mathcal{M}(M), s \in \mathcal{G}_n^+\}$$

and refer to the elements as *spin functionals*.

- Assert the weak-* topology on $\mathcal{M}^*(M)$ so that every $x \in M$ corresponds to a continuous map on $\mathcal{M}^*(M)$.

Characters

Characters

- Define the algebra $\mathbb{A}_{\mathbf{B}}$ to be the algebra generated by 1 and \mathbf{B} .

Characters

- Define the algebra $\mathbb{A}_{\mathbf{B}}$ to be the algebra generated by 1 and \mathbf{B} .
- The *spinor spectrum* $\mathfrak{M}(M)$ is the set of algebra homomorphisms

$$\mathfrak{M}(M) := \{ \delta \in \mathcal{M}^*(M) \mid \delta(f) \in \mathbb{A}_{\mathbf{B}}, \delta(fg) = \delta(f)\delta(g), \forall f, g \in \mathcal{A}_{\mathbf{B}}(M), \mathbf{B} \in \text{Gr}(2, n) \}$$

and refer to the elements as *spin characters*.

Characters

- Define the algebra $\mathbb{A}_{\mathbf{B}}$ to be the algebra generated by 1 and \mathbf{B} .
- The *spinor spectrum* $\mathfrak{M}(M)$ is the set of algebra homomorphisms

$$\mathfrak{M}(M) := \{ \delta \in \mathcal{M}^*(M) \mid \delta(f) \in \mathbb{A}_{\mathbf{B}}, \delta(fg) = \delta(f)\delta(g), \forall f, g \in \mathcal{A}_{\mathbf{B}}(M), \mathbf{B} \in \text{Gr}(2, n) \}$$

and refer to the elements as *spin characters*.

- One example of such characters are point evaluations $\delta(f) = f(x^\delta)$.

Characters

- Define the algebra $\mathbb{A}_{\mathbf{B}}$ to be the algebra generated by 1 and \mathbf{B} .
- The *spinor spectrum* $\mathfrak{M}(M)$ is the set of algebra homomorphisms

$$\mathfrak{M}(M) := \{\delta \in \mathcal{M}^*(M) \mid \delta(f) \in \mathbb{A}_{\mathbf{B}}, \delta(fg) = \delta(f)\delta(g), \forall f, g \in \mathcal{A}_{\mathbf{B}}(M), \mathbf{B} \in \text{Gr}(2, n)\}$$

and refer to the elements as *spin characters*.

- One example of such characters are point evaluations $\delta(f) = f(x^\delta)$.
- We show these are the only elements in the spectrum.

z analogs

z analogs

- Take the standard basis for \mathbb{R}^n , and consider $M = \mathbb{B}_{R,w}$.

z analogs

- Take the standard basis for \mathbb{R}^n , and consider $M = \mathbb{B}_{R,w}$.
- Let $\mathbf{B}_{ij} = \mathbf{e}_i \mathbf{e}_j$, and define

$$z_{ij} = x_j - x_i \mathbf{B}_{ij}$$

z analogs

- Take the standard basis for \mathbb{R}^n , and consider $M = \mathbb{B}_{R,w}$.
- Let $\mathbf{B}_{ij} = \mathbf{e}_i \mathbf{e}_j$, and define

$$z_{ij} = x_j - x_i \mathbf{B}_{ij}$$

- Note $z_{ij} \in \mathcal{A}_{\mathbf{B}_{ij}}(\mathbb{B}_{R,w})$.

Monogenic polynomials

Monogenic polynomials

- Let σ be a permutation of $\{2, 3, \dots, n\}$, then

$$p_{j_2 \dots j_n}(x) = \frac{1}{j!} \sum_{\text{permutations}} z_{1\sigma(1)}(x) \cdots z_{1\sigma(j)}(x)$$

is a monogenic homogeneous polynomial of degree j .

Monogenic polynomials

- Let σ be a permutation of $\{2, 3, \dots, n\}$, then

$$p_{j_2 \dots j_n}(x) = \frac{1}{j!} \sum_{\text{permutations}} z_{1\sigma(1)}(x) \cdots z_{1\sigma(j)}(x)$$

is a monogenic homogeneous polynomial of degree j .

- Collect these into the set of *monogenic polynomials*

$$\mathcal{M}^{\mathcal{P}}(\mathbb{B}_{R,w}) = \left\{ \sum_{j=0}^N \left(\sum_{\substack{j_2 \dots j_n \\ j_2 + \dots + j_n = j}} p_{j_2 \dots j_n} a_{j_2 \dots j_n} \right) \mid j_2 + \dots + j_n = j, N \in \mathbb{N}, a_{j_2 \dots j_n} \in \mathcal{G}_n \right\}.$$

Lemma (Density)

The space $\mathcal{M}^{\mathcal{P}}(\mathbb{B}_{R,w})$ is dense in $\mathcal{M}^+(\mathbb{B}_{R,w})$.

Proof sketch.

- Let $f \in \mathcal{M}^+(\mathbb{B}_{R,w})$ and use the Cauchy integral formula to define the coefficients $a_{j_2 \dots j_n} \in \mathcal{G}_n^+$ by

$$a_{j_2 \dots j_n} = \int_{\partial \mathbb{B}_{R,w}} \frac{\partial^j G(w, y)}{\partial y_2^{j_2} \dots \partial y_n^{j_n}} \nu(y) f(y) \mu_{\partial}(y),$$

Lemma (Density)

The space $\mathcal{M}^{\mathcal{P}}(\mathbb{B}_{R,w})$ is dense in $\mathcal{M}^+(\mathbb{B}_{R,w})$.

Proof sketch.

- Let $f \in \mathcal{M}^+(\mathbb{B}_{R,w})$ and use the Cauchy integral formula to define the coefficients $a_{j_2 \dots j_n} \in \mathcal{G}_n^+$ by

$$a_{j_2 \dots j_n} = \int_{\partial \mathbb{B}_{R,w}} \frac{\partial^j G(w, y)}{\partial y_2^{j_2} \dots \partial y_n^{j_n}} \nu(y) f(y) \mu_{\partial}(y),$$

- Then

$$f(x) = \sum_{j=0}^{\infty} \left(\sum_{\substack{j_2 \dots j_n \\ j_2 + \dots + j_n = j}} p_{j_2 \dots j_n}(x - w) a_{j_2 \dots j_n} \right),$$

converges pointwise for $x \in \mathbb{B}_{R,w}$ by **[Ryan, 2004]**.

Idea

Idea

- By linearity, we can note that for $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$

$$\delta(f(x)) = \sum_{j=0}^{\infty} \left(\sum_{\substack{j_2 \dots j_n \\ j_2 + \dots + j_n = j}} \delta(p_{j_2 \dots j_n}(x - w)) a_{j_2 \dots j_n} \right)$$

Idea

- By linearity, we can note that for $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$

$$\delta(f(x)) = \sum_{j=0}^{\infty} \left(\sum_{\substack{j_2 \dots j_n \\ j_2 + \dots + j_n = j}} \delta(p_{j_2 \dots j_n}(x-w)) a_{j_2 \dots j_n} \right)$$

- On each monogenic polynomial

$$\delta(p_{j_2 \dots j_n}(x)) = \frac{1}{j!} \sum_{\text{permutations}} \delta((z_{1\sigma(1)}(x)) \cdots \delta(z_{1\sigma(j)}(x)))$$

by the multiplicativity of δ .

Lemma (Point evaluation)

Let $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and $z_{ij} \in \mathcal{A}_{\mathbf{B}_{ij}}(\mathbb{B}_{R,w})$, then $\delta(z_{ij}) = z_{ij}(x^\delta)$ for some $x^\delta \in \mathbb{R}^n$.

Lemma (Point evaluation)

Let $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and $z_{ij} \in \mathcal{A}_{\mathbf{B}_{ij}}(\mathbb{B}_{R,w})$, then $\delta(z_{ij}) = z_{ij}(x^\delta)$ for some $x^\delta \in \mathbb{R}^n$.

Proof sketch.

Lemma (Point evaluation)

Let $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and $z_{ij} \in \mathcal{A}_{\mathbf{B}_{ij}}(\mathbb{B}_{R,w})$, then $\delta(z_{ij}) = z_{ij}(x^\delta)$ for some $x^\delta \in \mathbb{R}^n$.

Proof sketch.

- We have $\delta(z_{ij}) = \alpha_{ij} + \beta_{ij}\mathbf{B}_{ij}$.

Lemma (Point evaluation)

Let $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and $z_{ij} \in \mathcal{A}_{\mathbf{B}_{ij}}(\mathbb{B}_{R,w})$, then $\delta(z_{ij}) = z_{ij}(x^\delta)$ for some $x^\delta \in \mathbb{R}^n$.

Proof sketch.

- We have $\delta(z_{ij}) = \alpha_{ij} + \beta_{ij}\mathbf{B}_{ij}$.
- Note $z_{ij}\mathbf{B}_{ji} = -z_{ji}$ and $z_{ij} = z_{kj} + z_{ik}\mathbf{B}_{kj}$ yield the relationships

$$\alpha_{ji} = -\beta_{ij} \quad \alpha_{ij} = \alpha_{kj} \quad \beta_{ij} = \beta_{ik} \quad \alpha_{ik} = -\beta_{kj}.$$

Lemma (Point evaluation)

Let $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and $z_{ij} \in \mathcal{A}_{\mathbf{B}_{ij}}(\mathbb{B}_{R,w})$, then $\delta(z_{ij}) = z_{ij}(x^\delta)$ for some $x^\delta \in \mathbb{R}^n$.

Proof sketch.

- We have $\delta(z_{ij}) = \alpha_{ij} + \beta_{ij}\mathbf{B}_{ij}$.
- Note $z_{ij}\mathbf{B}_{ji} = -z_{ji}$ and $z_{ij} = z_{kj} + z_{ik}\mathbf{B}_{kj}$ yield the relationships

$$\alpha_{ji} = -\beta_{ij} \quad \alpha_{ij} = \alpha_{kj} \quad \beta_{ij} = \beta_{ik} \quad \alpha_{ik} = -\beta_{kj}.$$

- The set of constants α and β are determined by n independent numbers, so we can say $\delta(z_{ij}) = z_{ij}(x^\delta)$ for some $x^\delta \in \mathbb{R}^n$.

Lemma (Identification)

Let $f \in \mathcal{M}(\mathbb{B}_{R,w})$, then $\delta(f) = f(x^\delta)$ for some $x^\delta \in \mathbb{B}_{R,w}$.

Lemma (Identification)

Let $f \in \mathcal{M}(\mathbb{B}_{R,w})$, then $\delta(f) = f(x^\delta)$ for some $x^\delta \in \mathbb{B}_{R,w}$.

Proof.

Lemma (Identification)

Let $f \in \mathcal{M}(\mathbb{B}_{R,w})$, then $\delta(f) = f(x^\delta)$ for some $x^\delta \in \mathbb{B}_{R,w}$.

Proof.

- Fix $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and suppose $x^\delta \notin \mathbb{B}_{R,w}$.

Lemma (Identification)

Let $f \in \mathcal{M}(\mathbb{B}_{R,w})$, then $\delta(f) = f(x^\delta)$ for some $x^\delta \in \mathbb{B}_{R,w}$.

Proof.

- Fix $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and suppose $x^\delta \notin \mathbb{B}_{R,w}$.
- Take a sequence $x_n \rightarrow x^\delta$ with $x_n \notin \mathbb{B}_{R,w}$.

Lemma (Identification)

Let $f \in \mathcal{M}(\mathbb{B}_{R,w})$, then $\delta(f) = f(x^\delta)$ for some $x^\delta \in \mathbb{B}_{R,w}$.

Proof.

- Fix $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and suppose $x^\delta \notin \mathbb{B}_{R,w}$.
- Take a sequence $x_n \rightarrow x^\delta$ with $x_n \notin \mathbb{B}_{R,w}$.
- Define $G_n(x) := G(x, x_n)\mathbf{e}_1 \in \mathcal{M}^+(\mathbb{B}_{R,w})$.

Lemma (Identification)

Let $f \in \mathcal{M}(\mathbb{B}_{R,w})$, then $\delta(f) = f(x^\delta)$ for some $x^\delta \in \mathbb{B}_{R,w}$.

Proof.

- Fix $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and suppose $x^\delta \notin \mathbb{B}_{R,w}$.
- Take a sequence $x_n \rightarrow x^\delta$ with $x_n \notin \mathbb{B}_{R,w}$.
- Define $G_n(x) := G(x, x_n)\mathbf{e}_1 \in \mathcal{M}^+(\mathbb{B}_{R,w})$.
- Note,

$$\lim_{n \rightarrow \infty} \delta(G_n) = \lim_{n \rightarrow \infty} G_n(x^\delta)$$

so this sequence not converge due to a singularity at x^δ .

Lemma (Identification)

Let $f \in \mathcal{M}(\mathbb{B}_{R,w})$, then $\delta(f) = f(x^\delta)$ for some $x^\delta \in \mathbb{B}_{R,w}$.

Proof.

- Fix $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$ and suppose $x^\delta \notin \mathbb{B}_{R,w}$.
- Take a sequence $x_n \rightarrow x^\delta$ with $x_n \notin \mathbb{B}_{R,w}$.
- Define $G_n(x) := G(x, x_n)\mathbf{e}_1 \in \mathcal{M}^+(\mathbb{B}_{R,w})$.
- Note,

$$\lim_{n \rightarrow \infty} \delta(G_n) = \lim_{n \rightarrow \infty} G_n(x^\delta)$$

so this sequence not converge due to a singularity at x^δ .

- Hence, it must be that $x^\delta \in \mathbb{B}_{R,w}$ by continuity of δ .

Theorem (Noncommutative Gelfand representation)

For any $\delta \in \mathfrak{M}(\mathbb{B}_{R,w})$, there is a point $x^\delta \in \mathbb{B}_{R,w}$ such that $\delta(f) = f(x^\delta)$ for any $f \in \mathcal{M}(\mathbb{B}_{R,w})$. Given the weak- $$ topology on $\mathcal{M}^*(\mathbb{B}_{r,w})$, the map*

$$\gamma: \mathfrak{M}(\mathbb{B}_{R,w}) \rightarrow \mathbb{B}_{R,w}, \quad \delta \mapsto x^\delta$$

is a homeomorphism.

Proof.

Proof.

- The lemmas show that $\gamma: \mathfrak{M}(\mathbb{B}_{R,w}) \rightarrow \mathbb{B}_{R,w}$ is bijective.

Proof.

- The lemmas show that $\gamma: \mathfrak{M}(\mathbb{B}_{R,w}) \rightarrow \mathbb{B}_{R,w}$ is bijective.
- To see that γ is a homeomorphism, take a sequence $\delta_n \rightarrow \delta$ in $\mathfrak{M}(\mathbb{B}_{R,w})$.

Proof.

- The lemmas show that $\gamma: \mathfrak{M}(\mathbb{B}_{R,w}) \rightarrow \mathbb{B}_{R,w}$ is bijective.
- To see that γ is a homeomorphism, take a sequence $\delta_n \rightarrow \delta$ in $\mathfrak{M}(\mathbb{B}_{R,w})$.
- For $f \in \mathcal{M}^+(\mathbb{B}_{R,w})$ we have

$$f(\gamma(\delta_n)) = f(x^{\delta_n}) = \delta_n(f) = \gamma^{-1}(x^{\delta_n})(f).$$

Proof.

- The lemmas show that $\gamma: \mathfrak{M}(\mathbb{B}_{R,w}) \rightarrow \mathbb{B}_{R,w}$ is bijective.
- To see that γ is a homeomorphism, take a sequence $\delta_n \rightarrow \delta$ in $\mathfrak{M}(\mathbb{B}_{R,w})$.
- For $f \in \mathcal{M}^+(\mathbb{B}_{R,w})$ we have

$$f(\gamma(\delta_n)) = f(x^{\delta_n}) = \delta_n(f) = \gamma^{-1}(x^{\delta_n})(f).$$

- Taking $n \rightarrow \infty$ shows γ and γ^{-1} are continuous so γ is a homeomorphism.

Section 4

Future work

Calderón problem

Calderón problem

- Question: Let (M, g) be an unknown Riemannian manifold with known boundary ∂M . Consider the forward problem

$$\begin{cases} \Delta\omega = 0 & \text{in } M \\ \iota^*\omega = \phi & \text{on } \partial M \end{cases}$$

Calderón problem

- Question: Let (M, g) be an unknown Riemannian manifold with known boundary ∂M . Consider the forward problem

$$\begin{cases} \Delta\omega = 0 & \text{in } M \\ \iota^*\omega = \phi & \text{on } \partial M \end{cases}$$

- Define the *Dirichlet-to-Neumann map* on forms by $\Lambda\phi = \iota^*(\star d\omega)$.

Calderón problem

- Question: Let (M, g) be an unknown Riemannian manifold with known boundary ∂M . Consider the forward problem

$$\begin{cases} \Delta\omega = 0 & \text{in } M \\ \iota^*\omega = \phi & \text{on } \partial M \end{cases}$$

- Define the *Dirichlet-to-Neumann map* on forms by $\Lambda\phi = \iota^*(\star d\omega)$.
- Can we determine (M, g) from Λ ?

Calderón problem

Calderón problem

- This problem is equivalent to the electrical impedance tomography problem in dimension 3.

Calderón problem

- This problem is equivalent to the electrical impedance tomography problem in dimension 3.
- The problem has been solved in dimension $n = 2$ [Belishev: 2003].

Calderón problem

- This problem is equivalent to the electrical impedance tomography problem in dimension 3.
- The problem has been solved in dimension $n = 2$ [Belishev: 2003].
- Solved in dimensions $n \geq 3$ when M is an analytic manifold [Lassas, Taylor, Uhlmann: 2003].

Calderón problem

- This problem is equivalent to the electrical impedance tomography problem in dimension 3.
- The problem has been solved in dimension $n = 2$ [Belishev: 2003].
- Solved in dimensions $n \geq 3$ when M is an analytic manifold [Lassas, Taylor, Uhlmann: 2003].
- The smooth cases is still unsolved.

Thoughts

Thoughts

- Even monogenic fields have harmonic components. Given a harmonic r -vector A_r , can we reconstruct a monogenic multivector containing A_r ?

Thoughts

- Even monogenic fields have harmonic components. Given a harmonic r -vector A_r , can we reconstruct a monogenic multivector containing A_r ?
- For $n = 3$, the scalar potential u and magnetic bivector field b are two parts of a monogenic field $f = u + b$ due to Ohm's and Ampere's laws

$$-\nabla \wedge u = \mathbf{j} = \nabla \rfloor b.$$

Thoughts

- Even monogenic fields have harmonic components. Given a harmonic r -vector A_r , can we reconstruct a monogenic multivector containing A_r ?
- For $n = 3$, the scalar potential u and magnetic bivector field b are two parts of a monogenic field $f = u + b$ due to Ohm's and Ampere's laws

$$-\nabla \wedge u = \mathbf{j} = \nabla \rfloor b.$$

- If Λ can provide us $b|_{\partial M}$, then we can possibly reconstruct $\mathcal{M}^+(M)$.

Thoughts

- Even monogenic fields have harmonic components. Given a harmonic r -vector A_r , can we reconstruct a monogenic multivector containing A_r ?
- For $n = 3$, the scalar potential u and magnetic bivector field b are two parts of a monogenic field $f = u + b$ due to Ohm's and Ampere's laws

$$-\nabla \wedge u = \mathbf{j} = \nabla \rfloor b.$$

- If Λ can provide us $b|_{\partial M}$, then we can possibly reconstruct $\mathcal{M}^+(M)$.
- Given the algebraic structure of each $\mathcal{A}_B(M) \subset \mathcal{M}^+(M)$, can this be used to determine g ?

Other inverse problems

Other inverse problems

- Can the magnetic impedance tomography problem can provide some extra insight on the EIT problem?

Other inverse problems

- Can the magnetic impedance tomography problem can provide some extra insight on the EIT problem?
- The Hodge-Morrey decomposition is an instrumental tool for boundary value problems that, for example, allows one to show that Λ determines the Betti numbers of M [Belishev, Sharafutdinov: 2008].

Other inverse problems

- Can the magnetic impedance tomography problem can provide some extra insight on the EIT problem?
- The Hodge-Morrey decomposition is an instrumental tool for boundary value problems that, for example, allows one to show that Λ determines the Betti numbers of M [Belishev, Sharafutdinov: 2008].
- Can the Clifford-Hodge-Morrey decomposition can allow us to work on other related boundary inverse problems?

Other inverse problems

- Can the magnetic impedance tomography problem can provide some extra insight on the EIT problem?
- The Hodge-Morrey decomposition is an instrumental tool for boundary value problems that, for example, allows one to show that Λ determines the Betti numbers of M [Belishev, Sharafutdinov: 2008].
- Can the Clifford-Hodge-Morrey decomposition can allow us to work on other related boundary inverse problems?
- These problems could include spacetime problems where the metric g is of mixed signature.

Section 5

Conclusions

Conclusion

Conclusion

- We have utilized multivector fields to serve as a meaningful generalization of both the complex numbers and differential forms.

Conclusion

- We have utilized multivector fields to serve as a meaningful generalization of both the complex numbers and differential forms.
- This provides a new way to decompose fields on domains of \mathbb{R}^n and this can likely be generalized to arbitrary compact orientable pseudo-Riemannian manifolds.

Conclusion

- We have utilized multivector fields to serve as a meaningful generalization of both the complex numbers and differential forms.
- This provides a new way to decompose fields on domains of \mathbb{R}^n and this can likely be generalized to arbitrary compact orientable pseudo-Riemannian manifolds.
- Likewise, we have proven that the monogenic fields contain a wealth of topological information and this information is supported on the boundary by the Cauchy integral formula.

Data Assimilation

Data Assimilation

- Over the past two years I have also worked with a team on developing new techniques for data assimilation.

Data Assimilation

- Over the past two years I have also worked with a team on developing new techniques for data assimilation.
- We have submitted *Model and Data Reduction for Data Assimilation: Particle Filters Employing Projected Forecasts and Data with Application to a Shallow Water Model*.

Data Assimilation

- Over the past two years I have also worked with a team on developing new techniques for data assimilation.
- We have submitted *Model and Data Reduction for Data Assimilation: Particle Filters Employing Projected Forecasts and Data with Application to a Shallow Water Model*.
- We are continuing to work to apply our scheme to new models such as the Modular Arbitrary-Order Ocean-Atmospheric Model (MAOOAM).

Thank you!