

Riemannian Geometry

for Dummies

Colin Roberts

Colorado
State
University

Section 1

Introduction

Riemannian geometry is the study of a *smooth manifold* M along with a *Riemannian metric* g .

The point of Riemannian geometry is to generalize the differentiable and metric structure of \mathbb{R}^n .

We think of living on the manifold. We refer to this as *intrinsic*.

We generalize to spaces that have interesting topology and geometry.

This will require us to rethink some notions we found “easy”
in \mathbb{R}^n .

But we will gain a very general framework for working with differentiable objects.

Section 2

Motivation

Why study this in the first place?

Example: Partial differential equations (PDEs) on spaces that are not flat.

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth
- Electrical Impedence Tomography (EIT)

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth
- Electrical Impedence Tomography (EIT)
- General relativity

Example: Optimization in interesting spaces.

Example: Optimization in interesting spaces.

- Matrix (symmetry) groups

Example: Optimization in interesting spaces.

- Matrix (symmetry) groups
- Grassmannians, Flags

Example: Optimization in interesting spaces.

- Matrix (symmetry) groups
- Grassmannians, Flags
- Curved spacetime

Example: Rephrasing classical problems.

Example: Rephrasing classical problems.

- EIT

Example: Rephrasing classical problems.

- EIT
- Polymer growth

Example: Rephrasing classical problems.

- EIT
- Polymer growth
- Electrodynamics

Section 3

Preliminaries

Subsection 1

Smooth Manifolds

Our To-Do List:

- Start with a topological space M

Our To-Do List:

- Start with a topological space M
- Look at open sets U that cover M

Our To-Do List:


- Start with a topological space M
- Look at open sets U that cover M
- Construct local coordinates φ

Our To-Do List:

- Start with a topological space M
- Look at open sets U that cover M
- Construct local coordinates φ
- Show coordinate transition functions are smooth

Working example: 2-sphere

$$S^2 := \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}.$$



Take open sets in \mathbb{R}^m

$$\mathcal{O}_\alpha \quad \mathcal{O}_\beta$$

Take open sets in \mathbb{R}^m

$$\mathcal{O}_\alpha \qquad \mathcal{O}_\beta$$

and maps

$$\varphi_\alpha\colon \mathcal{O}_\alpha \rightarrow U_\alpha \subset M \qquad \varphi_\beta\colon \mathcal{O}_\beta \rightarrow U_\beta \subset M.$$


Take open sets in \mathbb{R}^m

$$\mathcal{O}_\alpha \quad \mathcal{O}_\beta$$

and maps

$$\varphi_\alpha: \mathcal{O}_\alpha \rightarrow U_\alpha \subset M \quad \varphi_\beta: \mathcal{O}_\beta \rightarrow U_\beta \subset M.$$

These are our *local coordinates*.



Our local coordinates must work together on overlaps

$$U_\alpha \cap U_\beta.$$


Our local coordinates must work together on overlaps

$$U_\alpha \cap U_\beta.$$

We check the *transition function*

$$\phi_{\alpha\beta} = \varphi_\alpha^{-1} \circ \varphi_\beta$$



is smooth and invertible as a function on \mathbb{R}^m .





Subsection 2

Vector Fields

Vector fields on \mathbb{R}^m are functions $\vec{f}:\mathbb{R}^m \rightarrow \mathbb{R}^m$.



Vector fields on \mathbb{R}^m are functions $\vec{f}: \mathbb{R}^m \rightarrow \mathbb{R}^m$.



Intrinsic vector fields on manifolds carry geometric information.

Our To-Do List:

- Construct the *tangent space* $T_p M$

Our To-Do List:

- Construct the *tangent space* T_pM
- Glue together tangent spaces to form the *tangent bundle* TM

Our To-Do List:


- Construct the *tangent space* T_pM
- Glue together tangent spaces to form the *tangent bundle* TM
- Properly define vector fields X as *sections* of the tangent bundle


- Start with a curve $\gamma(-1, 1) \rightarrow M$ with $\gamma(0) = p$

- Start with a curve $\gamma(-1, 1) \rightarrow M$ with $\gamma(0) = p$
- Find the velocity vector $\dot{\gamma} = \left. \frac{d\gamma}{dt} \right|_{t=0}$

- Start with a curve $\gamma(-1, 1) \rightarrow M$ with $\gamma(0) = p$
- Find the velocity vector $\dot{\gamma} = \left. \frac{d\gamma}{dt} \right|_{t=0}$
- This defines a tangent vector at p

- Start with a curve $\gamma(-1, 1) \rightarrow M$ with $\gamma(0) = p$
- Find the velocity vector $\dot{\gamma} = \left. \frac{d\gamma}{dt} \right|_{t=0}$
- This defines a tangent vector at p
- All possible tangent vectors form the tangent space $T_p M$.






We need to understand how tangent vectors on M relate to vectors in our local coordinates φ .

- The differential $d\varphi$ is a map of tangent vectors

We need to understand how tangent vectors on M relate to vectors in our local coordinates φ .

- The differential $d\varphi$ is a map of tangent vectors
- If $\varphi(x) = p$, then $d\varphi(x): T_x\mathbb{R}^m \rightarrow T_pM$




- We need to understand how different tangent spaces relate


- We need to understand how different tangent spaces relate
- Properly gluing tangent spaces T_pM to the manifold M allows us to build a larger manifold TM .

- We need to understand how different tangent spaces relate
- Properly gluing tangent spaces T_pM to the manifold M allows us to build a larger manifold TM .
- This allows us to see how tangent vectors move around the whole manifold.

We briefly drop a dimension to the 1-sphere

$$S^1 := \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}.$$







- Points in TM are (p, v) with $p \in M$ and $v \in T_pM$

- Points in TM are (p, v) with $p \in M$ and $v \in T_pM$
- So, a function $X: M \rightarrow TM$ selects a tangent vector at every point


- Points in TM are (p, v) with $p \in M$ and $v \in T_pM$
- So, a function $X: M \rightarrow TM$ selects a tangent vector at every point
- For example, $X_p = v \in T_pM$

- Points in TM are (p, v) with $p \in M$ and $v \in T_pM$
- So, a function $X: M \rightarrow TM$ selects a tangent vector at every point
- For example, $X_p = v \in T_pM$
- We have the projection $\pi: TM \rightarrow M$ by $\pi(p, v) = p$

- Points in TM are (p, v) with $p \in M$ and $v \in T_p M$
- So, a function $X: M \rightarrow TM$ selects a tangent vector at every point
- For example, $X_p = v \in T_p M$
- We have the projection $\pi: TM \rightarrow M$ by $\pi(p, v) = p$
- X is a *section* if $\pi \circ X = \text{Id}_M$ (vertical line test)




Back to the 2-sphere.




Subsection 3

Specific Coordinates

We should work with specific coordinates on S^2 .






- We can take a vector field in \mathbb{R}^m and push it forward onto M

- We can take a vector field in \mathbb{R}^m and push it forward onto M
- We extend the differential $d\varphi(x): T_x M \rightarrow T_p M$ to a map on bundles

- We can take a vector field in \mathbb{R}^m and push it forward onto M
- We extend the differential $d\varphi(x): T_x M \rightarrow T_p M$ to a map on bundles
- This bundle map $\varphi_*: T\mathbb{R}^m \rightarrow TM$ is the *pushforward*



Section 4

Riemannian Geometry

Our To-Do List:

- Build an inner product on the tangent space $T_p M$;

Our To-Do List:

- Build an inner product on the tangent space T_pM ;
- Have the inner product vary smoothly as we vary the point p ;

Our To-Do List:

- Build an inner product on the tangent space T_pM ;
- Have the inner product vary smoothly as we vary the point p ;
- Define this as our Riemannian metric tensor field g ;

Our To-Do List:

- Build an inner product on the tangent space T_pM ;
- Have the inner product vary smoothly as we vary the point p ;
- Define this as our Riemannian metric tensor field g ;
- Extract geometrical and analytical qualities of the underlying manifold M .

Subsection 1

Riemannian Metric

We use the differential and dot product to form a matrix at each point

$$g_{ij}(x) = d\varphi(x) e_i \cdot d\varphi(x) e_j.$$

We use the differential and dot product to form a matrix at each point

$$g_{ij}(x) = d\varphi(x)e_i \cdot d\varphi(x)e_j.$$

This matrix is the *Riemannian metric*.

Riemannian metric provides an inner product for tangent vectors on M . Thus, we know

- how lengths are distorted;
- how volume is distorted.

This allows us to integrate or differentiate in our coordinates but think of it as intrinsic to the manifold.

- The Riemannian metric gives us a distance function $d(p, q)$ on M

- The Riemannian metric gives us a distance function $d(p, q)$ on M
- Compute this by finding the length of the shortest curve

$$\gamma: [0, 1] \rightarrow M \quad \gamma(0) = p, \quad \gamma(1) = q$$

- The Riemannian metric gives us a distance function $d(p, q)$ on M
- Compute this by finding the length of the shortest curve

$$\gamma: [0, 1] \rightarrow M \quad \gamma(0) = p, \quad \gamma(1) = q$$


We need to solve

$$\inf_{\gamma} \ell(\gamma) := \int_0^1 \sqrt{g(\dot{\gamma}, \dot{\gamma})} dt$$

- Reminder: in \mathbb{R}^m , the speed of a curve is $\sqrt{\dot{\gamma}, \dot{\gamma}}$

- Reminder: in \mathbb{R}^m , the speed of a curve is $\sqrt{\dot{\gamma}, \dot{\gamma}}$
- $g(\dot{\gamma}, \dot{\gamma})$ is the speed on M

- Reminder: in \mathbb{R}^m , the speed of a curve is $\sqrt{\dot{\gamma}, \dot{\gamma}}$
- $g(\dot{\gamma}, \dot{\gamma})$ is the speed on M
- We put $g(\dot{\gamma}, \dot{\gamma})$ to mean $\sum_{i,j=1}^m g_{ij} \dot{\gamma}_i \dot{\gamma}_j$.



Solving this optimization problem yields the *geodesic equation*

$$\ddot{x}^l + \dot{x}^j \dot{x}^k \Gamma_{jk}^l = 0$$

where Γ_{jk}^l are the *Christoffel symbols* which are formed by derivatives of the metric.

This defines an intrinsic derivative ∇ called the *Levi-Civita connection*


- Since we know how vectors are transformed, combining those describes transformed volumes.

- Since we know how vectors are transformed, combining those describes transformed volumes.
- The determinant gives us area information.

- Since we know how vectors are transformed, combining those describes transformed volumes.
- The determinant gives us area information.
- Then $\sqrt{|\det(g(x))|}$ gives us the volume on M

In spherical coordinates, $\sqrt{|\det(g)|} = \sin \varphi$ which gives us the integrand

$$\sin \varphi d\varphi d\theta.$$



Section 5

Conclusions

- We constructed a smooth manifold M

- We constructed a smooth manifold M
- We generalized vector fields X to M

- We constructed a smooth manifold M
- We generalized vector fields X to M
- We created an inner product g on M to measure these fields

- We constructed a smooth manifold M
- We generalized vector fields X to M
- We created an inner product g on M to measure these fields
- No measurement depends on the choice of coordinates

- g allows us to measure vectors and understand the geometry of M in coordinates φ

- g allows us to measure vectors and understand the geometry of M in coordinates φ
- Hence, we can define lengths and volumes

- g allows us to measure vectors and understand the geometry of M in coordinates φ
- Hence, we can define lengths and volumes
- Thus, we can integrate

- g induces a derivative ∇

- g induces a derivative ∇
- g induces a Laplacian Δ

- g induces a derivative ∇
- g induces a Laplacian Δ
- g provides an intrinsic length function on M

This is just the beginning!