

The Calderón Problem

on Riemannian Manifolds

Colin Roberts

Colorado
State
University

Outline

Outline

- Introduce the Calderón problem.

Outline

- Introduce the Calderón problem.
- Discuss some of the current results and issues.

Outline

- Introduce the Calderón problem.
- Discuss some of the current results and issues.
- Rephrase the problem in a geometrical way.

Outline

- Introduce the Calderón problem.
- Discuss some of the current results and issues.
- Rephrase the problem in a geometrical way.
- Prove the problem in 2 dimensions using the boundary control method.

Outline

- Introduce the Calderón problem.
- Discuss some of the current results and issues.
- Rephrase the problem in a geometrical way.
- Prove the problem in 2 dimensions using the boundary control method.
- What can and can't we do to generalize this method?

Section 1

Introduction

Subsection 1

Calderón Problem

In 1980, Alberto Calderón proposed a problem in his paper *On an inverse boundary value problem*.

In 1980, Alberto Calderón proposed a problem in his paper *On an inverse boundary value problem*.

- He wanted to know if one can determine the conductivity of a domain by making voltage and current measurements along the boundary.

In 1980, Alberto Calderón proposed a problem in his paper *On an inverse boundary value problem*.


- He wanted to know if one can determine the conductivity of a domain by making voltage and current measurements along the boundary.
- This is the Electrical Impedance Tomography problem.

In 1980, Alberto Calderón proposed a problem in his paper *On an inverse boundary value problem*.

- He wanted to know if one can determine the conductivity of a domain by making voltage and current measurements along the boundary.
- This is the Electrical Impedance Tomography problem.
- Originally his motivation was for oil prospecting.

In 1980, Alberto Calderón proposed a problem in his paper *On an inverse boundary value problem*.

- He wanted to know if one can determine the conductivity of a domain by making voltage and current measurements along the boundary.
- This is the Electrical Impedance Tomography problem.
- Originally his motivation was for oil prospecting.
- This problem sparked interest due to its usefulness in geophysical and medical imaging.



The two main groups working on this problem now are...

The two main groups working on this problem now are...

- Practitioners: Work with incomplete and noisy data to recover information in the real world.

The two main groups working on this problem now are...

- Practitioners: Work with incomplete and noisy data to recover information in the real world.
- Theorists: Work in ideal scenarios with a chosen amount of information to see the scope of possibilities.

The two main groups working on this problem now are...

- Practitioners: Work with incomplete and noisy data to recover information in the real world.
- Theorists: Work in ideal scenarios with a chosen amount of information to see the scope of possibilities.

Problem Statement

Idea: Given a domain Ω with interior Ω^+ that we cannot probe, can we determine the conductivity γ matrix by studying the boundary $\partial\Omega$?

Problem Statement

Idea: Given a domain Ω with interior Ω^+ that we cannot probe, can we determine the conductivity γ matrix by studying the boundary $\partial\Omega$?

- Ω^+ is free of charges, hence $\nabla \cdot (\gamma \nabla u) = 0$ in Ω^+ where u is the electrostatic potential.

Problem Statement

Idea: Given a domain Ω with interior Ω^+ that we cannot probe, can we determine the conductivity γ matrix by studying the boundary $\partial\Omega$?

- Ω^+ is free of charges, hence $\nabla \cdot (\gamma \nabla u) = 0$ in Ω^+ where u is the electrostatic potential.
- Apply a known voltage f at the boundary $\partial\Omega$. Hence $f = u|_{\partial\Omega}$.

Problem Statement

Idea: Given a domain Ω with interior Ω^+ that we cannot probe, can we determine the conductivity γ matrix by studying the boundary $\partial\Omega$?

- Ω^+ is free of charges, hence $\nabla \cdot (\gamma \nabla u) = 0$ in Ω^+ where u is the electrostatic potential.
- Apply a known voltage f at the boundary $\partial\Omega$. Hence $f = u|_{\partial\Omega}$.
- Measure the current flux h through the boundary $\partial\Omega$. Hence, $h = \frac{\partial u}{\partial \nu}$.

Problem Statement

Idea: Given a domain Ω with interior Ω^+ that we cannot probe, can we determine the conductivity γ matrix by studying the boundary $\partial\Omega$?

- Ω^+ is free of charges, hence $\nabla \cdot (\gamma \nabla u) = 0$ in Ω^+ where u is the electrostatic potential.
- Apply a known voltage f at the boundary $\partial\Omega$. Hence $f = u|_{\partial\Omega}$.
- Measure the current flux h through the boundary $\partial\Omega$. Hence, $h = \frac{\partial u}{\partial \nu}$.
- This defines the voltage-to-current map Λ so that $\Lambda(f) = h$.

Problem Statement

Idea: Given a domain Ω with interior Ω^+ that we cannot probe, can we determine the conductivity γ matrix by studying the boundary $\partial\Omega$?

- Ω^+ is free of charges, hence $\nabla \cdot (\gamma \nabla u) = 0$ in Ω^+ where u is the electrostatic potential.
- Apply a known voltage f at the boundary $\partial\Omega$. Hence $f = u|_{\partial\Omega}$.
- Measure the current flux h through the boundary $\partial\Omega$. Hence, $h = \frac{\partial u}{\partial \nu}$.
- This defines the voltage-to-current map Λ so that $\Lambda(f) = h$.
- Can we determine the conductivity matrix γ from Λ ?

Section 2

The Calderón Problem on Riemannian Manifolds

Subsection 1

Preliminaries

Geometry

Geometry

- *Smooth n -dimensional manifold*: A space that locally looks like (is C^∞ diffeomorphic to) an open subset of \mathbb{R}^n .

Geometry


- *Smooth n -dimensional manifold*: A space that locally looks like (is C^∞ diffeomorphic to) an open subset of \mathbb{R}^n .
- *Riemannian metric*: A smoothly varying inner product defined on Ω . In coordinates, g takes the form of a symmetric and positive definite matrix with entries g_{jk} with inverse g^{jk} .


Geometry


- *Smooth n -dimensional manifold*: A space that locally looks like (is C^∞ diffeomorphic to) an open subset of \mathbb{R}^n .
- *Riemannian metric*: A smoothly varying inner product defined on Ω . In coordinates, g takes the form of a symmetric and positive definite matrix with entries g_{jk} with inverse g^{jk} .
- *Exterior algebra*: Differential forms with the wedge product \wedge .

Geometry


- *Smooth n -dimensional manifold*: A space that locally looks like (is C^∞ diffeomorphic to) an open subset of \mathbb{R}^n .
- *Riemannian metric*: A smoothly varying inner product defined on Ω . In coordinates, g takes the form of a symmetric and positive definite matrix with entries g_{jk} with inverse g^{jk} .
- *Exterior algebra*: Differential forms with the wedge product \wedge .
- *Hodge Star*: Attached to the exterior algebra when we also have a Riemannian metric. Gives an isomorphism between k and $n - k$ -forms.









1-Forms



2-Forms



3-Forms



Partial Differential Equations

Partial Differential Equations

- *k-Form Inner Product:* $\langle \alpha, \beta \rangle = \int_{\Omega} \alpha \wedge \star \beta.$

Partial Differential Equations

- *k-Form Inner Product*: $\langle \alpha, \beta \rangle = \int_{\Omega} \alpha \wedge \star \beta$.
- *Exterior Derivative*: Derivative operator d defined on k -forms.

Partial Differential Equations

- *k-Form Inner Product*: $\langle \alpha, \beta \rangle = \int_{\Omega} \alpha \wedge \star \beta$.
- *Exterior Derivative*: Derivative operator d defined on k -forms.
- *Codifferential*: Formal adjoint to d written as δ .

Partial Differential Equations

- *k-Form Inner Product*: $\langle \alpha, \beta \rangle = \int_{\Omega} \alpha \wedge \star \beta$.
- *Exterior Derivative*: Derivative operator d defined on k -forms.
- *Codifferential*: Formal adjoint to d written as δ .
- *Dirac Operator*: $D = d + \delta$.

Partial Differential Equations

- *k-Form Inner Product*: $\langle \alpha, \beta \rangle = \int_{\Omega} \alpha \wedge \star \beta$.
- *Exterior Derivative*: Derivative operator d defined on k -forms.
- *Codifferential*: Formal adjoint to d written as δ .
- *Dirac Operator*: $D = d + \delta$.
- *Laplace-Beltrami Operator*: $\Delta = d\delta + \delta d = D^2$ and in coordinates

$$\Delta f = \frac{1}{\sqrt{|g|}} \sum_{j=1}^n \sum_{k=1}^n \frac{\partial}{\partial x^i} \sqrt{|g|} g^{ij} \frac{\partial}{\partial x^j} f$$

Subsection 2

Raphrasing EIT Problem in a Geometrical Language

- Let (unknown) connected Ω be a smooth Riemannian manifold with boundary $\partial\Omega$.

- Let (unknown) connected Ω be a smooth Riemannian manifold with boundary $\partial\Omega$.
- Replace conductivity be represented by (unknown) g since $\gamma^{jk} = \sqrt{|g|}g^{jk}$.

- Let (unknown) connected Ω be a smooth Riemannian manifold with boundary $\partial\Omega$.
- Replace conductivity be represented by (unknown) g since $\gamma^{jk} = \sqrt{|g|}g^{jk}$.
- Let $\Delta u = 0$ in Ω^+ and $u = f$ on $\partial\Omega$.

- Let (unknown) connected Ω be a smooth Riemannian manifold with boundary $\partial\Omega$.
- Replace conductivity be represented by (unknown) g since $\gamma^{jk} = \sqrt{|g|}g^{jk}$.
- Let $\Delta u = 0$ in Ω^+ and $u = f$ on $\partial\Omega$.
- *Dirichlet-to-Neumann operator* Λ maps Dirichlet data $f = u|_{\partial\Omega}$ to $g = \frac{\partial u}{\partial \nu} = \iota^*(\star du)$.

- Let (unknown) connected Ω be a smooth Riemannian manifold with boundary $\partial\Omega$.
- Replace conductivity be represented by (unknown) g since $\gamma^{jk} = \sqrt{|g|}g^{jk}$.
- Let $\Delta u = 0$ in Ω^+ and $u = f$ on $\partial\Omega$.
- *Dirichlet-to-Neumann operator* Λ maps Dirichlet data $f = u|_{\partial\Omega}$ to $g = \frac{\partial u}{\partial \nu} = \iota^*(\star du)$.
- Recover g from knowing Λ .

Subsection 3

Expected and Current Results

Formal Variable Count

Formal Variable Count

$f \mapsto \Lambda(f)$ approximated by

$$\Lambda(f)_j = \sum_{k=1}^m \lambda_{jk} f_k.$$

Formal Variable Count

$f \mapsto \Lambda(f)$ approximated by

$$\Lambda(f)_j = \sum_{k=1}^m \lambda_{jk} f_k.$$

In the smooth setting,

$$\Lambda(f) = \int_{\partial\Omega} \lambda(x, y) f(y) dS(y).$$

Formal Variable Count

$f \mapsto \Lambda(f)$ approximated by

$$\Lambda(f)_j = \sum_{k=1}^m \lambda_{jk} f_k.$$

In the smooth setting,

$$\Lambda(f) = \int_{\partial\Omega} \lambda(x, y) f(y) dS(y).$$

So, g is a function of n variables that needs to be determined by the kernel λ which is $2n - 2$ variables.

Formal Variable Count

Formal Variable Count

- $n = 1$ gives us an undetermined system.

Formal Variable Count

- $n = 1$ gives us an undetermined system.
- $n = 2$ is well determined.

Formal Variable Count

- $n = 1$ gives us an undetermined system.
- $n = 2$ is well determined.
- $n \geq 3$ is overdetermined.

Issue In 2-Dimensions

Issue In 2-Dimensions

In 2D, Δ is conformally invariant.

Issue In 2-Dimensions

In 2D, Δ is conformally invariant.

Indeed, let $\tilde{g} = e^{2\phi}g$, then

$$\Delta_{\tilde{g}} = e^{-2\phi} \Delta_g + (n-2) e^{-2\phi} g^{jk} \frac{\partial \phi}{\partial x^k} \frac{\partial}{\partial x^j}$$

Issue In 2-Dimensions

In 2D, Δ is conformally invariant.

Indeed, let $\tilde{g} = e^{2\phi}g$, then

$$\Delta_{\tilde{g}} = e^{-2\phi} \Delta_g + (n-2) e^{-2\phi} g^{jk} \frac{\partial \phi}{\partial x^k} \frac{\partial}{\partial x^j}$$

When $n = 2$, the extra term cancels.

1 Dimension

1 Dimension

- The problem in 1 dimension is trivial.

1 Dimension

- The problem in 1 dimension is trivial.
- Can only know the total impedance between the two electrodes.

Isotropic Case

For g isotropic and $n \geq 3$ one can determine g from Λ .

(*Sylvester-Uhlmann 1987*)

2 Dimensional Anisotropic

- Can recover g up to conformal class and can't do better.
- Proven by Lassas and Uhlmann in *On Determining the Riemannian manifold from the Dirichlet to Neumann map*.

3+ Dimensional Anisotropic

3+ Dimensional Anisotropic

- For real analytic manifolds, the (scalar/classical) DN map determines the manifold up to isometry. This gives the topological information as well.
(Lassas, Taylor, Uhlmann *The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary*.)

3+ Dimensional Anisotropic

- For real analytic manifolds, the (scalar/classical) DN map determines the manifold up to isometry. This gives the topological information as well.
(Lassas, Taylor, Uhlmann *The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary*.)
- Can determine the boundary C^∞ -jet of g in Lee and Uhlmann's *Determining anisotropic real-analytic conductivities by boundary measurements*.

3+ Dimensional Anisotropic

- For real analytic manifolds, the (scalar/classical) DN map determines the manifold up to isometry. This gives the topological information as well. (Lassas, Taylor, Uhlmann *The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary*).
- Can determine the boundary C^∞ -jet of g in Lee and Uhlmann's *Determining anisotropic real-analytic conductivities by boundary measurements*.
- For smooth manifolds, the anisotropic problem is open. The goal is to recover the metric up to isometry.

Section 3

Boundary Control Method in 2 Dimensions

Theorem

Theorem

Two 2-dimensional compact orientable manifolds with single common boundary are conformally equivalent iff their DN-maps coincide.

Belishev's *The Calderón Problem for Two-Dimensional Manifolds by the BC-Method.*

Key Ideas

Key Ideas

- Surfaces are two dimensional and can be related to \mathbb{C} .

Key Ideas

- Surfaces are two dimensional and can be related to \mathbb{C} .
- Holomorphic functions have components that are harmonic.

Key Ideas

- Surfaces are two dimensional and can be related to \mathbb{C} .
- Holomorphic functions have components that are harmonic.
- Hilbert transform converts one harmonic function to another by connecting them via a single holomorphic function.

Key Ideas

- Surfaces are two dimensional and can be related to \mathbb{C} .
- Holomorphic functions have components that are harmonic.
- Hilbert transform converts one harmonic function to another by connecting them via a single holomorphic function.
- Gelfand transform relates an algebra \mathcal{A} to the algebra of continuous functions on the spectrum of that algebra, $C(\text{spec}\mathcal{A})$.

Key Ideas

- Surfaces are two dimensional and can be related to \mathbb{C} .
- Holomorphic functions have components that are harmonic.
- Hilbert transform converts one harmonic function to another by connecting them via a single holomorphic function.
- Gelfand transform relates an algebra \mathcal{A} to the algebra of continuous functions on the spectrum of that algebra, $C(\text{spec}\mathcal{A})$.
- This gives us a way to realize Ω from functions defined on Ω that we have access to.

Outline of the Proof

Outline of the Proof

- From DN map, recover the algebra of holomorphic functions. (Lemma 1)

Outline of the Proof

- From DN map, recover the algebra of holomorphic functions. (Lemma 1)
- Show that this algebra is generic. (Lemma 2)

Outline of the Proof

- From DN map, recover the algebra of holomorphic functions. (Lemma 1)
- Show that this algebra is generic. (Lemma 2)
- Represent the trace algebra with the DN map. (Lemma 3)

Outline of the Proof

- From DN map, recover the algebra of holomorphic functions. (Lemma 1)
- Show that this algebra is generic. (Lemma 2)
- Represent the trace algebra with the DN map. (Lemma 3)
- Construct the manifold. (Theorem)

Some Notes

Some Notes

- We have the inclusion of the boundary $\iota: \partial\Omega \rightarrow \Omega$.

Some Notes


- We have the inclusion of the boundary $\iota: \partial\Omega \rightarrow \Omega$.
- The pullback $\iota^*: T^*\Omega \rightarrow T^*\partial\Omega$.

Some Notes

- We have the inclusion of the boundary $\iota: \partial\Omega \rightarrow \Omega$.
- The pullback $\iota^*: T^*\Omega \rightarrow T^*\partial\Omega$.
- Define Λ by $\iota^*(\star du)$ for a harmonic u .

Some Notes

- We have the inclusion of the boundary $\iota: \partial\Omega \rightarrow \Omega$.
- The pullback $\iota^*: T^*\Omega \rightarrow T^*\partial\Omega$.
- Define Λ by $\iota^*(\star du)$ for a harmonic u .
- Λ maps boundary k -forms to boundary $n - k - 1$ forms.



Subsection 1

Lemma 1

Lemma 1

Lemma 1

- A function u satisfying $\Delta u = 0$ has a conjugate function v if and only if the trace $\iota^* u$ satisfies

$$[\Lambda + d\Lambda^{-1}d]\iota^* u = 0.$$

Lemma 1

- A function u satisfying $\Delta u = 0$ has a conjugate function v if and only if the trace $\iota^* u$ satisfies

$$\left[\Lambda + d\Lambda^{-1}d \right] \iota^* u = 0.$$

- $\dim \text{Ran} \left[\Lambda + d\Lambda^{-1}d \right] = \beta_1(\Omega).$

Corollary

Λ completely determines the topology of Ω .

Proof

Proof

- Since Ω is a single connected component, $\beta_0(\Omega) = 1$.
- We have $\beta_1(\Omega)$ from before.
- Since Ω is a surface with boundary, $\beta_2(\Omega) = 0$.
- Since Ω is two dimensional, $\beta_n(\Omega) = 0$ for $n \geq 3$.

Conjugate Function Intuition

Conjugate Function Intuition

- Suppose we have holomorphic complex function $w = u + iv$.

Conjugate Function Intuition

- Suppose we have holomorphic complex function $w = u + iv$.
- Let u be a 0-form and v as a 1-form.

Conjugate Function Intuition

- Suppose we have holomorphic complex function $w = u + iv$.
- Let u be a 0-form and v as a 1-form.
- Then $\frac{\partial}{\partial \bar{z}}$ is given by $D = d + \delta$.

Conjugate Function Intuition

- Suppose we have holomorphic complex function $w = u + iv$.
- Let u be a 0-form and v as a 1-form.
- Then $\frac{\partial}{\partial \bar{z}}$ is given by $D = d + \delta$.
- $Dw = 0$ gives us the Cauchy-Riemann equations
- We call u and v conjugate by CREs.

Hilbert Transform

Hilbert Transform

- We can get v from u via the Hilbert transform.

Hilbert Transform

- We can get v from u via the Hilbert transform.
- Define $\mathcal{H} = d\Lambda^{-1}$.

Construct the Algebra $\mathcal{A}(\Omega)$

Construct the Algebra $\mathcal{A}(\Omega)$

- By Lemma 1 we can now create the algebra $\mathcal{A}(\Omega) \subset C(\Omega)$ from harmonic functions u with conjugates v by

$$\mathcal{A}(\Omega) := \{w = u + iv\}.$$

Algebra since product of two holomorphic functions is holomorphic.

Construct the Algebra $\mathcal{A}(\Omega)$

- By Lemma 1 we can now create the algebra $\mathcal{A}(\Omega) \subset C(\Omega)$ from harmonic functions u with conjugates v by

$$\mathcal{A}(\Omega) := \{w = u + iv\}.$$

Algebra since product of two holomorphic functions is holomorphic.

- In isothermal coordinates, each $w \in \mathcal{A}(\Omega)$ is holomorphic.

Construct the Algebra $\mathcal{A}(\Omega)$

- By Lemma 1 we can now create the algebra $\mathcal{A}(\Omega) \subset C(\Omega)$ from harmonic functions u with conjugates v by

$$\mathcal{A}(\Omega) := \{w = u + iv\}.$$

Algebra since product of two holomorphic functions is holomorphic.

- In isothermal coordinates, each $w \in \mathcal{A}(\Omega)$ is holomorphic.
- This gives Ω a complex structure.

Construct the Algebra $\mathcal{A}(\Omega)$

- By Lemma 1 we can now create the algebra $\mathcal{A}(\Omega) \subset C(\Omega)$ from harmonic functions u with conjugates v by

$$\mathcal{A}(\Omega) := \{w = u + iv\}.$$

Algebra since product of two holomorphic functions is holomorphic.

- In isothermal coordinates, each $w \in \mathcal{A}(\Omega)$ is holomorphic.
- This gives Ω a complex structure.
- This is analogous to having the Hodge star on a surface.

Subsection 2

Lemma 2

Gelfand Representation

Gelfand Representation

- Let \mathcal{M} be the set of multiplicative linear functionals on a commutative Banach algebra \mathcal{A} .

Gelfand Representation

- Let \mathcal{M} be the set of multiplicative linear functionals on a commutative Banach algebra \mathcal{A} .
- The Gelfand transform gives a way of representing an algebra \mathcal{A} as a function algebra $\hat{\mathcal{A}}$.

Gelfand Representation

- Let \mathcal{M} be the set of multiplicative linear functionals on a commutative Banach algebra \mathcal{A} .
- The Gelfand transform gives a way of representing an algebra \mathcal{A} as a function algebra $\hat{\mathcal{A}}$.
- The *Gelfand transform* maps $a \in \mathcal{A}$ to a function \hat{a} on \mathcal{M} by

$$\hat{a}(\delta) := \delta(a), \quad \delta \in \mathcal{M}.$$

Gelfand Representation

- Let \mathcal{M} be the set of multiplicative linear functionals on a commutative Banach algebra \mathcal{A} .
- The Gelfand transform gives a way of representing an algebra \mathcal{A} as a function algebra $\hat{\mathcal{A}}$.
- The *Gelfand transform* maps $a \in \mathcal{A}$ to a function \hat{a} on \mathcal{M} by

$$\hat{a}(\delta) := \delta(a), \quad \delta \in \mathcal{M}.$$

- The *Gelfand topology* is the weakest topology on \mathcal{M} in which all \hat{a} are continuous. This makes \mathcal{M} compact.

Gelfand Representation

- Let \mathcal{M} be the set of multiplicative linear functionals on a commutative Banach algebra \mathcal{A} .
- The Gelfand transform gives a way of representing an algebra \mathcal{A} as a function algebra $\hat{\mathcal{A}}$.
- The *Gelfand transform* maps $a \in \mathcal{A}$ to a function \hat{a} on \mathcal{M} by

$$\hat{a}(\delta) := \delta(a), \quad \delta \in \mathcal{M}.$$

- The *Gelfand topology* is the weakest topology on \mathcal{M} in which all \hat{a} are continuous. This makes \mathcal{M} compact.
- \mathcal{M} with this topology is called the *spectrum* $\text{spec}\mathcal{A}$.

Generic Algebras

Generic Algebras

- The Gelfand transform $\hat{\mathcal{A}}$ is a subalgebra of $C(\operatorname{spec}\mathcal{A})$ and $a \mapsto \hat{a}$ is an isometric isomorphism.

Generic Algebras

- The Gelfand transform $\hat{\mathcal{A}}$ is a subalgebra of $C(\operatorname{spec}\mathcal{A})$ and $a \mapsto \hat{a}$ is an isometric isomorphism.
- For a function algebra $\mathcal{A} \subset C(X)$, take $\epsilon: X \rightarrow \operatorname{spec}\mathcal{A}$ with $\epsilon(x) = \delta_x$.

Generic Algebras

- The Gelfand transform $\hat{\mathcal{A}}$ is a subalgebra of $C(\operatorname{spec}\mathcal{A})$ and $a \mapsto \hat{a}$ is an isometric isomorphism.
- For a function algebra $\mathcal{A} \subset C(X)$, take $\epsilon: X \rightarrow \operatorname{spec}\mathcal{A}$ with $\epsilon(x) = \delta_x$.
- A function algebra $\mathcal{A} \subset C(X)$ is *generic* if ϵ is a homeomorphism.

Generic Algebras

- The Gelfand transform $\hat{\mathcal{A}}$ is a subalgebra of $C(\operatorname{spec}\mathcal{A})$ and $a \mapsto \hat{a}$ is an isometric isomorphism.
- For a function algebra $\mathcal{A} \subset C(X)$, take $\epsilon: X \rightarrow \operatorname{spec}\mathcal{A}$ with $\epsilon(x) = \delta_x$.
- A function algebra $\mathcal{A} \subset C(X)$ is *generic* if ϵ is a homeomorphism.
- A generic algebra is (spatially) isomorphic to its Gelfand transform.

Lemma 2

Lemma 2

The algebra of holomorphic functions $\mathcal{A}(\Omega)$ is generic.

Importance

Importance

- $\hat{\mathcal{A}}(\partial\Omega)$ is (spatially) isomorphic to $\mathcal{A}(\Omega)$ by taking the Gelfand transform of the trace.

Importance

- $\hat{\mathcal{A}}(\partial\Omega)$ is (spatially) isomorphic to $\mathcal{A}(\Omega)$ by taking the Gelfand transform of the trace.
- The lemma shows that $\epsilon: \Omega \rightarrow \text{spec}\mathcal{A}(\Omega)$ is a homeomorphism, so we have determined Ω up to homeomorphism.

What's Left?

What's Left?

We can only have hope access to the trace algebra $\mathcal{A}(\partial\Omega)$. So we need to determine this to reach our goal.

Subsection 3

Lemma 3

Trace Algebra

Trace Algebra

- The trace algebra $\mathcal{A}(\partial\Omega) := \iota^* \mathcal{A}(\Omega)$ is isometrically isomorphic to $\mathcal{A}(\Omega)$ since

$$\|w\|_{\mathcal{A}(\Omega)} = \|\iota^* w\|_{\mathcal{A}(\partial\Omega)}$$

and since a holomorphic function is uniquely determined by its boundary values.

Lemma 3

We have the representation

$$\mathcal{A}(\partial\Omega) = \text{clos}_{C(\partial\Omega)} \{f + ih\},$$

where h is conjugate to f by \mathcal{H} .

Subsection 4

Proof of the Main Theorem

Construction of the Manifold

Construction of the Manifold

Following these steps yields a manifold (Ω, g) with the DN map Λ .

Construction of the Manifold

Following these steps yields a manifold (Ω, g) with the DN map Λ .

- Step 1: We know $g|_{\partial\Omega}$ by Lee and Uhlmann, and thus we know \mathcal{H} and $C^\infty(\partial\Omega)$. This allows us to recover the trace algebra $\mathcal{A}(\partial\Omega)$ using the representation in Lemma 3.

Construction of the Manifold

Following these steps yields a manifold (Ω, g) with the DN map Λ .

- Step 1: We know $g|_{\partial\Omega}$ by Lee and Uhlmann, and thus we know \mathcal{H} and $C^\infty(\partial\Omega)$. This allows us to recover the trace algebra $\mathcal{A}(\partial\Omega)$ using the representation in Lemma 3.
- Step 2: Then find $\text{spec}\mathcal{A}(\partial\Omega) = \Omega$ by Lemma 2.

Construction of the Manifold

Following these steps yields a manifold (Ω, g) with the DN map Λ .

- Step 1: We know $g|_{\partial\Omega}$ by Lee and Uhlmann, and thus we know \mathcal{H} and $C^\infty(\partial\Omega)$. This allows us to recover the trace algebra $\mathcal{A}(\partial\Omega)$ using the representation in Lemma 3.
- Step 2: Then find $\text{spec}\mathcal{A}(\partial\Omega) = \Omega$ by Lemma 2.
- Step 3: Next, the Gelfand transform $\hat{\mathcal{A}}(\partial\Omega) = \mathcal{A}(\Omega)$ by Lemma 2.

Construction of the Manifold

Following these steps yields a manifold (Ω, g) with the DN map Λ .

- Step 1: We know $g|_{\partial\Omega}$ by Lee and Uhlmann, and thus we know \mathcal{H} and $C^\infty(\partial\Omega)$. This allows us to recover the trace algebra $\mathcal{A}(\partial\Omega)$ using the representation in Lemma 3.
- Step 2: Then find $\text{spec}\mathcal{A}(\partial\Omega) = \Omega$ by Lemma 2.
- Step 3: Next, the Gelfand transform $\hat{\mathcal{A}}(\partial\Omega) = \mathcal{A}(\Omega)$ by Lemma 2.
- Step 4: $\mathcal{A}(\Omega)$ gives us the complex structure on Ω by Lemma 1.

Construction of the Manifold

Following these steps yields a manifold (Ω, g) with the DN map Λ .

- Step 1: We know $g|_{\partial\Omega}$ by Lee and Uhlmann, and thus we know \mathcal{H} and $C^\infty(\partial\Omega)$. This allows us to recover the trace algebra $\mathcal{A}(\partial\Omega)$ using the representation in Lemma 3.
- Step 2: Then find $\text{spec}\mathcal{A}(\partial\Omega) = \Omega$ by Lemma 2.
- Step 3: Next, the Gelfand transform $\hat{\mathcal{A}}(\partial\Omega) = \mathcal{A}(\Omega)$ by Lemma 2.
- Step 4: $\mathcal{A}(\Omega)$ gives us the complex structure on Ω by Lemma 1.
- Step 5: Equip Ω with a metric g conforming to this complex structure.

Section 4

Generalizing This Method

First Issue

First Issue

- No complex structure in higher dimensions.

Resolution

Resolution

- Use a Clifford algebra/calculus structure to replace \mathbb{C} .

Resolution

- Use a Clifford algebra/calculus structure to replace \mathbb{C} .
- The tools of Clifford analysis allow us to recover a notion of holomorphicity known as *monogenicity*.

Resolution

- Use a Clifford algebra/calculus structure to replace \mathbb{C} .
- The tools of Clifford analysis allow us to recover a notion of holomorphicity known as *monogenicity*.
- We can recover a similar algebra (Hardy space) of monogenic functions.

Clifford Structure

A Clifford algebra builds upon the exterior algebra of forms. Given a quadratic space (V, Q) , we have

Clifford Structure

A Clifford algebra builds upon the exterior algebra of forms. Given a quadratic space (V, Q) , we have

- The quotient of the tensor algebra

$$Cl(V, Q) = \bigoplus_{j=0}^{\infty} V^{\otimes j} / \langle v \otimes v - Q(v) \rangle.$$

Clifford Structure

A Clifford algebra builds upon the exterior algebra of forms. Given a quadratic space (V, Q) , we have

- The quotient of the tensor algebra

$$\text{Cl}(V, Q) = \bigoplus_{j=0}^{\infty} V^{\otimes j} / \langle v \otimes v - Q(v) \rangle.$$

- If $Q = g$ is an inner product, this yields a geometric product on vectors $u, v \in \text{Cl}(V, g)$

$$uv = g(u, v) + u \wedge v.$$

Clifford Structure

A Clifford algebra builds upon the exterior algebra of forms. Given a quadratic space (V, Q) , we have

- The quotient of the tensor algebra

$$\text{Cl}(V, Q) = \bigoplus_{j=0}^{\infty} V^{\otimes j} / \langle v \otimes v - Q(v) \rangle.$$

- If $Q = g$ is an inner product, this yields a geometric product on vectors $u, v \in \text{Cl}(V, g)$

$$uv = g(u, v) + u \wedge v.$$

- Geometric product can be extended to multivectors.

Familiar Examples

Familiar Examples

We have all seen Clifford algebras before. Indeed,

Familiar Examples

We have all seen Clifford algebras before. Indeed,

- $\mathbb{C} \cong \text{Cl}(\mathbb{R}, -\cdot).$

Familiar Examples

We have all seen Clifford algebras before. Indeed,

- $\mathbb{C} \cong \text{Cl}(\mathbb{R}, -\cdot)$.
- \mathbb{C} also lives inside of $\text{Cl}(\mathbb{R}^2, \cdot)$ as the even subalgebra. That is,

$$x + iy \iff 1x + e_1 \wedge e_2 y,$$

where $e_1 \wedge e_2$ is the bivector (and psuedoscalar).

Familiar Examples

We have all seen Clifford algebras before. Indeed,

- $\mathbb{C} \cong \text{Cl}(\mathbb{R}, -\cdot)$.
- \mathbb{C} also lives inside of $\text{Cl}(\mathbb{R}^2, \cdot)$ as the even subalgebra. That is,

$$x + iy \iff 1x + e_1 \wedge e_2 y,$$

where $e_1 \wedge e_2$ is the bivector (and psuedoscalar).

- \mathbb{H} lives inside of $\text{Cl}(\mathbb{R}^3, \cdot)$ as the even subalgebra.

Clifford Analysis

Clifford Analysis

- We can replace $d + \delta$ with the Dirac operator D

$$D = \sum_{j=1}^n e_j \frac{\partial}{\partial x^j}.$$

Clifford Analysis

- We can replace $d + \delta$ with the Dirac operator D

$$D = \sum_{j=1}^n e_j \frac{\partial}{\partial x^j}.$$

- $D^2 = \Delta$.

Clifford Analysis

- We can replace $d + \delta$ with the Dirac operator D

$$D = \sum_{j=1}^n e_j \frac{\partial}{\partial x^j}.$$

- $D^2 = \Delta$.
- Elements in the kernel of D are monogenic.

Clifford Analysis

- We can replace $d + \delta$ with the Dirac operator D

$$D = \sum_{j=1}^n e_j \frac{\partial}{\partial x^j}.$$

- $D^2 = \Delta$.
- Elements in the kernel of D are monogenic.
- Monogenic objects in even subalgebra have components that are harmonic.

Clifford Analysis

- We can replace $d + \delta$ with the Dirac operator D

$$D = \sum_{j=1}^n e_j \frac{\partial}{\partial x^j}.$$

- $D^2 = \Delta$.
- Elements in the kernel of D are monogenic.
- Monogenic objects in even subalgebra have components that are harmonic.
- There are Cauchy integral and Hilbert transform type operators for D in arbitrary dimension.

Second Issue

In dimensions greater than 2, even subalgebra is noncommutative.

Second Issue

In dimensions greater than 2, even subalgebra is noncommutative.

- The spectral theory for Belishev's solution required a commutative Banach algebra.

Second Issue

In dimensions greater than 2, even subalgebra is noncommutative.

- The spectral theory for Belishev's solution required a commutative Banach algebra.
- The spectral theory for noncommutative Banach algebras is not as developed.

Second Issue

In dimensions greater than 2, even subalgebra is noncommutative.

- The spectral theory for Belishev's solution required a commutative Banach algebra.
- The spectral theory for noncommutative Banach algebras is not as developed.
- There is still work to do here to find some way around this.

Section 5

Conclusion

Main Story

Main Story

- Calderón proposed a useful and challenging problem for both theorists and practitioners.

Main Story

- Calderón proposed a useful and challenging problem for both theorists and practitioners.
- Advances have been made in both areas.

Main Story

- Calderón proposed a useful and challenging problem for both theorists and practitioners.
- Advances have been made in both areas.
- Ideal results are still not yet obtained.

Novel Approach

Novel Approach

- Belishev solved the 2D problem using the boundary control method.

Novel Approach

- Belishev solved the 2D problem using the boundary control method.
- It relies heavily on complex analysis and the spectral theory for commutative Banach algebras.

Novel Approach

- Belishev solved the 2D problem using the boundary control method.
- It relies heavily on complex analysis and the spectral theory for commutative Banach algebras.
- These issues remain if we try to naively generalize this approach.

New Input

New Input

- Clifford algebras and analysis replace the complex structure in arbitrary dimension.

New Input

- Clifford algebras and analysis replace the complex structure in arbitrary dimension.
- We can still construct the same algebra of holomorphic functions.

New Input

- Clifford algebras and analysis replace the complex structure in arbitrary dimension.
- We can still construct the same algebra of holomorphic functions.
- The relevant algebras are noncommutative for dimensions ≥ 3 .

New Input

- Clifford algebras and analysis replace the complex structure in arbitrary dimension.
- We can still construct the same algebra of holomorphic functions.
- The relevant algebras are noncommutative for dimensions ≥ 3 .
- There are possibly other tools at our disposal that may be able to replace the loss of commutivity.

Thank you!