

# **Riemannian Geometry**

for Dummies

**Colin Roberts**

**Colorado  
State**  
University

# Section 1

## **Introduction**

Riemannian geometry is the study of a *smooth manifold*  $M$  along with a *metric tensor field*  $g$ .

The point of Riemannian geometry is to generalize the differentiable and metric structure of  $\mathbb{R}^n$ .

We generalize to spaces that have interesting topology and geometry.

This will require us to rethink some notions we found “easy” in  $\mathbb{R}^n$ .

But we will gain a very general framework for working with differentiable objects.

## Section 2

### **Motivation**



Why study this in the first place?

Example: Partial differential equations (PDEs) on spaces that are not flat.

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;
- Electrical Impedence Tomography (EIT);

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;
- Electrical Impedence Tomography (EIT);
- General relativity.

Example: Optimization in interesting spaces.

Example: Optimization in interesting spaces.

- Grassmannians;

Example: Optimization in interesting spaces.

- Grassmannians;
- Flags.



## Section 3

### **Preliminaries**

A *topological space*  $M$  is a set of points with a collection of subsets  $\mathcal{O}$  that we define to be open. These sets satisfy

A **topological space**  $M$  is a set of points with a collection of subsets  $\mathcal{O}$  that we define to be open. These sets satisfy

- $\emptyset, M \in \mathcal{O}$ ;

A **topological space**  $M$  is a set of points with a collection of subsets  $\mathcal{O}$  that we define to be open. These sets satisfy

- $\emptyset, M \in \mathcal{O}$ ;
- Arbitrary unions of open sets are open;

A **topological space**  $M$  is a set of points with a collection of subsets  $\mathcal{O}$  that we define to be open.

These sets satisfy

- $\emptyset, M \in \mathcal{O}$ ;
- Arbitrary unions of open sets are open;
- Finite intersections of open sets are open.

- A **homeomorphism** is a continuous bijection  $f: M \rightarrow N$  with continuous inverse  $f^{-1}$ .

- A **homeomorphism** is a continuous bijection  $f: M \rightarrow N$  with continuous inverse  $f^{-1}$ .
- We say  $M$  and  $N$  are homeomorphic.

A *manifold* (with boundary) is a space that locally looks like  $\mathbb{R}^n$  (or  $\mathbb{R}^{n+}$ ).



A **manifold** (with boundary) is a space that locally looks like  $\mathbb{R}^n$  (or  $\mathbb{R}^{n+}$ ).




A **manifold**  $M$  (with boundary) is a topological space such that each open set in  $\mathcal{O}$  is homeomorphic to  $\mathbb{R}^n$  (or  $\mathbb{R}^{n+}$ ).





# Manifolds



## Section 4

# **Applications**

## Section 5

### **Conclusions**