

Riemannian Geometry

for Dummies

Colin Roberts

**Colorado
State**
University

Section 1

Introduction

Riemannian geometry is the study of a *smooth manifold* M along with a *metric tensor field* g .

The point of Riemannian geometry is to generalize the differentiable and metric structure of \mathbb{R}^n .

We generalize to spaces that have interesting topology and geometry.

This will require us to rethink some notions we found “easy” in \mathbb{R}^n .

But we will gain a very general framework for working with differentiable objects.

Section 2

Motivation

Why study this in the first place?

Example: Partial differential equations (PDEs) on spaces that are not flat.

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;
- Electrical Impedence Tomography (EIT);

Example: Partial differential equations (PDEs) on spaces that are not flat.

- Fluid flow on Earth;
- Electrical Impedance Tomography (EIT);
- General relativity.

Example: Optimization in interesting spaces.

Example: Optimization in interesting spaces.


- Grassmannians;


Example: Optimization in interesting spaces.


- Grassmannians;
- Flags.


Section 3


Preliminaries











A *topological space* M is a set of points with a collection of subsets \mathcal{O} that we define to be open. These sets satisfy

A **topological space** M is a set of points with a collection of subsets \mathcal{O} that we define to be open. These sets satisfy

- $\emptyset, M \in \mathcal{O}$;

A **topological space** M is a set of points with a collection of subsets \mathcal{O} that we define to be open. These sets satisfy

- $\emptyset, M \in \mathcal{O}$;
- Arbitrary unions of open sets are open;

A **topological space** M is a set of points with a collection of subsets \mathcal{O} that we define to be open.

These sets satisfy

- $\emptyset, M \in \mathcal{O}$;
- Arbitrary unions of open sets are open;
- Finite intersections of open sets are open.

Just do diffeomorphisms and forget this

- A **homeomorphism** is a continuous bijection $f: M \rightarrow N$ with continuous inverse f^{-1} .

Just do diffeomorphisms and forget this

- A **homeomorphism** is a continuous bijection $f: M \rightarrow N$ with continuous inverse f^{-1} .
- We say M and N are homeomorphic.

ignore boundary

A *manifold* (with boundary) is a space that locally looks like \mathbb{R}^n (or \mathbb{R}^{n+}).


ignore boundary

A *manifold* (with boundary) is a space that locally looks like \mathbb{R}^n (or \mathbb{R}^{n+}).



A *manifold* M (with boundary) is a topological space such that each open set in \mathcal{O} is homeomorphic to \mathbb{R}^n (or \mathbb{R}^{n+}).

Manifolds



Section 4

Applications

Section 5

Conclusions