

# A Multiscale approach to modeling the municipal spread of COVID-19

Colin Roberts

Joint work with

- Elijah Pivo, MIT Institute for Data, Systems, and Society.
- Claire Valva, NYU Courant Center for Atmosphere and Ocean Science.

Note that the phrase,

*“All models are wrong, but some are useful”*

is in play.

# Outline

# Outline

1. Discuss agent, compartmental, and multiscale modeling.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.



# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.
    - i. Class sizes.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.
    - i. Class sizes.
    - ii. Class periods.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.
    - i. Class sizes.
    - ii. Class periods.
  - b. (Compartmental): municipal disease spread.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.
    - i. Class sizes.
    - ii. Class periods.
  - b. (Compartmental): municipal disease spread.
    - i. University-city coupling.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.
    - i. Class sizes.
    - ii. Class periods.
  - b. (Compartmental): municipal disease spread.
    - i. University-city coupling.
  - c. (Multiscale): interactions involving both levels.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.
    - i. Class sizes.
    - ii. Class periods.
  - b. (Compartmental): municipal disease spread.
    - i. University-city coupling.
  - c. (Multiscale): interactions involving both levels.
    - i. Staggered schedules.

# Outline

1. Discuss agent, compartmental, and multiscale modeling.
2. Describe the multiscale modeling approach.
3. Give results
  - a. (Agent): contact rate in the university.
    - i. Class sizes.
    - ii. Class periods.
  - b. (Compartmental): municipal disease spread.
    - i. University-city coupling.
  - c. (Multiscale): interactions involving both levels.
    - i. Staggered schedules.
    - ii. Quarantining/presymptomatic.

# Question



# Question

How do schools and universities impact the spread of COVID-19 in the surrounding community?

# Ideas

# Ideas

- Use an (*non deterministic* and *heterogeneous*) agent based approach.

# Ideas

- Use an (*non deterministic* and *heterogeneous*) agent based approach.
- Use a (*deterministic* and *homogeneous*) compartmental model approach.

# Agent based simulations

# Agent based simulations

- Treat every individual as a single *agent* (entity).

# Agent based simulations

- Treat every individual as a single *agent* (entity).
- Describe every agent's schedule, movement, and infection status at every instant in time.

# Agent based simulations


- Treat every individual as a single *agent* (entity).
- Describe every agent's schedule, movement, and infection status at every instant in time.
- Let agents interact with one another and keep track of the disease progression.



# Examples

# Examples

- Particle based simulations. (3Blue1Brown)



# Examples

# Examples

- Network based simulations. (covasim)



# Benefits

Benefits of agent models:

# Benefits

## Benefits of agent models:

- Heterogeneous social structure.

# Benefits

## Benefits of agent models:

- Heterogeneous social structure.
- Stochastic.

# Benefits

## Benefits of agent models:

- Heterogeneous social structure.
- Stochastic.
- (Typically) less ad-hoc parameter tuning.



# Drawbacks

Drawbacks of agent models:

# Drawbacks

## Drawbacks of agent models:

- Complicated to design.

# Drawbacks

## Drawbacks of agent models:

- Complicated to design.
- Slow to run.

# Drawbacks

## Drawbacks of agent models:

- Complicated to design.
- Slow to run.
- Stochastic nature requires ensembles to generate statistics.

# Compartmental (ODE) based simulations

# Compartmental (ODE) based simulations

- Consider an entire homogeneous population.

# Compartmental (ODE) based simulations

- Consider an entire homogeneous population.
- Ignore individualistic behavior for coarse-grained homogeneity.

# Compartmental (ODE) based simulations


- Consider an entire homogeneous population.
- Ignore individualistic behavior for coarse-grained homogeneity.
- Assume efficient and homogeneous mixing.



# Example

# Example

- SIR Model (Kermack and McKendrick, 1927)



# SIR equations

# SIR equations

We can write *any* ODE as a first order update of a state  $\vec{\mathbf{x}}$  by

$$\dot{\vec{\mathbf{x}}} = \vec{\mathbf{f}}(t, \vec{\mathbf{x}}).$$

# SIR equations

We can write *any* ODE as a first order update of a state  $\vec{\mathbf{x}}$  by

$$\dot{\vec{\mathbf{x}}} = \vec{\mathbf{f}}(t, \vec{\mathbf{x}}).$$

The SIR equations then read

$$\begin{pmatrix} \dot{S} \\ \dot{I} \\ \dot{R} \end{pmatrix} = \begin{pmatrix} -\beta C \frac{I}{N} \\ +\beta C \frac{I}{N} - \gamma I \\ +\gamma I \end{pmatrix},$$


where  $S$ ,  $I$ , and  $R$  denotes the *susceptible*, *infected*, and *removed* populations respectively. Note,  $N = S + I + R$  is the total (conserved) population size.

# Relation to chemistry

The equation

$$\dot{S} = -\beta C \frac{I}{N},$$

can be thought of as a first order chemical reaction




# Relation to chemistry

The equation

$$\dot{S} = -\beta C \frac{I}{N},$$

can be thought of as a first order chemical reaction




- $C$  describes how many contacts with other molecules per unit time species  $S$  experiences.

# Relation to chemistry

The equation

$$\dot{S} = -\beta C \frac{I}{N},$$

can be thought of as a first order chemical reaction



- $C$  describes how many contacts with other molecules per unit time species  $S$  experiences.
- $\frac{I}{N}$  is the proportion of these molecules of the proper type.




# Relation to chemistry

The equation

$$\dot{S} = -\beta C \frac{I}{N},$$

can be thought of as a first order chemical reaction



- $C$  describes how many contacts with other molecules per unit time species  $S$  experiences.
- $\frac{I}{N}$  is the proportion of these molecules of the proper type.
- $\beta$  is the likelihood of reaction.

# Understanding SIR parameters

# Understanding SIR parameters

The parameters  $\beta$ ,  $C$ , and  $\gamma$  can be thought of as:

# Understanding SIR parameters

The parameters  $\beta$ ,  $C$ , and  $\gamma$  can be thought of as:

- $\beta \in [0, 1]$  is likelihood of transmission.

# Understanding SIR parameters

The parameters  $\beta$ ,  $C$ , and  $\gamma$  can be thought of as:

- $\beta \in [0, 1]$  is likelihood of transmission.
- $C \in [0, \infty)$  is the contact rate.

# Understanding SIR parameters

The parameters  $\beta$ ,  $C$ , and  $\gamma$  can be thought of as:

- $\beta \in [0, 1]$  is likelihood of transmission.
- $C \in [0, \infty)$  is the contact rate.
- $\gamma \in [0, \infty)$  is the recovery + death rate.

# Benefits

Benefits of compartmental models:

# Benefits

## Benefits of compartmental models:

- Easy to build and analyze.



# Benefits

## Benefits of compartmental models:

- Easy to build and analyze.
- Quick to compute.

# Benefits

## Benefits of compartmental models:

- Easy to build and analyze.
- Quick to compute.
- Captures large scale behavior.

# Drawbacks

Drawbacks of compartmental models:

# Drawbacks

## Drawbacks of compartmental models:

- Homogeneous.

# Drawbacks

## Drawbacks of compartmental models:

- Homogeneous.
- Deterministic.

# Drawbacks

## Drawbacks of compartmental models:

- Homogeneous.
- Deterministic.
- Ad-hoc parameter changes.

# Multiscale modeling

# Multiscale modeling

- Interesting dynamics for a single system can take place on various spatio-temporal scales.



# Multiscale modeling

- Interesting dynamics for a single system can take place on various spatio-temporal scales.
- Small scale interactions drive large scale phenomenon.

# Multiscale modeling

- Interesting dynamics for a single system can take place on various spatio-temporal scales.
- Small scale interactions drive large scale phenomenon.
- Couple together small and large scale models to study complicated systems.

# Multiscale modeling

- Interesting dynamics for a single system can take place on various spatio-temporal scales.
- Small scale interactions drive large scale phenomenon.
- Couple together small and large scale models to study complicated systems.
- E.g., quantum mechanics  $\rightarrow$  molecular dynamics  $\rightarrow$  kinetic theory  $\rightarrow$  statistical mechanics  $\rightarrow$  thermodynamics.

Idea

# Idea

Can we couple an agent based model alongside a compartmental model to remove the drawbacks and gain benefits?

# Hierarchical compartmental model

# Hierarchical compartmental model

Assume the following:

# Hierarchical compartmental model

Assume the following:

- Two coupled SIR systems  $S_1$ ,  $I_1$ ,  $R_1$ , and  $S_2$ ,  $I_2$ , and  $R_2$ .



# Hierarchical compartmental model


Assume the following:

- Two coupled SIR systems  $S_1$ ,  $I_1$ ,  $R_1$ , and  $S_2$ ,  $I_2$ , and  $R_2$ .
- System 1 refers to the university students and System 2 all other citizens in the city that aren't university members.

# Hierarchical compartmental model

Assume the following:

- Two coupled SIR systems  $S_1, I_1, R_1$ , and  $S_2, I_2$ , and  $R_2$ .
- System 1 refers to the university students and System 2 all other citizens in the city that aren't university members.



# Hierarchical compartmental model

# Hierarchical compartmental model

In general, for  $n$  systems, we have the equations

$$\begin{pmatrix} \dot{S}_i \\ \dot{I}_i \\ \dot{R}_i \end{pmatrix} = \begin{pmatrix} -\beta S_i \sum_j C_{ij} \frac{I_j}{N_j} \\ +\beta S_i \sum_j C_{ij} \frac{I_j}{N_j} - \gamma I_j \\ +\gamma I_j \end{pmatrix}.$$

# Hierarchical compartmental model

In general, for  $n$  systems, we have the equations

$$\begin{pmatrix} \dot{S}_i \\ \dot{I}_i \\ \dot{R}_i \end{pmatrix} = \begin{pmatrix} -\beta S_i \sum_j C_{ij} \frac{I_j}{N_j} \\ +\beta S_i \sum_j C_{ij} \frac{I_j}{N_j} - \gamma I_j \\ +\gamma I_j \end{pmatrix}.$$

In this case, the contact rate  $C$  becomes the **contact matrix**  $C_{ij}$  that describes the contact rate between the systems  $i$  and  $j$ . Note  $C_{ij}$  is symmetric.

# Aggregating systems

# Aggregating systems

- Many systems may comprise a larger system (e.g., System 1 + System 2 from before comprise the entire city).

# Aggregating systems

- Many systems may comprise a larger system (e.g., System 1 + System 2 from before comprise the entire city).
- This decomposition allows us to aggregate larger system dynamics by summing over relevant systems.



# Adding complexity

SIR is lacking. We prefer the SEQIRD compartmental model governed by

$$\begin{pmatrix} \dot{S}_i \\ \dot{E}_i \\ \dot{Q}_i \\ \dot{I}_i \\ \dot{R}_i \\ \dot{D}_i \end{pmatrix} = \begin{pmatrix} -\beta S_i \sum_j C_{ij} \frac{I_j}{N_j} \\ \beta S_i \sum_j C_{ij} \frac{I_j}{N_j} - (\gamma_I - \gamma_Q) E_i \\ \gamma_I E_i - (\lambda + \kappa) I_i \\ \gamma_Q E_i - (\lambda + \kappa) Q_i \\ \lambda(I_i + Q_i) \\ \kappa(I_i + Q_i) \end{pmatrix}$$

- $S \equiv$  susceptible.
- $E \equiv$  exposed.
- $Q \equiv$  quarantined.
- $I \equiv$  infected.
- $R \equiv$  recovered.
- $D \equiv$  dead.

# Adding complexity


SIR is lacking. We prefer the SEQIRD compartmental model governed by

$$\begin{pmatrix} \dot{S}_i \\ \dot{E}_i \\ \dot{Q}_i \\ \dot{I}_i \\ \dot{R}_i \\ \dot{D}_i \end{pmatrix} = \begin{pmatrix} -\beta S_i \sum_j C_{ij} \frac{I_j}{N_j} \\ \beta S_i \sum_j C_{ij} \frac{I_j}{N_j} - (\gamma_I - \gamma_Q) E_i \\ \gamma_I E_i - (\lambda + \kappa) I_i \\ \gamma_Q E_i - (\lambda + \kappa) Q_i \\ \lambda(I_i + Q_i) \\ \kappa(I_i + Q_i) \end{pmatrix}$$

- $\beta = 0.3 \equiv$  transmission rate.
- $C \equiv$  contact matrix.
- $\gamma = 0.07 \equiv$  latent time to infection.
- $\lambda = 0.1 \equiv$  recovery rate.
- $\kappa = 0.002 \equiv$  death rate.
- $q_{\text{percent}} \equiv$  percentage quarantining.
- $\gamma_Q = q_{\text{percent}} \gamma$ .
- $\gamma_I = (1 - q_{\text{percent}}) \gamma$ .


# Determining parameters

# Determining parameters




- All parameters but  $C$  were determined via measurements in other sources.

# Determining parameters



- All parameters but  $C$  were determined via measurements in other sources.
- We determine the values for  $C$  via data assimilation and a agent based model.

# Determining parameters




- All parameters but  $C$  were determined via measurements in other sources.
- We determine the values for  $C$  via data assimilation and a agent based model.
- *This is where the model becomes multiscale.*

# Assimilating data

# Assimilating data


Using measured data from Fort Collins and Larimer County, we estimated a value for  $C$  assuming  $\beta = 1$ .





# Assimilating data

Using measured data from Fort Collins and Larimer County, we estimated a value for  $C$  assuming  $\beta = 1$ .




# Counting contacts

# Counting contacts

**Idea:** Agent based simulator tracks (various types of) contact between students each day.


## Adjustable parameters:

- Student body size.
- Number of class periods.
- Class sizes.
- Students per major.
- Schedule staggering.




# Contacts and school population

- Students are randomly assigned courses of a certain size.
- We count the number of unique contacts a student makes vs. class size for a three class period day.
- The growth is sublinear, but as the population increases we approach linear behavior.
- This is due to diminishing chances of having repeat students in other randomly drawn classes.




# Contacts and major grouping

- To add realism or intervention, we can group students within majors.
- This leads to a decrease in new contacts and sublinear growth due to the effective population size decreasing to the size of the majors (500).



# Contacts and class periods

- Increasing the number of in-person class periods increases contact rate almost linearly when class sizes are small (30).



# Integrating the agent model

# Integrating the agent model

- At the beginning of each day, the agent model is ran to compute a contact rate within the university.



# Integrating the agent model

- At the beginning of each day, the agent model is ran to compute a contact rate within the university.
- This contact rate is used in the compartmental model and we integrate this ODE for a day.

# Integrating the agent model

- At the beginning of each day, the agent model is ran to compute a contact rate within the university.
- This contact rate is used in the compartmental model and we integrate this ODE for a day.
- University members in the Q and D department are removed and the contact rate is recomputed

# City-university coupling

- City-university coupling changes the strength of the off diagonal  $C_{21}$  element.
- With high coupling,  $C_{21} = C_{22}$  and students mix just like typical city members.
- Coupling minimally affects the university, but greatly affects the city.




Figure:  $I(t)$  for class size = 15, major size = 500, and no quarantining.

# University quarantining

- Quarantining only delays infection within the university.
- Significantly alters aggregated city dynamics and prevents second wave.




Figure:  $I(t)$  with 50% quarantine rate, class size of 15, and major size

# Staggering schedules

- Day and week staggering flatten the curve and reduce total infected.
- Week staggering performs slightly better due to latent time for infection.




Figure:  $I(t)$  with staggering, 50% of infected population quarantining,

# Future directions

- Improve and extend the data assimilation for the compartmental model.
- Incorporate different types of contacts.
- Masks, hygiene, and other measures of contact severity.
- Quantify the stochasticity of the agent model.
- More heterogeneity from the agent model (scheduling, transportation, etc.).

# Acknowledgements

