

MATH 271, HOMEWORK 8
DUE NOVEMBER 16TH

Problem 1. Consider the following vectors in space \mathbb{R}^3

$$\vec{u} = \hat{e}_1 + 2\hat{e}_2 + 3\hat{e}_3 \quad \text{and} \quad \vec{v} = -2\hat{e}_1 + \hat{e}_2 - 2\hat{e}_3.$$

- (a) Compute the dot product $\vec{u} \cdot \vec{v}$.
- (b) Compute the lengths $|\vec{u}|$ and $|\vec{v}|$ using the dot product.
- (c) Compute the projection of \vec{u} in the direction of \vec{v} . *Hint: don't forget to normalize the vectors before you build your projection.*
- (d) Compute the cross product $\vec{u} \times \vec{v}$.
- (e) Find the area of the parallelogram generated by \vec{u} and \vec{v} .

Problem 2. Write down the matrix for the following linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$:

$$T \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x + y + z \\ 2x \\ 3y + z \end{pmatrix}.$$

Problem 3. Consider the linear transformation $J: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$J(\hat{e}_1) = \hat{e}_2 \quad \text{and} \quad J(\hat{e}_2) = -\hat{e}_1.$$

This linear transformation is fundamental in understanding how we can reconstruct complex numbers using matrices.

- (a) Show that $J^2 = J \circ J = -1$.
- (b) Determine a matrix representation for J and denote it by $[J]$.
- (c) Recall that we can represent a complex number as $z = x + iy$ and that we can represent z as a vector in \mathbb{R}^2 as $\vec{\zeta} = x\hat{e}_1 + y\hat{e}_2$. Show that $J\vec{\zeta}$ corresponds to iz . *Hint: just show the multiplications are analogous.*
- (d) We can completely reconstruct a representation of \mathbb{C} by using a matrix representation. In particular, we can take

$$[z] = x[I] + y[J].$$

Show that we recover the complex addition and multiplication using this representation.

- (e) We can represent a unit complex number as $z = e^{i\theta}$. Show that the representation described before leads to

$$[z] = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}.$$

Problem 4. Take the following matrices:

$$[A] = \begin{pmatrix} 4 & 3 & 10 & 2 \\ 1 & 1 & 0 & 9 \end{pmatrix}, \quad [B] = \begin{pmatrix} 8 & 5 & 8 \\ 10 & 9 & 2 \\ 4 & 6 & 3 \end{pmatrix}, \quad [C] = \begin{pmatrix} 0 & 0 & 9 \\ 7 & 9 & 9 \\ 1 & 9 & 9 \\ 3 & 3 & 1 \end{pmatrix}$$

- (a) Compute either $[A][C]$ or $[C][A]$ and state which multiplication is not possible.
- (b) Compute either $[B][C]$ or $[C][B]$ and state which multiplication is not possible.
- (c) Can you add any of these matrices?
- (d) Describe each matrix as linear transformation $T: \mathbb{R}^m \rightarrow \mathbb{R}^n$. What is m and n for each? How does this relate to the number of rows and columns?

Problem 5. Solve the following equation.

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 2 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 6 \\ 8 \\ 11 \end{pmatrix}.$$

Problem 6. Consider the space of polynomials of degree at most 3, $P_3(\mathbb{C})$.

- (a) Using the basis

$$B = \{1, x, x^2, x^3\},$$

determine a matrix representation for the linear transformation $\frac{d}{dx}: P_3(\mathbb{C}) \rightarrow P_3(\mathbb{C})$.

- (b) Show that the set of Legendre polynomials

$$B_L = \left\{ f_0 = \sqrt{\frac{1}{2}}, f_1 = \sqrt{\frac{3}{2}}x, f_2 = \sqrt{\frac{5}{8}}(1 - 3x^2), f_3 = \sqrt{\frac{63}{8}} \left(x - \frac{5x^3}{3} \right) \right\}$$

is a basis for $P_3(\mathbb{C})$.

- (c) Using the basis B_L instead, compute a matrix representation for the linear transformation $\frac{d}{dx}$.

Remark 1. This should go to show you that a matrix representation depends on a basis!!!

Problem 7. Let $C^\omega(\mathbb{C})$ be the set of analytic functions (functions that have a power series representation), i.e., functions of the form

$$f(x) = \sum_{n=0}^{\infty} a_n x^n,$$

where $a_n \in \mathbb{C}$ for $n = 0, 1, 2, \dots$. Let us compare this with the vector space of polynomials $P_N(\mathbb{C})$.

- (a) Argue that $C^\omega(\mathbb{C})$ is a vector space. *Hint: show what addition and scalar multiplication look like.*
- (b) Show that the space of polynomials of degree at most N , $P_N(\mathbb{C})$ is a subspace of $C^\omega(\mathbb{C})$.
- (c) Let $f, g \in C^\omega(\mathbb{C})$ be given by

$$f(x) = \sum_{n=0}^{\infty} a_n x^n \quad \text{and} \quad g(x) = \sum_{n=0}^{\infty} b_n x^n,$$

then define an inner product on $C^\omega(\mathbb{C})$ by taking

$$\langle f, g \rangle := \sum_{n=0}^{\infty} a_n b_n^*.$$

Now, write $h(x)$ as a Taylor series centered at $x = 0$ and show that

$$h^{(n)}(0) = n! \langle h, x^n \rangle.$$

- (d) Show that the N^{th} order Taylor approximation for the function $h(x)$ centered at $x = 0$ is the projection onto the subspace spanned by the functions

$$S = \{1, x, x^2, \dots, x^N\}.$$

This projection is given by

$$\text{proj}_S(h) = \sum_{n=0}^N \langle h, x^n \rangle x^n.$$