


Lecture 11: KKT Conditions

Shi Pu

School of Data Science (SDS)
The Chinese University of Hong Kong, Shenzhen



- 1 KKT Conditions
- 2 The Convex Case
- 3 Constrained Least Squares
- 4 Second Order Optimality Conditions
- 5 Optimality Conditions for the Trust Region Subproblem
- 6 Total Least Squares

- 1 KKT Conditions
- 2 The Convex Case
- 3 Constrained Least Squares
- 4 Second Order Optimality Conditions
- 5 Optimality Conditions for the Trust Region Subproblem
- 6 Total Least Squares

The Karush-Kuhn-Tucker Conditions

- The Karush-Kuhn-Tucker conditions are optimality conditions for inequality constrained problems discovered in 1951 (originating from Karush's thesis from 1939).
- Modern nonlinear optimization essentially begins with the discovery of these conditions.

The basic notion that we will require is the one of **feasible descent directions**.

Definition

Consider the problem

$$\begin{array}{ll}\min & f(\mathbf{x}) \\ \text{s.t.} & \mathbf{x} \in C\end{array}$$

where f is continuously differentiable over the set $C \subset \mathbb{R}^n$. Then a vector $\mathbf{d} \neq \mathbf{0}$ is called a **feasible descent direction** at $\mathbf{x} \in C$ if $\nabla f(\mathbf{x})^\top \mathbf{d} < 0$ and there exists $\epsilon > 0$ such that $\mathbf{x} + t\mathbf{d} \in C$ for all $t \in [0, \epsilon]$.

The Basic Necessary Condition - No Feasible Descent Directions

Lemma

Consider the problem

$$(P) \quad \begin{array}{ll} \min & f(\mathbf{x}) \\ \text{s.t.} & \mathbf{x} \in C \end{array}$$

where f is continuously differentiable over the set C . If \mathbf{x}^* is a local optimal solution of (P), then there are no feasible descent directions at \mathbf{x}^* .

Proof.

- By contradiction, assume that there exists a vector \mathbf{d} and $\epsilon_1 > 0$ such that $\mathbf{x} + t\mathbf{d} \in C$ for all $t \in [0, \epsilon_1]$ and $\nabla f(\mathbf{x}^*)^\top \mathbf{d} < 0$.
- By definition of the directional derivative there exists $\epsilon_2 < \epsilon_1$ such that $f(\mathbf{x}^* + t\mathbf{d}) < f(\mathbf{x}^*)$ for all $t \in [0, \epsilon_2] \Rightarrow$ contradiction to the local optimality of \mathbf{x}^* .



Lemma

Let \mathbf{x}^* be a local minimum of the problem

$$\begin{array}{ll}\min & f(\mathbf{x}) \\ \text{s.t.} & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m\end{array}$$

where f, g_1, \dots, g_m are continuously differentiable over \mathbb{R}^n . Let $I(\mathbf{x}^*)$ be the set of active constraints at \mathbf{x}^* :

$$I(\mathbf{x}^*) = \{i : g_i(\mathbf{x}^*) = 0\}$$

Then there does not exist a vector $\mathbf{d} \in \mathbb{R}^n$ such that

$$\nabla f(\mathbf{x}^*)^\top \mathbf{d} < 0$$

$$\nabla g_i(\mathbf{x}^*)^\top \mathbf{d} < 0, \quad i \in I(\mathbf{x}^*)$$

Proof.

- Suppose that \mathbf{d} satisfies the system of inequalities.
- Then $\exists \epsilon_1 > 0$ such that $f(\mathbf{x}^* + t\mathbf{d}) < f(\mathbf{x}^*)$ and $g_i(\mathbf{x}^* + t\mathbf{d}) < g_i(\mathbf{x}^*) = 0$ for any $t \in (0, \epsilon_1)$ and $i \in I(\mathbf{x}^*)$.
- For any $i \notin I(\mathbf{x}^*)$ we have that $g_i(\mathbf{x}^*) < 0$, and hence, by the continuity of g_i , there exists $\epsilon_2 > 0$ such that $g_i(\mathbf{x}^* + t\mathbf{d}) < 0$ for any $t \in (0, \epsilon_2)$ and $i \notin I(\mathbf{x}^*)$.
- Consequently,

$$\begin{aligned} f(\mathbf{x}^* + t\mathbf{d}) &< f(\mathbf{x}^*) \\ g_i(\mathbf{x}^* + t\mathbf{d}) &< 0, \quad i = 1, 2, \dots, m \end{aligned}$$

for all $t \in (0, \min\{\epsilon_1, \epsilon_2\})$

- Contradiction to the local optimality of \mathbf{x}^* .



Theorem

Let \mathbf{x}^* be a local minimum of the problem

$$\begin{array}{ll}\min & f(\mathbf{x}) \\ \text{s.t.} & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m\end{array}$$

where f, g_1, \dots, g_m are continuously differentiable functions over \mathbb{R}^n . Then there exist multipliers $\lambda_0, \lambda_1, \dots, \lambda_m \geq 0$, which are not all zeros, such that

$$\begin{aligned}\lambda_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) &= \mathbf{0} \\ \lambda_i g_i(\mathbf{x}^*) &= 0, \quad i = 1, 2, \dots, m\end{aligned}$$

Proof.

- The following system is infeasible

$$(S) \quad \nabla f(\mathbf{x}^*)^\top \mathbf{d} < 0, \quad \nabla g_i(\mathbf{x}^*)^\top \mathbf{d} < 0, \quad i \in I(\mathbf{x}^*) = \{i_1, \dots, i_k\}$$

- System (S) is the same as $\mathbf{A}\mathbf{d} < 0$ where $\mathbf{A} = (\nabla f(\mathbf{x}^*), \nabla g_{i_1}(\mathbf{x}^*), \dots, \nabla g_{i_k}(\mathbf{x}^*))^\top$
- By Gordan's theorem of alternative, system (S) is infeasible if and only if there exists a vector $\eta = (\lambda_0, \lambda_{i_1}, \dots, \lambda_{i_k})^\top \neq \mathbf{0}$ such that $\mathbf{A}^\top \eta = \mathbf{0}$, $\eta \geq \mathbf{0}$, which is the same as $\lambda_0 \nabla f(\mathbf{x}^*) + \sum_{i \in I(\mathbf{x}^*)} \lambda_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$.
- Define $\lambda_i = 0$ for any $i \notin I(\mathbf{x}^*)$, and we obtain that

$$\lambda_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}, \quad \lambda_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m$$



The KKT Conditions for Inequality Constrained Problems

A major drawback of the Fritz-John conditions is, they allow λ_0 to be zero. Under an additional **regularity** condition, we can assume $\lambda_0 = 1$.

Theorem

Let \mathbf{x}^* be a local minimum of the problem

$$\begin{array}{ll}\min & f(\mathbf{x}) \\ \text{s.t.} & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m\end{array}$$

where f, g_1, \dots, g_m are continuously differentiable functions over \mathbb{R}^n . Suppose that the gradients of the active constraints $\{\nabla g_i(\mathbf{x}^*)\}_{i \in I(\mathbf{x}^*)}$ are **linearly independent**. Then there exist multipliers $\lambda_1, \lambda_2, \dots, \lambda_m \geq 0$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$$

$$\lambda_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m$$

Proof.

- By the Fritz-John conditions, there exists $\tilde{\lambda}_0, \tilde{\lambda}_1, \dots, \tilde{\lambda}_m$, not all zeros, such that

$$\tilde{\lambda}_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m \tilde{\lambda}_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$$

$$\tilde{\lambda}_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m$$

- $\tilde{\lambda}_0 \neq 0$ since otherwise, if $\tilde{\lambda}_0 = 0$, $\sum_{i \in I(\mathbf{x}^*)} \tilde{\lambda}_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}$, where not all the scalars $\tilde{\lambda}_i, i \in I(\mathbf{x}^*)$ are zeros, which is a contradiction to the regularity condition.
- $\tilde{\lambda}_0 > 0$. Defining $\lambda_i = \frac{\tilde{\lambda}_i}{\tilde{\lambda}_0}$, the result follows.



Theorem

Let \mathbf{x}^* be a local minimum of the problem

$$\begin{aligned} \min \quad & f(\mathbf{x}) \\ \text{s.t.} \quad & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \\ & h_j(\mathbf{x}) = 0, \quad j = 1, 2, \dots, p \end{aligned} \quad (1)$$

where $f, g_1, \dots, g_m, h_1, h_2, \dots, h_p$ are continuously differentiable functions over \mathbb{R}^n . Suppose that the gradients of the active constraints and the equality constraints: $\{\nabla g_i(\mathbf{x}^*), \nabla h_j(\mathbf{x}^*), i \in I(\mathbf{x}^*), j = 1, 2, \dots, p\}$ are *linearly independent*. Then there exist multipliers $\lambda_1, \dots, \lambda_m \geq 0, \mu_1, \mu_2, \dots, \mu_p \in \mathbb{R}$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j \nabla h_j(\mathbf{x}^*) = \mathbf{0}$$

$$\lambda_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m$$

Definition (KKT point)

Consider problem (1) where $g_1, \dots, g_m, h_1, h_2, \dots, h_p$ are continuously differentiable functions over \mathbb{R}^n . A feasible point \mathbf{x}^* is called a KKT point if there exist $\lambda_1, \dots, \lambda_m \geq 0, \mu_1, \mu_2, \dots, \mu_p \in \mathbb{R}$ such that

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j \nabla h_j(\mathbf{x}^*) = \mathbf{0}$$

$$\lambda_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m$$

Definition (Regularity)

A feasible point \mathbf{x}^* is called regular if the set $\{\nabla g_i(\mathbf{x}^*), \nabla h_j(\mathbf{x}^*), i \in I(\mathbf{x}^*), j = 1, 2, \dots, p\}$ is linearly independent.

- The KKT theorem states that a necessary local optimality condition of a regular point is that it is a KKT point.
- The additional requirement of regularity is not required in linearly constrained problems in which no such assumption is needed.

1.

$$\begin{array}{ll}\min & x_1 + x_2 \\ \text{s.t.} & x_1^2 + x_2^2 = 1\end{array}$$

2.

$$\begin{array}{ll}\min & x_1 + x_2 \\ \text{s.t.} & (x_1^2 + x_2^2 - 1)^2 = 0\end{array}$$

In class.

- 1 KKT Conditions
- 2 The Convex Case
- 3 Constrained Least Squares
- 4 Second Order Optimality Conditions
- 5 Optimality Conditions for the Trust Region Subproblem
- 6 Total Least Squares

Sufficiency of KKT Conditions in the Convex Case

In the convex case the KKT conditions are **always** sufficient.

Theorem

Let \mathbf{x}^* be a feasible solution of the problem

$$\begin{aligned} \min \quad & f(\mathbf{x}) \\ \text{s.t.} \quad & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \\ & h_j(\mathbf{x}) = 0, \quad j = 1, 2, \dots, p \end{aligned} \tag{2}$$

where f, g_1, \dots, g_m are continuously differentiable **convex** functions over \mathbb{R}^n and h_1, h_2, \dots, h_p are affine functions. Suppose that there exist multipliers $\lambda_1, \dots, \lambda_m \geq 0, \mu_1, \mu_2, \dots, \mu_p \in \mathbb{R}$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j \nabla h_j(\mathbf{x}^*) = \mathbf{0}$$

$$\lambda_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m$$

Then \mathbf{x}^* is the optimal solution of (2).

Proof.

- Let \mathbf{x} be a feasible solution of (2). We will show that $f(\mathbf{x}) \geq f(\mathbf{x}^*)$.
- The function $s(\mathbf{x}) = f(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j h_j(\mathbf{x})$ is convex.
- Since $\nabla s(\mathbf{x}^*) = \nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j \nabla h_j(\mathbf{x}^*) = \mathbf{0}$, it follows that \mathbf{x}^* is a minimizer of s over \mathbb{R}^n , and in particular $s(\mathbf{x}^*) \leq s(\mathbf{x})$.
- Thus,

$$\begin{aligned} f(\mathbf{x}^*) &= f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}^*) + \sum_{j=1}^p \mu_j h_j(\mathbf{x}^*) \\ &= s(\mathbf{x}^*) \\ &\leq s(\mathbf{x}) \\ &= f(\mathbf{x}) + \sum_{i=1}^m \lambda_i g_i(\mathbf{x}) + \sum_{j=1}^p \mu_j h_j(\mathbf{x}) \leq f(\mathbf{x}) \end{aligned}$$



Convex Constraints - Necessity under Slater's Condition

If constraints are convex, regularity can be replaced by [Slater's condition](#).

Theorem

Let \mathbf{x}^* be a local minimum of the problem

$$\begin{array}{ll} \min & f(\mathbf{x}) \\ \text{s.t.} & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \end{array} \quad (3)$$

where f, g_1, \dots, g_m are continuously differentiable functions over \mathbb{R}^n . In addition, g_1, \dots, g_m are [convex](#) over \mathbb{R}^n . Suppose $\exists \hat{\mathbf{x}} \in \mathbb{R}^n$ such that

$$g_i(\hat{\mathbf{x}}) < 0, \quad i = 1, 2, \dots, m$$

Then there exist multipliers $\lambda_1, \dots, \lambda_m \geq 0$, such that

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) = \mathbf{0} \quad (4)$$

$$\lambda_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m \quad (5)$$

Proof.

- Since \mathbf{x}^* is an optimal solution of (3), the Fritz-John conditions are satisfied: there exist $\tilde{\lambda}_0, \dots, \tilde{\lambda}_m \geq 0$ not all zeros, such that

$$\begin{aligned}\tilde{\lambda}_0 \nabla f(\mathbf{x}^*) + \sum_{i=1}^m \tilde{\lambda}_i \nabla g_i(\mathbf{x}^*) &= \mathbf{0} \\ \tilde{\lambda}_i g_i(\mathbf{x}^*) &= 0, \quad i = 1, 2, \dots, m\end{aligned}\tag{6}$$

- We will prove that $\tilde{\lambda}_0 > 0$, and then conditions (4) and (5) will be satisfied with $\lambda_i = \frac{\tilde{\lambda}_i}{\tilde{\lambda}_0}, i = 1, 2, \dots, m$.
- Assume in contradiction that $\tilde{\lambda}_0 = 0$. Then

$$\sum_{i=1}^m \tilde{\lambda}_i \nabla g_i(\mathbf{x}^*) = \mathbf{0}\tag{7}$$

Proof Contd.

- By the gradient inequality,

$$0 > g_i(\hat{\mathbf{x}}) \geq g_i(\mathbf{x}^*) + \nabla g_i(\mathbf{x}^*)^\top (\hat{\mathbf{x}} - \mathbf{x}^*), \quad i = 1, 2, \dots, m$$

- Multiplying the i -th equation by $\tilde{\lambda}_i$ and summing over $i = 1, 2, \dots, m$ we obtain

$$0 > \sum_{i=1}^m \tilde{\lambda}_i g_i(\mathbf{x}^*) + \left[\sum_{i=1}^m \tilde{\lambda}_i \nabla g_i(\mathbf{x}^*) \right]^\top (\hat{\mathbf{x}} - \mathbf{x}^*) \quad (8)$$

- Plugging the identities (7) and (6) into (8) we obtain the impossible statement that $0 > 0$, thus establishing the result.



1.

$$\begin{array}{ll}\min & x_1^2 - x_2 \\ \text{s.t.} & x_2 = 0\end{array}$$

2.

$$\begin{array}{ll}\min & x_1^2 - x_2 \\ \text{s.t.} & x_2^2 \leq 0\end{array}$$

The optimal solution is $(x_1, x_2) = (0, 0)$. Satisfies KKT conditions for problem 1, but not for problem 2.

In class.

Definition (Generalized Slater's Condition)

Consider the system

$$g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m$$

$$h_j(\mathbf{x}) \leq 0, \quad j = 1, 2, \dots, p$$

$$s_k(\mathbf{x}) = 0, \quad k = 1, 2, \dots, q$$

where $g_i, i = 1, 2, \dots, m$ are **convex** functions over \mathbb{R}^n and $h_j, s_k, j = 1, 2, \dots, p, k = 1, 2, \dots, q$ are affine functions over \mathbb{R}^n . Then we say that the **generalized Slater's condition** is satisfied if there exists $\hat{\mathbf{x}} \in \mathbb{R}^n$ for which

$$g_i(\hat{\mathbf{x}}) < 0, \quad i = 1, 2, \dots, m$$

$$h_j(\hat{\mathbf{x}}) \leq 0, \quad j = 1, 2, \dots, p$$

$$s_k(\hat{\mathbf{x}}) = 0, \quad k = 1, 2, \dots, q$$

Theorem

Let \mathbf{x}^* be an optimal solution of the problem

$$\begin{aligned} \min \quad & f(\mathbf{x}) \\ \text{s.t.} \quad & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \\ & h_j(\mathbf{x}) \leq 0, \quad j = 1, 2, \dots, p \\ & s_k(\mathbf{x}) = 0, \quad k = 1, 2, \dots, q \end{aligned} \tag{9}$$

where f, g_1, \dots, g_m are continuously differentiable functions and $\{g_i\}$ are **convex** functions over \mathbb{R}^n , and $\{h_j\}, \{s_k\}$ are affine. Suppose the generalized Slater's condition holds. Then there exist multipliers $\lambda_1, \lambda_2, \dots, \lambda_m, \eta_1, \eta_2, \dots, \eta_p \geq 0, \mu_1, \mu_2, \dots, \mu_q \in \mathbb{R}$ such that

$$\nabla f(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^p \eta_j \nabla h_j(\mathbf{x}^*) + \sum_{k=1}^q \mu_k \nabla s_k(\mathbf{x}^*) = \mathbf{0}$$

$$\lambda_i g_i(\mathbf{x}^*) = 0, \quad i = 1, 2, \dots, m$$

$$\eta_j h_j(\mathbf{x}^*) = 0, \quad j = 1, 2, \dots, p$$

Example

$$\begin{array}{ll}\min & 4x_1^2 + x_2^2 - x_1 - 2x_2 \\ \text{s.t.} & 2x_1 + x_2 \leq 1 \\ & x_1^2 \leq 1\end{array}$$

In class.

- 1 KKT Conditions
- 2 The Convex Case
- 3 Constrained Least Squares
- 4 Second Order Optimality Conditions
- 5 Optimality Conditions for the Trust Region Subproblem
- 6 Total Least Squares

$$\begin{aligned} (CLS) \quad & \min \quad \|\mathbf{Ax} - \mathbf{b}\|^2 \\ & \text{s.t.} \quad \|\mathbf{x}\|^2 \leq \alpha \end{aligned}$$

where $\mathbf{A} \in \mathbb{R}^{m \times n}$ has full column rank, $\mathbf{b} \in \mathbb{R}^m$, $\alpha > 0$.

- Problem (CLS) is a convex problem and satisfies Slater's condition.
- Lagrangian: $L(\mathbf{x}, \lambda) = \|\mathbf{Ax} - \mathbf{b}\|^2 + \lambda(\|\mathbf{x}\|^2 - \alpha)$ ($\lambda \geq 0$)
- KKT conditions:

$$\begin{aligned} \nabla_{\mathbf{x}} L &= 2\mathbf{A}^T(\mathbf{Ax} - \mathbf{b}) + 2\lambda\mathbf{x} = 0 \\ \lambda(\|\mathbf{x}\|^2 - \alpha) &= 0 \\ \|\mathbf{x}\|^2 &\leq \alpha, \quad \lambda \geq 0 \end{aligned}$$

- If $\lambda = 0$, then by the first equation

$$\mathbf{x} = \mathbf{x}_{LS} \equiv (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}.$$

Optimal iff $\|\mathbf{x}_{LS}\|^2 \leq \alpha$.

- On the other hand, if $\|\mathbf{x}_{LS}\|^2 > \alpha$, then necessarily $\lambda > 0$. By the C-S condition we have that $\|\mathbf{x}\|^2 = \alpha$ and the first equation implies that

$$\mathbf{x} = \mathbf{x}_\lambda \equiv (\mathbf{A}^\top \mathbf{A} + \lambda \mathbf{I})^{-1} \mathbf{A}^\top \mathbf{b}$$

The multiplier $\lambda > 0$ should be chosen to satisfy $\|\mathbf{x}_\lambda\|^2 = \alpha$, that is, λ is the solution of

$$f(\lambda) = \|(\mathbf{A}^\top \mathbf{A} + \lambda \mathbf{I})^{-1} \mathbf{A}^\top \mathbf{b}\|^2 - \alpha = 0$$

- $f(0) = \|(\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{b}\|^2 - \alpha = \|\mathbf{x}_{LS}\|^2 - \alpha > 0$, f strictly decreasing and $f(\lambda) \rightarrow -\alpha$ as $\lambda \rightarrow \infty$.
- Conclusion: the optimal solution of the CLS problem is given by

$$\mathbf{x} = \begin{cases} \mathbf{x}_{LS} & \|\mathbf{x}_{LS}\|^2 \leq \alpha \\ (\mathbf{A}^\top \mathbf{A} + \lambda \mathbf{I})^{-1} \mathbf{A}^\top \mathbf{b} & \|\mathbf{x}_{LS}\|^2 > \alpha \end{cases}$$

where λ is the unique root of $f(\lambda)$ over $(0, \infty)$.

- 1 KKT Conditions
- 2 The Convex Case
- 3 Constrained Least Squares
- 4 Second Order Optimality Conditions**
- 5 Optimality Conditions for the Trust Region Subproblem
- 6 Total Least Squares

Theorem

Consider the problem

$$\begin{aligned} \min \quad & f(\mathbf{x}) \\ \text{s.t.} \quad & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \end{aligned}$$

where f, g_1, \dots, g_m are twice continuously differentiable functions over \mathbb{R}^n . Let \mathbf{x}^ be a local minimum, and suppose that \mathbf{x}^* is regular meaning that $\{\nabla g_i(\mathbf{x}^*)\}_{i \in I(\mathbf{x}^*)}$ are linearly independent. Then $\exists \lambda_1, \lambda_2, \dots, \lambda_m \geq 0$ such that*

$$\begin{aligned} \nabla_{\mathbf{x}} L(\mathbf{x}^*, \lambda) &= \mathbf{0} \\ \lambda_i g_i(\mathbf{x}^*) &= 0, \quad i = 1, 2, \dots, m \end{aligned}$$

and $\mathbf{d}^\top \nabla_{\mathbf{xx}}^2 L(\mathbf{x}^, \lambda) \mathbf{d} \geq 0$ for all $\mathbf{d} \in \Lambda(\mathbf{x}^*)$ where*

$$\Lambda(\mathbf{x}^*) \equiv \{\mathbf{d} \in \mathbb{R}^n : \nabla g_i(\mathbf{x}^*)^\top \mathbf{d} = 0, \quad i \in I(\mathbf{x}^*)\}.$$

See the proof of Theorem 11.18 in the textbook.

Second Order Necessary Optimality Conditions for Inequality/Equality Constrained Problems

Theorem

Consider the problem

$$\begin{aligned} \min \quad & f(\mathbf{x}) \\ \text{s.t.} \quad & g_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \\ & h_j(\mathbf{x}) = 0, \quad j = 1, 2, \dots, p \end{aligned}$$

where $f, g_1, \dots, g_m, h_1, \dots, h_p$ are twice continuously differentiable functions. Let \mathbf{x}^ be a local minimum and suppose that \mathbf{x}^* is regular meaning that the set $\{\nabla g_i(\mathbf{x}^*), \nabla h_j(\mathbf{x}^*), i \in I(\mathbf{x}^*), j = 1, 2, \dots, p\}$ is linearly independent. Then $\exists \lambda_1, \lambda_2, \dots, \lambda_m \geq 0$ and $\mu_1, \mu_2, \dots, \mu_p \in \mathbb{R}$ such that*

$$\begin{aligned} \nabla_{\mathbf{x}} L(\mathbf{x}^*, \lambda, \mu) &= \mathbf{0} \\ \lambda_i g_i(\mathbf{x}^*) &= 0, \quad i = 1, 2, \dots, m \end{aligned}$$

and $\mathbf{d}^\top \nabla_{\mathbf{xx}}^2 L(\mathbf{x}^, \lambda, \mu) \mathbf{d} \geq 0$ for all $\mathbf{d} \in \Lambda(\mathbf{x}^*) \equiv \{\mathbf{d} \in \mathbb{R}^n : \nabla g_i(\mathbf{x}^*)^\top \mathbf{d} = 0, \nabla h_j(\mathbf{x}^*)^\top \mathbf{d} = 0, i \in I(\mathbf{x}^*), j = 1, 2, \dots, p\}$.*

- 1 KKT Conditions
- 2 The Convex Case
- 3 Constrained Least Squares
- 4 Second Order Optimality Conditions
- 5 Optimality Conditions for the Trust Region Subproblem**
- 6 Total Least Squares

Optimality Conditions for the Trust Region Subproblem

The Trust Region Subproblem (TRS) is the problem consisting of minimizing an indefinite quadratic function subject to an l_2 -norm constraint:

$$(TRS) : \quad \min\{f(\mathbf{x}) \equiv \mathbf{x}^\top \mathbf{A} \mathbf{x} + 2\mathbf{b}^\top \mathbf{x} + c : \|\mathbf{x}\|^2 \leq \alpha\}$$

where $\mathbf{A} = \mathbf{A}^\top \in \mathbb{R}^{n \times n}$, $\mathbf{b} \in \mathbb{R}^n$ and $c \in \mathbb{R}$. Although the problem is nonconvex, it possesses necessary and sufficient optimality conditions.

Theorem

A vector \mathbf{x}^ is an optimal solution of problem (TRS) if and only if there exists $\lambda^* \geq 0$ such that*

$$(\mathbf{A} + \lambda^* \mathbf{I})\mathbf{x}^* = -\mathbf{b} \tag{10}$$

$$\|\mathbf{x}^*\|^2 \leq \alpha \tag{11}$$

$$\lambda^*(\|\mathbf{x}^*\|^2 - \alpha) = 0 \tag{12}$$

$$\mathbf{A} + \lambda^* \mathbf{I} \succeq \mathbf{0} \tag{13}$$

Proof.

Sufficiency:

- Assume that \mathbf{x}^* satisfies (10)-(13) for some $\lambda^* \geq 0$.
- Define the function

$$\begin{aligned} h(\mathbf{x}) &= \mathbf{x}^\top \mathbf{A} \mathbf{x} + 2\mathbf{b}^\top \mathbf{x} + c + \lambda^*(\|\mathbf{x}\|^2 - \alpha) \\ &= \mathbf{x}^\top (\mathbf{A} + \lambda^* \mathbf{I}) \mathbf{x} + 2\mathbf{b}^\top \mathbf{x} + c - \alpha \lambda^* \end{aligned} \quad (14)$$

- Then by (13) we have that h is a convex quadratic function. By (10) it follows that $\nabla h(\mathbf{x}^*) = 0$, which implies that \mathbf{x}^* is the unconstrained minimizer of h over \mathbb{R}^n .
- Let \mathbf{x} be a feasible point, i.e., $\|\mathbf{x}\|^2 \leq \alpha$. Then

$$\begin{aligned} f(\mathbf{x}) &\geq f(\mathbf{x}) + \lambda^*(\|\mathbf{x}\|^2 - \alpha) && (\lambda^* \geq 0, \|\mathbf{x}\|^2 - \alpha \leq 0) \\ &= h(\mathbf{x}) && \text{(by (14))} \\ &\geq h(\mathbf{x}^*) && (\mathbf{x}^* \text{ is the minimizer of } h) \\ &= f(\mathbf{x}^*) + \lambda^*(\|\mathbf{x}^*\|^2 - \alpha) \\ &= f(\mathbf{x}^*) && \text{(by (12))} \end{aligned}$$

Proof Contd.

Necessity:

- Regularity always satisfied. Suppose \mathbf{x}^* is a minimizer of (TRS). If $\|\mathbf{x}^*\|^2 < \alpha$, done. If $\|\mathbf{x}^*\|^2 = \alpha$, by SONC, $\exists \lambda^* \geq 0$ such that

$$(\mathbf{A} + \lambda^* \mathbf{I})\mathbf{x}^* = -\mathbf{b} \quad (15)$$

$$\|\mathbf{x}^*\|^2 \leq \alpha \quad (16)$$

$$\lambda^*(\|\mathbf{x}^*\|^2 - \alpha) = 0 \quad (17)$$

$$\mathbf{d}^\top (\mathbf{A} + \lambda^* \mathbf{I}) \mathbf{d} \geq 0 \quad \text{for all } \mathbf{d} \text{ satisfying } \mathbf{d}^\top \mathbf{x}^* = 0 \quad (18)$$

- Need to show that (18) is true for any \mathbf{d} .
- Suppose on the contrary: $\exists \mathbf{d}$ s.t. $\mathbf{d}^\top \mathbf{x}^* > 0$ and $\mathbf{d}^\top (\mathbf{A} + \lambda^* \mathbf{I}) \mathbf{d} < 0$.
- Consider $\bar{\mathbf{x}} = \mathbf{x}^* + t\mathbf{d}$, where $t = -2\frac{\mathbf{d}^\top \mathbf{x}^*}{\|\mathbf{d}\|^2}$. $\bar{\mathbf{x}}$ is feasible since

$$\begin{aligned} \|\bar{\mathbf{x}}\|^2 &= \|\mathbf{x}^* + t\mathbf{d}\|^2 = \|\mathbf{x}^*\|^2 + 2t\mathbf{d}^\top \mathbf{x}^* + t^2\|\mathbf{d}\|^2 \\ &= \|\mathbf{x}^*\|^2 - 4\frac{(\mathbf{d}^\top \mathbf{x}^*)^2}{\|\mathbf{d}\|^2} + 4\frac{(\mathbf{d}^\top \mathbf{x}^*)^2}{\|\mathbf{d}\|^2} = \|\mathbf{x}^*\|^2 \leq \alpha. \end{aligned}$$

Proof Contd.

■ In addition,

$$\begin{aligned}
 f(\bar{\mathbf{x}}) &= \bar{\mathbf{x}}^\top \mathbf{A} \bar{\mathbf{x}} + 2\mathbf{b}^\top \bar{\mathbf{x}} + c \\
 &= (\mathbf{x}^* + t\mathbf{d})^\top \mathbf{A}(\mathbf{x}^* + t\mathbf{d}) + 2\mathbf{b}^\top (\mathbf{x}^* + t\mathbf{d}) + c \\
 &= \underbrace{(\mathbf{x}^*)^\top \mathbf{A} \mathbf{x}^* + 2\mathbf{b}^\top \mathbf{x}^* + c}_{f(\mathbf{x}^*)} + t^2 \mathbf{d}^\top \mathbf{A} \mathbf{d} + 2t\mathbf{d}^\top (\mathbf{A} \mathbf{x}^* + \mathbf{b}) \\
 &= f(\mathbf{x}^*) + t^2 \mathbf{d}^\top (\mathbf{A} + \lambda^* \mathbf{I}) \mathbf{d} + 2t\mathbf{d}^\top \underbrace{((\mathbf{A} + \lambda^* \mathbf{I}) \mathbf{x}^* + \mathbf{b})}_{=0 \text{ by (15)}} \\
 &\quad - \lambda^* t \underbrace{[t\|\mathbf{d}\|^2 + 2\mathbf{d}^\top \mathbf{x}^*]}_{=0} \\
 &= f(\mathbf{x}^*) + t^2 \mathbf{d}^\top (\mathbf{A} + \lambda^* \mathbf{I}) \mathbf{d} \\
 &< f(\mathbf{x}^*)
 \end{aligned}$$

which is a contradiction to the optimality of \mathbf{x}^* .



- 1 KKT Conditions
- 2 The Convex Case
- 3 Constrained Least Squares
- 4 Second Order Optimality Conditions
- 5 Optimality Conditions for the Trust Region Subproblem
- 6 Total Least Squares

Total Least Squares

Consider the approximate set of linear equations:

$$\mathbf{Ax} \approx \mathbf{b}$$

- In the **Least Squares (LS)** approach we only assume \mathbf{b} is subjected to noise.

$$\min_{\mathbf{w}, \mathbf{x}} \quad \|\mathbf{w}\|^2$$

$$\text{s.t.} \quad \mathbf{Ax} = \mathbf{b} + \mathbf{w}$$

$$\mathbf{w} \in \mathbb{R}^m$$

- In the **Total Least Squares (TLS)** we assume that both the RHS vector \mathbf{b} and the model matrix \mathbf{A} are subjected to noise

$$\min_{\mathbf{E}, \mathbf{w}, \mathbf{x}} \quad \|\mathbf{E}\|_F^2 + \|\mathbf{w}\|^2$$

$$(TLS) \quad \text{s.t.} \quad (\mathbf{A} + \mathbf{E})\mathbf{x} = \mathbf{b} + \mathbf{w}$$

$$\mathbf{E} \in \mathbb{R}^{m \times n}, \mathbf{w} \in \mathbb{R}^m$$

The TLS problem – as formulated – seems like a difficult nonconvex problem. Will see that it can be solved efficiently.

Eliminating the \mathbf{E} and \mathbf{w} variables

- Fixing \mathbf{x} , we will solve the problem

$$(P_{\mathbf{x}}) \quad \begin{array}{ll} \min_{\mathbf{E}, \mathbf{w}} & \|\mathbf{E}\|_F^2 + \|\mathbf{w}\|^2 \\ \text{s.t.} & (\mathbf{A} + \mathbf{E})\mathbf{x} = \mathbf{b} + \mathbf{w} \end{array}$$

- The KKT conditions are necessary and sufficient for problem $(P_{\mathbf{x}})$.
- Lagrangian: $L(\mathbf{E}, \mathbf{w}, \lambda) = \|\mathbf{E}\|_F^2 + \|\mathbf{w}\|^2 + 2\lambda^\top [(\mathbf{A} + \mathbf{E})\mathbf{x} - \mathbf{b} - \mathbf{w}]$
- By the KKT conditions, (\mathbf{E}, \mathbf{w}) is an optimal solution of $(P_{\mathbf{x}})$ if and only if there exists $\lambda \in \mathbb{R}^m$ such that

$$2\mathbf{E} + 2\lambda\mathbf{x}^\top = \mathbf{0} \quad (\nabla_{\mathbf{E}}L = \mathbf{0}) \quad (19)$$

$$2\mathbf{w} - 2\lambda = \mathbf{0} \quad (\nabla_{\mathbf{w}}L = \mathbf{0}) \quad (20)$$

$$(\mathbf{A} + \mathbf{E})\mathbf{x} = \mathbf{b} + \mathbf{w} \quad (\text{feasibility}) \quad (21)$$

- By (19), (20) and (21), $\mathbf{E} = -\lambda\mathbf{x}^\top$, $\mathbf{w} = \lambda$ and $\lambda = \frac{\mathbf{A}\mathbf{x} - \mathbf{b}}{\|\mathbf{x}\|^2 + 1}$. Plugging this into the objective function, a reduced formulation in the variables \mathbf{x} is obtained.

$$(TLS') \quad \min_{\mathbf{x} \in \mathbb{R}^n} \frac{\|\mathbf{Ax} - \mathbf{b}\|^2}{\|\mathbf{x}\|^2 + 1}$$

Theorem

\mathbf{x} is an optimal solution of (TLS') if and only if $(\mathbf{x}, \mathbf{E}, \mathbf{w})$ is an optimal solution of (TLS) where $\mathbf{E} = -\frac{(\mathbf{Ax} - \mathbf{b})\mathbf{x}^\top}{\|\mathbf{x}\|^2 + 1}$ and $\mathbf{w} = \frac{\mathbf{Ax} - \mathbf{b}}{\|\mathbf{x}\|^2 + 1}$

- Still a nonconvex problem.
- Resembles the problem of minimizing the Rayleigh quotient.

Solving the Fractional Quadratic Formulation

Under a rather mild condition, the optimal solution of (TLS') can be derived via a [homogenization argument](#).

- (TLS') is the same as

$$\min_{\mathbf{x} \in \mathbb{R}^n, t \in \mathbb{R}} \left\{ \frac{\|\mathbf{Ax} - t\mathbf{b}\|^2}{\|\mathbf{x}\|^2 + t^2} : t = 1 \right\}$$

- The same as (denoting $\mathbf{y} = \begin{pmatrix} \mathbf{x} \\ t \end{pmatrix}$):

$$f^* = \min_{\mathbf{y} \in \mathbb{R}^{n+1}} \left\{ \frac{\mathbf{y}^\top \mathbf{B} \mathbf{y}}{\|\mathbf{y}\|^2} : y_{n+1} = 1 \right\} \quad (22)$$

where

$$\mathbf{B} = \begin{pmatrix} \mathbf{A}^\top \mathbf{A} & -\mathbf{A}^\top \mathbf{b} \\ -\mathbf{b}^\top \mathbf{A} & \|\mathbf{b}\|^2 \end{pmatrix}$$

Solving the Fractional Quadratic Formulation Contd.

We will consider the following relaxed version:

$$g^* = \min_{\mathbf{y} \in \mathbb{R}^{n+1}} \left\{ \frac{\mathbf{y}^\top \mathbf{B} \mathbf{y}}{\|\mathbf{y}\|^2} : \mathbf{y} \neq \mathbf{0} \right\} \quad (23)$$

Lemma

Let \mathbf{y}^* be an optimal solution of (23) and assume that $y_{n+1}^* \neq 0$. Then $\tilde{\mathbf{y}} = \frac{1}{y_{n+1}^*} \mathbf{y}^*$ is an optimal solution of (22).

Proof.

- $f^* \geq g^*$.
- $\tilde{\mathbf{y}}$ is feasible for (22) and we have

$$f^* \leq \frac{\tilde{\mathbf{y}}^\top \mathbf{B} \tilde{\mathbf{y}}}{\|\tilde{\mathbf{y}}\|^2} = \frac{\frac{1}{(y_{n+1}^*)^2} (\mathbf{y}^*)^\top \mathbf{B} \mathbf{y}^*}{\frac{1}{(y_{n+1}^*)^2} \|\mathbf{y}^*\|^2} = \frac{(\mathbf{y}^*)^\top \mathbf{B} \mathbf{y}^*}{\|\mathbf{y}^*\|^2} = g^*$$

- Therefore, $\tilde{\mathbf{y}}$ is an optimal solution of both (22) and (23).



Theorem

Assume that the following condition holds:

$$\lambda_{\min}(\mathbf{B}) < \lambda_{\min}(\mathbf{A}^\top \mathbf{A}) \quad (24)$$

where

$$\mathbf{B} = \begin{pmatrix} \mathbf{A}^\top \mathbf{A} & -\mathbf{A}^\top \mathbf{b} \\ -\mathbf{b}^\top \mathbf{A} & \|\mathbf{b}\|^2 \end{pmatrix}$$

Then the optimal solution of problem (TLS') is given by $\frac{1}{y_{n+1}^} \mathbf{v}$, where*

$\mathbf{y}^ = \begin{pmatrix} \mathbf{v} \\ y_{n+1}^* \end{pmatrix}$ is an eigenvector corresponding to the minimum eigenvalue of \mathbf{B} .*

Proof.

- All we need to prove is that under condition (24), an optimal solution \mathbf{y}^* of (23) must satisfy $y_{n+1}^* \neq 0$.
- Assume on the contrary that $y_{n+1}^* = 0$. Then

$$\lambda_{\min}(\mathbf{B}) = \frac{(\mathbf{y}^*)^\top \mathbf{B} \mathbf{y}^*}{\|\mathbf{y}^*\|^2} = \frac{\mathbf{v}^\top \mathbf{A}^\top \mathbf{A} \mathbf{v}}{\|\mathbf{v}\|^2} \geq \lambda_{\min}(\mathbf{A}^\top \mathbf{A})$$

which is a contradiction to (24).

