


# Lecture 6: Convex Sets

Shi Pu

School of Data Science (SDS)  
The Chinese University of Hong Kong, Shenzhen



- 1 Definition and Examples
- 2 Algebraic Operations with Convex Sets
- 3 The Convex Hull
- 4 Convex Cones
- 5 Topological Properties of Convex Sets


- 1 Definition and Examples
- 2 Algebraic Operations with Convex Sets
- 3 The Convex Hull
- 4 Convex Cones
- 5 Topological Properties of Convex Sets

## Definition


A set  $C \subseteq \mathbb{R}^n$  is called **convex** if for any  $\mathbf{x}, \mathbf{y} \in C$  and  $\lambda \in [0, 1]$ , the point  $\lambda\mathbf{x} + (1 - \lambda)\mathbf{y}$  belongs to  $C$ .

- The above definition is equivalent to saying that for any  $\mathbf{x}, \mathbf{y} \in C$ , the line segment  $[\mathbf{x}, \mathbf{y}]$  is also in  $C$ .

convex sets



nonconvex sets



- Lines: a line in  $\mathbb{R}^n$  is a set of the form

$$L = \{\mathbf{z} + t\mathbf{d} : t \in \mathbb{R}\},$$

where  $\mathbf{z}, \mathbf{d} \in \mathbb{R}^n$  and  $\mathbf{d} \neq \mathbf{0}$ .

- $[\mathbf{x}, \mathbf{y}]$ ,  $(\mathbf{x}, \mathbf{y})$  for  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  ( $\mathbf{x} \neq \mathbf{y}$ ) .
- $\emptyset, \mathbb{R}^n$ .
- A **hyperplane** is a set of the form

$$H = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^\top \mathbf{x} = b\} \quad (\mathbf{a} \in \mathbb{R}^n \setminus \{\mathbf{0}\}, b \in \mathbb{R})$$

The associated **half-space** is the set

$$H^- = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}^\top \mathbf{x} \leq b\}$$

Both hyperplanes and half-spaces are convex sets.

## Lemma

Let  $\mathbf{c} \in \mathbb{R}^n$  and  $r > 0$ . Then the open ball

$$B(\mathbf{c}, r) = \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x} - \mathbf{c}\| < r\}$$

and the closed ball




$$B[\mathbf{c}, r] = \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x} - \mathbf{c}\| \leq r\}$$

are convex.

Note that the norm is an arbitrary norm defined over  $\mathbb{R}^n$ .

**Proof.** In class

# $l_1$ , $l_2$ and $l_\infty$ Balls



An **ellipsoid** is a set of the form

$$E = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x}^\top \mathbf{Q} \mathbf{x} + 2\mathbf{b}^\top \mathbf{x} + c \leq 0\},$$

where  $\mathbf{Q} \in \mathbb{R}^{n \times n}$  is positive semidefinite,  $\mathbf{b} \in \mathbb{R}^n$  and  $c \in \mathbb{R}$ .

**Lemma**

*$E$  is convex.*



## Proof.

- Write  $E$  as  $E = \{\mathbf{x} \in \mathbb{R}^n : f(\mathbf{x}) \leq 0\}$  where  $f(\mathbf{x}) \equiv \mathbf{x}^\top \mathbf{Q} \mathbf{x} + 2\mathbf{b}^\top \mathbf{x} + c$ .
- Take  $\mathbf{x}, \mathbf{y} \in E$  and  $\lambda \in [0, 1]$ . Then  $f(\mathbf{x}) \leq 0, f(\mathbf{y}) \leq 0$ .
- The vector  $\mathbf{z} = \lambda \mathbf{x} + (1 - \lambda) \mathbf{y}$  satisfies
$$\mathbf{z}^\top \mathbf{Q} \mathbf{z} = \lambda^2 \mathbf{x}^\top \mathbf{Q} \mathbf{x} + (1 - \lambda)^2 \mathbf{y}^\top \mathbf{Q} \mathbf{y} + 2\lambda(1 - \lambda) \mathbf{x}^\top \mathbf{Q} \mathbf{y}.$$
- $\mathbf{x}^\top \mathbf{Q} \mathbf{y} \leq \|\mathbf{Q}^{1/2} \mathbf{x}\| \cdot \|\mathbf{Q}^{1/2} \mathbf{y}\| = \sqrt{\mathbf{x}^\top \mathbf{Q} \mathbf{x}} \sqrt{\mathbf{y}^\top \mathbf{Q} \mathbf{y}} \leq \frac{1}{2}(\mathbf{x}^\top \mathbf{Q} \mathbf{x} + \mathbf{y}^\top \mathbf{Q} \mathbf{y})$
- $\mathbf{z}^\top \mathbf{Q} \mathbf{z} \leq \lambda \mathbf{x}^\top \mathbf{Q} \mathbf{x} + (1 - \lambda) \mathbf{y}^\top \mathbf{Q} \mathbf{y}$
- $$\begin{aligned} f(\mathbf{z}) &= \mathbf{z}^\top \mathbf{Q} \mathbf{z} + 2\mathbf{b}^\top \mathbf{z} + c \\ &\leq \lambda \mathbf{x}^\top \mathbf{Q} \mathbf{x} + (1 - \lambda) \mathbf{y}^\top \mathbf{Q} \mathbf{y} + 2\lambda \mathbf{b}^\top \mathbf{x} + 2(1 - \lambda) \mathbf{b}^\top \mathbf{y} + \lambda c + (1 - \lambda)c \\ &= \lambda f(\mathbf{x}) + (1 - \lambda) f(\mathbf{y}) \leq 0. \end{aligned}$$



- 1 Definition and Examples
- 2 Algebraic Operations with Convex Sets**
- 3 The Convex Hull
- 4 Convex Cones
- 5 Topological Properties of Convex Sets

## Lemma

*Let  $C_i \subseteq \mathbb{R}^n$  be a convex set for any  $i \in I$  where  $I$  is an index set (possibly infinite). Then the set  $\bigcap_{i \in I} C_i$  is convex.*

**Proof.** In class

Example: Consider the set

$$P = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{Ax} \leq \mathbf{b}\}$$

where  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and  $\mathbf{b} \in \mathbb{R}^m$ .  $P$  is called a **convex polyhedron** and it is indeed convex. Why?

## Theorem (preservation under addition, cartesian product, forward and inverse linear mappings)

1. Let  $C_1, C_2, \dots, C_k \subseteq \mathbb{R}^n$  be convex sets and let  $\mu_1, \mu_2, \dots, \mu_k \in \mathbb{R}$ . Then the set  $\mu_1 C_1 + \mu_2 C_2 + \dots + \mu_k C_k$  is convex.
2. Let  $C_i \subseteq \mathbb{R}^{k_i}, i = 1, 2, \dots, m$  be convex sets. Then the cartesian product

$$C_1 \times C_2 \times \dots \times C_m = \{(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m) : \mathbf{x}_i \in C_i, i = 1, 2, \dots, m\}$$

is convex.

3. Let  $M \subseteq \mathbb{R}^n$  be a convex set and let  $\mathbf{A} \in \mathbb{R}^{m \times n}$ . Then the set

$$\mathbf{A}(M) = \{\mathbf{Ax} : \mathbf{x} \in M\}$$

is convex.

4. Let  $D \subseteq \mathbb{R}^m$  be a convex set and let  $\mathbf{A} \in \mathbb{R}^{m \times n}$ . Then the set

$$\mathbf{A}^{-1}(D) = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{Ax} \in D\}$$

is convex.

Given  $m$  points  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m \in \mathbb{R}^n$ , a **convex combination** of these  $m$  points is a vector of the form

$$\lambda_1 \mathbf{x}_1 + \lambda_2 \mathbf{x}_2 + \dots + \lambda_m \mathbf{x}_m,$$

where  $\lambda_1, \lambda_2, \dots, \lambda_m$  are nonnegative numbers satisfying  $\lambda_1 + \lambda_2 + \dots + \lambda_m = 1$ .

- A convex set is defined by the property that any convex combination of two points from the set is also in the set.
- We will now show that a convex combination of **any** number of points from a convex set is in the set.

## Theorem

*Let  $C \subseteq \mathbb{R}^n$  be a convex set and let  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m \in C$ . Then for any  $\boldsymbol{\lambda} \in \Delta_m$ , the relation  $\sum_{i=1}^m \lambda_i \mathbf{x}_i \in C$  holds.*

## Proof.

- Proof by induction on  $m$ . For  $m = 1$  the result is obvious.
- Induction hypothesis: for any  $m$  vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m \in C$  and any  $\boldsymbol{\lambda} \in \Delta_m$ ,  $\sum_{i=1}^m \lambda_i \mathbf{x}_i \in C$ . Will prove the theorem for  $m + 1$  vectors.
- Suppose that  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{m+1} \in C$  and that  $\boldsymbol{\lambda} \in \Delta_{m+1}$ . We will show that  $\mathbf{z} \equiv \sum_{i=1}^{m+1} \lambda_i \mathbf{x}_i \in C$ .
- If  $\lambda_{m+1} = 1$ , then  $\mathbf{z} = \mathbf{x}_{m+1} \in C$  and the result obviously follows.
- If  $\lambda_{m+1} < 1$  then

$$\mathbf{z} = \sum_{i=1}^m \lambda_i \mathbf{x}_i + \lambda_{m+1} \mathbf{x}_{m+1} = (1 - \lambda_{m+1}) \underbrace{\sum_{i=1}^m \frac{\lambda_i}{1 - \lambda_{m+1}} \mathbf{x}_i}_{\mathbf{v}} + \lambda_{m+1} \mathbf{x}_{m+1}$$

- $\mathbf{v} \in C$  and hence  $\mathbf{z} = (1 - \lambda_{m+1})\mathbf{v} + \lambda_{m+1}\mathbf{x}_{m+1} \in C$ .



- 1 Definition and Examples
- 2 Algebraic Operations with Convex Sets
- 3 The Convex Hull**
- 4 Convex Cones
- 5 Topological Properties of Convex Sets



## Definition

Let  $S \subseteq \mathbb{R}^n$ . The **convex hull** of  $S$ , denoted by  $\text{conv}(S)$ , is the set comprising all the convex combinations of vectors from  $S$ :

$$\text{conv}(S) \equiv \left\{ \sum_{i=1}^k \lambda_i \mathbf{x}_i : \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in S, \lambda \in \Delta_k, k \in \mathbb{N} \right\}.$$




Figure: A nonconvex set and its convex hull

The convex hull  $\text{conv}(S)$  is the “smallest” convex set containing  $S$ .

## Lemma

*Let  $S \subseteq \mathbb{R}^n$ . If  $S \subseteq T$  for some convex set  $T$ , then  $\text{conv}(S) \subseteq T$ .*

The convex hull  $\text{conv}(S)$  is the “smallest” convex set containing  $S$ .

## Lemma

Let  $S \subseteq \mathbb{R}^n$ . If  $S \subseteq T$  for some convex set  $T$ , then  $\text{conv}(S) \subseteq T$ .

## Proof.

- Suppose that indeed  $S \subseteq T$  for some convex set  $T$ .
- To prove that  $\text{conv}(S) \subseteq T$ , take  $\mathbf{z} \in \text{conv}(S)$ .
- There exist  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in S \subseteq T$  (where  $k$  is a positive integer), and  $\boldsymbol{\lambda} \in \Delta_k$  such that  $\mathbf{z} = \sum_{i=1}^k \lambda_i \mathbf{x}_i$ .
- Since  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in T$ , it follows that  $\mathbf{z} \in T$ , showing the desired result.



Any element in the convex hull of a given set  $S \subseteq \mathbb{R}^n$  can be represented as a convex combination of no more than  $n + 1$  vectors from  $S$ .

## Theorem

Let  $S \subseteq \mathbb{R}^n$  and let  $\mathbf{x} \in \text{conv}(S)$ . Then there exist  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1} \in S$  such that  $\mathbf{x} \in \text{conv}(\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1}\})$ , that is, there exist  $\lambda \in \Delta_{n+1}$  such that

$$\mathbf{x} = \sum_{i=1}^{n+1} \lambda_i \mathbf{x}_i.$$

## Proof.

- Let  $\mathbf{x} \in \text{conv}(S)$ . Then  $\exists \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in S$  and  $\boldsymbol{\lambda} \in \Delta_k$  s.t.

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i.$$

- We can assume that  $\lambda_i > 0$  for all  $i = 1, 2, \dots, k$ .
- If  $k \leq n + 1$ , the result is proven.
- Otherwise, if  $k \geq n + 2$ , then the vectors  $\mathbf{x}_2 - \mathbf{x}_1, \mathbf{x}_3 - \mathbf{x}_1, \dots, \mathbf{x}_k - \mathbf{x}_1$ , being more than  $n$  vectors in  $\mathbb{R}^n$ , are necessarily linearly dependent  $\Rightarrow \exists \mu_2, \mu_3, \dots, \mu_k$  not all zeros s.t.

$$\sum_{i=2}^k \mu_i (\mathbf{x}_i - \mathbf{x}_1) = \mathbf{0}.$$

## Proof Contd.

- Defining  $\mu_1 = -\sum_{i=2}^k \mu_i$ , we obtain that  $\sum_{i=1}^k \mu_i \mathbf{x}_i = \mathbf{0}$ .
- Not all of the coefficients  $\mu_1, \mu_2, \dots, \mu_k$  are zeros and  $\sum_{i=1}^k \mu_i = 0$ .
- There exists an index  $i$  for which  $\mu_i < 0$ . Let  $\alpha \in \mathbb{R}_+$ . Then

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i = \sum_{i=1}^k \lambda_i \mathbf{x}_i + \alpha \sum_{i=1}^k \mu_i \mathbf{x}_i = \sum_{i=1}^k (\lambda_i + \alpha \mu_i) \mathbf{x}_i. \quad (1)$$

- We have  $\sum_{i=1}^k (\lambda_i + \alpha \mu_i) = 1$ , so (1) is a convex combination representation iff

$$\lambda_i + \alpha \mu_i \geq 0 \text{ for all } i = 1, \dots, k. \quad (2)$$

- Since  $\lambda_i > 0$  for all  $i$ , it follows that (2) is satisfied for all  $\alpha \in [0, \epsilon]$  where  $\epsilon = \min_{i: \mu_i < 0} \left\{ -\frac{\lambda_i}{\mu_i} \right\}$ .

## Proof Contd.

- If we substitute  $\alpha = \epsilon$ , then (2) still holds, but  $\lambda_j + \epsilon\mu_j = 0$  for  $j \in \operatorname{argmin}_{i:\mu_i < 0} \{-\frac{\lambda_i}{\mu_i}\}$ .
- This means that we found a representation of  $\mathbf{x}$  as a convex combination of  $k - 1$  (or less) vectors.
- This process can be carried on until a representation of  $\mathbf{x}$  as a convex combination of no more than  $n + 1$  vectors is derived.



## Example

For  $n = 2$ , consider the four vectors

$$\mathbf{x}_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \mathbf{x}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \mathbf{x}_3 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \mathbf{x}_4 = \begin{pmatrix} 2 \\ 2 \end{pmatrix},$$

and let  $\mathbf{x} \in \text{conv}(\{\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4\})$  be given by

$$\mathbf{x} = \frac{1}{8}\mathbf{x}_1 + \frac{1}{4}\mathbf{x}_2 + \frac{1}{2}\mathbf{x}_3 + \frac{1}{8}\mathbf{x}_4 = \begin{pmatrix} \frac{13}{8} \\ \frac{11}{8} \end{pmatrix}.$$

Find a representation of  $\mathbf{x}$  as a convex combination of no more than 3 vectors.

In class



- 1 Definition and Examples
- 2 Algebraic Operations with Convex Sets
- 3 The Convex Hull
- 4 Convex Cones**
- 5 Topological Properties of Convex Sets

- A set  $S$  is called a **cone** if it satisfies the following property: for any  $\mathbf{x} \in S$  and  $\lambda \geq 0$ , the inclusion  $\lambda \mathbf{x} \in S$  is satisfied.
- The following lemma shows that there is a very simple and elegant characterization of convex cones.

## Lemma

*A set  $S$  is a convex cone if and only if the following properties hold:*

1.  $\mathbf{x}, \mathbf{y} \in S \Rightarrow \mathbf{x} + \mathbf{y} \in S$ .
2.  $\mathbf{x} \in S, \lambda \geq 0 \Rightarrow \lambda \mathbf{x} \in S$ .

Simple exercise

# Examples of Convex Cones

- The convex polyhedron

$$C = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{Ax} \leq \mathbf{0}\}$$


where  $\mathbf{A} \in \mathbb{R}^{m \times n}$ .

- The **Lorenz cone**, or **ice cream cone** is given by

$$L^n = \left\{ \begin{pmatrix} \mathbf{x} \\ t \end{pmatrix} \in \mathbb{R}^{n+1} : \|\mathbf{x}\| \leq t, \mathbf{x} \in \mathbb{R}^n, t \in \mathbb{R} \right\}.$$

- **Nonnegative polynomials**: set consisting of all possible coefficients of polynomials of degree  $n - 1$  which are nonnegative over  $\mathbb{R}$ :

$$K^n = \{\mathbf{x} \in \mathbb{R}^n : x_1 t^{n-1} + x_2 t^{n-2} + \dots + x_{n-1} t + x_n \geq 0, \forall t \in \mathbb{R}\}.$$



# The Conic Hull

## Definition

Given  $m$  points  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m \in \mathbb{R}^n$ , a **conic combination** of these  $m$  points is a vector of the form  $\lambda_1 \mathbf{x}_1 + \lambda_2 \mathbf{x}_2 + \dots + \lambda_m \mathbf{x}_m$ , where  $\boldsymbol{\lambda} \in \mathbb{R}_+^m$ .

The definition of the **conic hull** is now quite natural.

## Definition

Let  $S \subseteq \mathbb{R}^n$ . Then the **conic hull** of  $S$ , denoted by  $\text{cone}(S)$ , is the set comprising all the conic combinations of vectors from  $S$ :

$$\text{cone}(S) \equiv \left\{ \sum_{i=1}^k \lambda_i \mathbf{x}_i : \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in S, \boldsymbol{\lambda} \in \mathbb{R}_+^k \right\}.$$

The conic hull of a set  $S$  is the smallest convex cone containing  $S$ .

## Lemma

Let  $S \subseteq \mathbb{R}^n$ . If  $S \subseteq T$  for some convex cone  $T$ , then  $\text{cone}(S) \subseteq T$ .

The following result is similar to Carathéodory Theorem.

## Theorem (Conic Representation Theorem)

Let  $S \subseteq \mathbb{R}^n$  and let  $\mathbf{x} \in \text{cone}(S)$ . Then there exist  $k$  *linearly independent* vectors  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in S$  such that  $\mathbf{x} \in \text{cone}(\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\})$ , that is, there exist  $\lambda \in \mathbb{R}_+^k$  such that

$$\mathbf{x} = \sum_{i=1}^k \lambda_i \mathbf{x}_i.$$

In particular,  $k \leq n$ .

Proof very similar to the proof of Carathéodory theorem; see page 107 of the textbook.

# Basic Feasible Solutions

- Consider the convex polyhedron.

$$P = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}, \quad (\mathbf{A} \in \mathbb{R}^{m \times n}, \mathbf{b} \in \mathbb{R}^m)$$

- The rows of  $\mathbf{A}$  are assumed to be linearly independent.
- The above is a standard formulation of the constraints of a linear programming problem.

## Definition

$\bar{\mathbf{x}}$  is a **basic feasible solution** (bfs) of  $P$  if the columns of  $\mathbf{A}$  corresponding to the indices of the positive values of  $\bar{\mathbf{x}}$  are linearly independent.

Example. Consider the linear system:

$$x_1 + x_2 + x_3 = 6$$

$$x_2 + x_4 = 3$$

$$x_1, x_2, x_3, x_4 \geq 0.$$

Find all the basic feasible solutions. **In class**

## Theorem

*Let  $P = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ , where  $\mathbf{A} \in \mathbb{R}^{m \times n}, \mathbf{b} \in \mathbb{R}^m$ . If  $P \neq \emptyset$ , then it contains at least one bfs.*

## Proof.

- $P \neq \emptyset \Rightarrow \mathbf{b} \in \text{cone}(\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n\})$  where  $\mathbf{a}_i$  denotes the  $i$ -th column of  $\mathbf{A}$ .
- By the conic representation theorem, there exist indices  $i_1 < i_2 < \dots < i_k$  and  $k$  numbers  $y_{i_1}, y_{i_2}, \dots, y_{i_k} \geq 0$  such that  $\mathbf{b} = \sum_{j=1}^k y_{i_j} \mathbf{a}_{i_j}$  and  $\mathbf{a}_{i_1}, \mathbf{a}_{i_2}, \dots, \mathbf{a}_{i_k}$  are linearly independent.
- Denote  $\bar{\mathbf{x}} = \sum_{j=1}^k y_{i_j} \mathbf{e}_{i_j}$ . Then obviously  $\bar{\mathbf{x}} \geq \mathbf{0}$  and in addition

$$\mathbf{A}\bar{\mathbf{x}} = \sum_{j=1}^k y_{i_j} \mathbf{A}\mathbf{e}_{i_j} = \sum_{j=1}^k y_{i_j} \mathbf{a}_{i_j} = \mathbf{b}.$$

- Therefore,  $\bar{\mathbf{x}}$  is contained in  $P$  and the columns of  $\mathbf{A}$  corresponding to the indices of the positive components of  $\bar{\mathbf{x}}$  are linearly independent, meaning that  $P$  contains a bfs.





- 1 Definition and Examples
- 2 Algebraic Operations with Convex Sets
- 3 The Convex Hull
- 4 Convex Cones
- 5 Topological Properties of Convex Sets**

## Theorem

*Let  $C \subseteq \mathbb{R}^n$  be a convex set. Then  $\text{cl}(C)$  is a convex set.*

## Theorem

Let  $C \subseteq \mathbb{R}^n$  be a convex set. Then  $\text{cl}(C)$  is a convex set.

## Proof.

- Let  $\mathbf{x}, \mathbf{y} \in \text{cl}(C)$  and let  $\lambda \in [0, 1]$ .
- There exist sequences  $\{\mathbf{x}_k\}_{k \geq 0} \subseteq C$  and  $\{\mathbf{y}_k\}_{k \geq 0} \subseteq C$  for which  $\mathbf{x}_k \rightarrow \mathbf{x}$  and  $\mathbf{y}_k \rightarrow \mathbf{y}$  as  $k \rightarrow \infty$ .
- (\*)  $\lambda \mathbf{x}_k + (1 - \lambda) \mathbf{y}_k \in C$  for any  $k \geq 0$ .
- (\*\*)  $\lambda \mathbf{x}_k + (1 - \lambda) \mathbf{y}_k \rightarrow \lambda \mathbf{x} + (1 - \lambda) \mathbf{y}$ .
- (\*) + (\*\*)  $\Rightarrow \lambda \mathbf{x} + (1 - \lambda) \mathbf{y} \in \text{cl}(C)$ .



## Theorem

*Let  $C$  be a convex set and assume that  $\text{int}(C) \neq \emptyset$ . Suppose that  $\mathbf{x} \in \text{int}(C)$  and  $\mathbf{y} \in \text{cl}(C)$ . Then  $(1 - \lambda)\mathbf{x} + \lambda\mathbf{y} \in \text{int}(C)$  for any  $\lambda \in [0, 1)$ .*

# The Line Segment Principle

## Proof.

- There exists  $\epsilon > 0$  such that  $B(\mathbf{x}, \epsilon) \subseteq C$ .
- Let  $\mathbf{z} = (1 - \lambda)\mathbf{x} + \lambda\mathbf{y}$ . We will show that  $B(\mathbf{z}, (1 - \lambda)\epsilon) \subseteq C$ .
- Let  $\mathbf{w} \in B(\mathbf{z}, (1 - \lambda)\epsilon)$ . Since  $\mathbf{y} \in \text{cl}(C)$ ,  $\exists \mathbf{w}_1 \in C$  s.t.

$$\|\mathbf{w}_1 - \mathbf{y}\| < \frac{(1 - \lambda)\epsilon - \|\mathbf{w} - \mathbf{z}\|}{\lambda}. \quad (3)$$

- Set  $\mathbf{w}_2 = \frac{1}{1 - \lambda}(\mathbf{w} - \lambda\mathbf{w}_1)$ . Then

$$\begin{aligned} \|\mathbf{w}_2 - \mathbf{x}\| &= \left\| \frac{\mathbf{w} - \lambda\mathbf{w}_1}{1 - \lambda} - \mathbf{x} \right\| = \frac{1}{1 - \lambda} \|(\mathbf{w} - \mathbf{z}) + \lambda(\mathbf{y} - \mathbf{w}_1)\| \\ &\leq \frac{1}{1 - \lambda} (\|\mathbf{w} - \mathbf{z}\| + \lambda\|\mathbf{w}_1 - \mathbf{y}\|) < \epsilon \quad (\text{using (3)}), \end{aligned}$$

- Hence, since  $B(\mathbf{x}, \epsilon) \subseteq C$ , it follows that  $\mathbf{w}_2 \in C$ . Finally, since  $\mathbf{w} = \lambda\mathbf{w}_1 + (1 - \lambda)\mathbf{w}_2$  with  $\mathbf{w}_1, \mathbf{w}_2 \in C$ , we have that  $\mathbf{w} \in C$ .



## Theorem

*Let  $C \subseteq \mathbb{R}^n$  be a convex set. Then  $\text{int}(C)$  is convex.*

## Proof.

- If  $\text{int}(C) = \emptyset$ , then the theorem is obviously true.
- Otherwise, let  $\mathbf{x}_1, \mathbf{x}_2 \in \text{int}(C)$ , and let  $\lambda \in (0, 1)$ .
- By the LSP,  $\lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2 \in \text{int}(C)$ , establishing the convexity of  $\text{int}(C)$ .



## Lemma

*Let  $C$  be a convex set with a nonempty interior. Then*

1.  $\text{cl}(\text{int}(C)) = \text{cl}(C)$ .
2.  $\text{int}(\text{cl}(C)) = \text{int}(C)$ .

## Lemma

*Let  $C$  be a convex set with a nonempty interior. Then*

1.  $\text{cl}(\text{int}(C)) = \text{cl}(C)$ .
2.  $\text{int}(\text{cl}(C)) = \text{int}(C)$ .

## Proof of 1.

- Obviously,  $\text{cl}(\text{int}(C)) \subseteq \text{cl}(C)$  holds.
- To prove that opposite, let  $\mathbf{x} \in \text{cl}(C)$ ,  $\mathbf{y} \in \text{int}(C)$ .
- Then  $\mathbf{x}_k = \frac{1}{k}\mathbf{y} + (1 - \frac{1}{k})\mathbf{x} \in \text{int}(C)$  for any  $k \geq 1$ .
- Since  $\mathbf{x}$  is the limit (as  $k \rightarrow \infty$ ) of the sequence  $\{\mathbf{x}_k\}_{k \geq 1} \subseteq \text{int}(C)$ , it follows that  $\mathbf{x} \in \text{cl}(\text{int}(C))$ .



For the proof of 2, see pages 109, 110 of the textbook for the proof of Lemma 6.30(b).



- In general, the convex hull of a closed set is not necessarily a closed set.
- Example:  $S = \{(0, 0)^\top\} \cup \{(x, y)^\top : xy \geq 1, x \geq 0, y \geq 0\}$ .

## Theorem

*Let  $S \subseteq \mathbb{R}^n$  be a compact set. Then  $\text{conv}(S)$  is compact.*

# Compactness of the Convex Hull of a Compact Set

## Proof.

- $\exists M > 0$  such that  $\|\mathbf{x}\| \leq M$  for any  $\mathbf{x} \in S$ .
- Let  $\mathbf{y} \in \text{conv}(S)$ . Then there exist  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1} \in S$  and  $\boldsymbol{\lambda} \in \Delta_{n+1}$  for which  $\mathbf{y} = \sum_{i=1}^{n+1} \lambda_i \mathbf{x}_i$  and therefore

$$\|\mathbf{y}\| = \left\| \sum_{i=1}^{n+1} \lambda_i \mathbf{x}_i \right\| \leq \sum_{i=1}^{n+1} \lambda_i \|\mathbf{x}_i\| \leq M \sum_{i=1}^{n+1} \lambda_i = M,$$

establishing the boundedness of  $\text{conv}(S)$ .

- To prove the closedness of  $\text{conv}(S)$ , let  $\{\mathbf{y}_k\}_{k \geq 1} \subseteq \text{conv}(S)$  be a sequence converging to  $\mathbf{y} \in \mathbb{R}^n$ .
- There exist  $\mathbf{x}_1^k, \mathbf{x}_2^k, \dots, \mathbf{x}_{n+1}^k \in S$  and  $\boldsymbol{\lambda}^k \in \Delta_{n+1}$  such that

$$\mathbf{y}_k = \sum_{i=1}^{n+1} \lambda_i^k \mathbf{x}_i^k. \quad (4)$$

## Proof Contd.

- By the compactness of  $S$  and  $\Delta_{n+1}$ , it follows that  $\{(\lambda^k, \mathbf{x}_1^k, \mathbf{x}_2^k, \dots, \mathbf{x}_{n+1}^k)\}_{k \geq 1}$  has a convergent subsequence  $\{(\lambda^{k_j}, \mathbf{x}_1^{k_j}, \mathbf{x}_2^{k_j}, \dots, \mathbf{x}_{n+1}^{k_j})\}_{j \geq 1}$  whose limit will be denoted by  $(\lambda, \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1})$

with  $\lambda \in \Delta_{n+1}$ ,  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{n+1} \in S$

- Taking the limit  $j \rightarrow \infty$  in

$$\mathbf{y}_{k_j} = \sum_{i=1}^{n+1} \lambda_i^{k_j} \mathbf{x}_i^{k_j},$$


we obtain that  $\mathbf{y} = \sum_{i=1}^{n+1} \lambda_i \mathbf{x}_i \in \text{conv}(S)$  as required.



## Definition

Let  $S \subseteq \mathbb{R}^n$  be a convex set. A point  $\mathbf{x} \in S$  is called an **extreme point** of  $S$  if there do not exist  $\mathbf{x}_1, \mathbf{x}_2 \in S$  ( $\mathbf{x}_1 \neq \mathbf{x}_2$ ) and  $\lambda \in (0, 1)$ , such that  $\mathbf{x} = \lambda \mathbf{x}_1 + (1 - \lambda) \mathbf{x}_2$ .

- The set of extreme point is denoted by  $\text{ext}(S)$ .
- For example, the set of extreme points of a convex polytope (bounded polyhedron) consists of all its vertices.



## Theorem

*Let  $P = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$ , where  $\mathbf{A} \in \mathbb{R}^{m \times n}$  has linearly independent rows and  $\mathbf{b} \in \mathbb{R}^m$ . Then  $\bar{\mathbf{x}}$  is a basic feasible solution of  $P$  if and only if it is an extreme point of  $P$ .*

Theorem 6.34 in the book.

## Theorem

*Let  $S \subseteq \mathbb{R}^n$  be a compact convex set. Then*

$$S = \text{conv}(\text{ext}(S)).$$