1

NCERT Discrete - 10.5.3.20

EE23BTECH1205 - Avani Chouhan*

Question: 10.5.3.20

In a potato race, a bucket is placed at the starting point, which is 5m from the first potato, and the other potatoes are placed 3m apart in a straight line. There are ten potatoes in the line.

A competitor starts from the bucket, picks up the nearest potato, runs back with it, drops it in the bucket, runs back to pick up the next potato, runs to the bucket to drop it in, and she continues in the same way until all the potatoes are in the bucket. What is the total distance the competitor has to run?

Solution: The distance covered forms an AP with given parameters

Parameter	Value	Description
x(0)	10	First term
d	6	Common difference
x(n)	6n + 10	General term
n	10	Number of terms

TABLE 0
INPUT PARAMETERS

using the residue theorem, expressed as R_i , which represents the residue of the Z-transform at z = 1 for the expression Y(z).

The sum of the terms of the sequence is computed

$$R_i = R_1 + R_2 \tag{7}$$

 R_1 and R_2 are residues calculated at the poles of the Z-transform.

$$R_1 = \frac{1}{(2-1)!} \frac{d(10z^{10})}{dz} |_{z=1}$$
 (8)

$$= 10 \times 10 \tag{9}$$

$$= 100$$
 (10)

$$R_2 = \frac{1}{(3-1)!} \cdot \frac{d^2(6z^{10})}{dz^2} \Big|_{z=1}$$
 (11)

$$=\frac{6}{2}(10)(9)\tag{12}$$

$$=270\tag{13}$$

The sum of terms is given by R_i :

$$100 + 270 = 370 \tag{14}$$

From equation (??): Applying Z transform:

$$\implies X(z) = \frac{10}{1 - z^{-1}} + \frac{6z^{-1}}{(1 - z^{-1})^2} \quad |z| > |1| \quad (1)$$

For AP, the sum of first n+1 terms can be written as :

$$y(n) = x(n) * u(n) \tag{2}$$

$$Y(z) = X(z)U(z) \tag{3}$$

$$= \frac{10}{(1-z^{-1})^2} + \frac{6z^{-1}}{(1-z^{-1})^3} \quad |z| > |1| \quad (4)$$

Using contour integration to find inverse Z transform:

$$y(n) = \frac{1}{2\pi j} \oint_C Y(z) z^{n-1} dz \tag{5}$$

$$= \frac{1}{2\pi j} \oint_C \left[\frac{10}{(1-z^{-1})^2} + \frac{6z^{-1}}{(1-z^{-1})^3} \right] z^8 dz \quad (6)$$




Fig. 0. Simulation vs Analysis of y(n)