

1°. (2 points). Let $\varphi : \mathbb{R}[x, n] \rightarrow \mathbb{R}[x, n]$ defined as $\varphi(p(x)) = x^2 \cdot (p(x-1) - p(x+1))'$.

(a) Find $\text{Spec}(\varphi)$ (b) For each $\lambda \in \text{Spec}(\varphi)$ find its algebraic and geometric multiplicity.

The matrix is upper-triangular in the standard basis (explain why). Avoid doing unnecessary calculations.

a) $1 \rightarrow 0 = v_1$

$x \rightarrow x^2((x-1) - (x+1))' = 0 = v_2$

$x^2 \rightarrow x^2((x-1)^2 - (x+1)^2)' = -4x^2 = v_3$

$x^3 \rightarrow x^2((x-1)^3 - (x+1)^3)' = -12x^3 = v_4$

$x^4 \rightarrow x^2((x-1)^4 - (x+1)^4)' = -24x^4 - 8x^2 = v_5$

\vdots

$$\begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -4 & 0 & -8 \\ 0 & 0 & 0 & -12 & 0 \\ 0 & 0 & 0 & 0 & -24 \end{bmatrix}$$

The matrix is upper triangular, as operator cannot increase the maximum power of the input. Therefore, spec consists of numbers from the diagonal.

$\text{Spec} = \{-2(n-1)(n-2), n = \dim(\mathbb{R}[x, n])\}$

b) a. m. of 0 is 2.

g. m. of 0 is 2.

a. m. and g. m. of $-2(n-1)(n-2)$ $n \geq 3$ is 1, as it is only encountered once and the rank of $(A - (-2(n-1)(n-2))I) = n - (n-1) = 1$

2°. (1 point). Let $A = [a_1, \dots, a_n]$ (at least one number is nonzero) and let $B = A^T A$. Consider $\varphi: \mathbb{R}^n \rightarrow \mathbb{R}^n$, $\varphi(v) = Bv$.

(a) Find $\text{Spec}(\varphi)$ (b) For each $\lambda \in \text{Spec}(\varphi)$ find its algebraic and geometric multiplicity.

What is the image of φ ? What is the dimension of the kernel? The smart solution takes about two lines.

$$\varphi: \mathbb{R}^n \rightarrow \mathbb{R} \rightarrow \mathbb{R}^n, \text{ hence the image is } \left\langle \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \right\rangle$$

$$\dim \ker \varphi = n-1$$

$$\begin{aligned} \text{a)} \quad A^T A \cdot \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} &= A^T [a_1 x_1 + \dots + a_n x_n] = \begin{bmatrix} a_1 \sum_{k=1}^n a_k x_k \\ \vdots \\ a_n \sum_{k=1}^n a_k x_k \end{bmatrix} = \\ &= \sum_{k=1}^n a_k x_k \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix} \end{aligned}$$

$$\text{Spec } \varphi = \left\{ 0, \sum_{k=1}^n a_k^2 \right\}$$

$$\text{b)} \quad \text{a.m.}(0) = \dim \ker \varphi = n-1$$

$$\text{g.m.}(0) = n - \text{rk } \varphi = n - (n-1) = 1$$

$$\text{a.m.} \sum_{k=1}^n a_k^2 = 1$$

$$\text{g.m.} \sum_{k=1}^n a_k^2 = 1$$

3°. (0.4 points per item). For each matrix A find its characteristic $\chi_A(x)$ and minimal $m_A(x)$ polynomials.

(a) $\begin{bmatrix} 2 & 1 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ (b) $\begin{bmatrix} 2 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ (c) $\begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 1 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ (d) $\begin{bmatrix} 2 & 1 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 1 \\ 0 & 0 & 0 & 3 \end{bmatrix}$ (e) $\begin{bmatrix} -3 & 2 & 4 & 0 \\ -7 & 5 & 5 & 0 \\ -3 & 1 & 5 & 0 \\ -3 & 1 & 2 & 3 \end{bmatrix}$

Writing down $m_A(x)$ without proof of minimality will not be accepted.


Recall that $\chi_A(x)$ and $m_A(x)$ have the same roots.

$$\begin{aligned} \text{a)} \quad \chi_A(x) &= (x-2)^3(x-3) \\ m_A(x) &= (x-2)^2(x-3) \Rightarrow \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}^2 \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0 \end{aligned}$$

$(x-2)(x-3)$ doesn't work \odot


$$b) X_A(x) = (x-2)^3(x-3)$$

$$m_A(x) = (x-2)^2(x-3) \Rightarrow \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = 0$$

$(x-2)(x-3)$ doesn't work either 


$$c) X_A(x) = (x-2)^2(x-3)^2$$

$$m_A(x) = (x-2)(x-3) \Rightarrow \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}^2 = 0$$

only dropped one power 

$$d) X_A(x) = (x-2)^2(x-3)^2$$

$$m_A(x) = (x-2)^2(x-3)^2 \Rightarrow \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}^2 \cdot \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}^2 = 0$$

It didn't drop a single power 

$$e) \begin{vmatrix} -3-\lambda & 2 & 4 & 0 \\ -7 & 5-\lambda & 5 & 0 \\ -3 & 1 & 5-\lambda & 0 \\ -3 & 1 & 2 & 3-\lambda \end{vmatrix} = (\lambda-3)^2(\lambda-2)^2$$

$$m_A(x) = (x-3)(x-2)^2 \Rightarrow \begin{bmatrix} -6 & 2 & 4 & 0 \\ -7 & 2 & 5 & 0 \\ -3 & 1 & 2 & 0 \\ -3 & 1 & 2 & 0 \end{bmatrix} \cdot \begin{bmatrix} -5 & 2 & 4 & 0 \\ -7 & 3 & 5 & 0 \\ -3 & 1 & 3 & 0 \\ -3 & 1 & 2 & 1 \end{bmatrix}^2 = 0$$

No emotions...

4°. (1 point). For $A = \begin{bmatrix} 4 & -1 & -1 \\ 2 & 1 & -1 \\ 3 & -1 & 0 \end{bmatrix}$ find $a, b, c \in \mathbb{R}$ such that $A^{-1} = aA^2 + bA + cI$.

$$\chi_A(x) = -(x-2)^2(x-1) = -x^3 + 5x^2 - 8x + 4$$

$$-A^3 + 5A^2 - 8A + 4I = 0$$

$$A \cdot \frac{1}{4} (A^2 - 5A + 8 \cdot I) = I$$

$$\frac{1}{4} \begin{bmatrix} 4 & -1 & -1 \\ 2 & 1 & -1 \\ 3 & -1 & 0 \end{bmatrix}^2 - \frac{5}{4} \begin{bmatrix} 4 & -1 & -1 \\ 2 & 1 & -1 \\ 3 & -1 & 0 \end{bmatrix} + \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix} = A^{-1}$$

$$a = \frac{1}{4}, \quad b = -\frac{5}{4}, \quad c = 2$$

5. (1 point). Does there exist a 10×10 matrix such that $A^{100} = 0$ and $A^{99} \neq 0$?

$$\dim = 10 \cdot 10 = 100$$

$$m_\varphi(x) = t^k, \quad k \leq n$$

k is 100 in our case $100 \leq 100$

Hence, it does exist.

6. (1 point). Find an example of a matrix A such that its minimal polynomial is equal to $(x+2)^5$ (the size of A is up to you).

$$\begin{bmatrix} -2 & 1 & 0 & 0 & 0 \\ 0 & -2 & 1 & 0 & 0 \\ 0 & 0 & -2 & 1 & 0 \\ 0 & 0 & 0 & -2 & 1 \\ 0 & 0 & 0 & 0 & -2 \end{bmatrix} = A \Rightarrow m_\varphi(x) = (x+2)^5$$

7. (2 points). Suppose that there are three types of pokemons: blue, red, and green pokemons.

It is known that in one day time:

1. a blue pokemon evolves into one red pokemon;
2. a red pokemon evolves into two green pokemons;
3. a green pokemon evolves into two blue and one red pokemons.

For example: if you start with one blue and one green pokemons, then, in one day you will have two blue and two red pokemons ($B \rightarrow R$ and $G \rightarrow 2B + R$) and in two days you will have two red and four green pokemons ($2B \rightarrow 2R$ and $2R \rightarrow 4G$).

Suppose you start with five blue pokemons, then, how many blue, red, and green pokemons will you have in 60 days?

You will need to find the remainder of x^{60} when divided by a certain polynomial $\chi(x)$ with complex roots. The remainder has real coefficients but you will need to use complex numbers in the intermediate steps.

$$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \xrightarrow{\text{day}} \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad T(\text{day, pokemons}) = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 2 & 0 \end{bmatrix} = A$$

$$\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \xrightarrow{\text{day}} \begin{bmatrix} 0 \\ 0 \\ 2 \end{bmatrix} \quad \chi_{\text{day}}(x) = -x^3 + 2x + 4$$

$$x^{60} = q(x) \cdot (-x^3 + 2x + 4) + r(x)$$

$$\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \xrightarrow{\text{day}} \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} \quad \begin{cases} r(2) = 2^{60} \\ r(-1+i) = \left(\left((-1+i)^2 \right)^2 \right)^{15} = -2^{30} \\ r(-1-i) = (-1-i)^{60} = -2^{30} \end{cases}$$

$$r(x) = 2^{60} \frac{(x+1-i)(x+1+i)}{(2+1-i)(2+1+i)} - 2^{30} \frac{(x-2)(x+1+i)}{(-1+i-2)(-1+i+1+i)} -$$

$$- 2^{30} \frac{(x-2)(x+1-i)}{(-1-i-2)(-1-i+1-i)} =$$

$$= 2^{60} \left(\frac{x^2 + 2x + 2}{10} \right) - 2^{30} \left(\frac{-x^2 - 2x + 8}{20} + \frac{3x^2 - 4x - 4}{20} i \right) + 2^{30} \left(\frac{x^2 + 2x - 8}{20} + \frac{3x^2 - 4x - 4}{20} i \right) =$$

$$= \frac{2^{60} x^2 + 2^{61} x + 2^{61}}{10} - \frac{2^{30} x^2 - 2^{31} x + 2^{33}}{20} + \frac{2^{30} x^2 + 2^{31} x - 2^{33}}{20} =$$

$$= \frac{2^{60} x^2 + 2^{61} x + 2^{61}}{10} + \frac{2^{31} x^2 + 2^{32} x - 2^{34}}{20} =$$

$$= \frac{2^{61} x^2 + 2^{61} x^2 + 2^{62} x + 2^{62} x + 2^{62} x - 2^{34}}{20} =$$

$$= \frac{(2^{61} + 2^{31})}{20} x^2 + \frac{(2^{62} + 2^{32})}{20} x + \frac{(2^{62} - 2^{34})}{20}$$

$$A^{60} = \frac{(2^{61} + 2^{31})}{20} A^2 + \frac{(2^{62} + 2^{32})}{20} A + \frac{(2^{62} - 2^{34})}{20} I$$

$$A^{60} = \frac{2^{61} + 2^{31}}{20} \begin{bmatrix} 0 & 4 & 0 \\ 0 & 2 & 2 \\ 2 & 0 & 2 \end{bmatrix} + \frac{2^{62} + 2^{32}}{20} \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 2 & 0 \end{bmatrix} + \frac{2^{62} - 2^{34}}{20} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} =$$

$$= \begin{bmatrix} 0 & \frac{2^{61} + 2^{31}}{5} & 0 \\ 0 & \frac{2^{61} + 2^{31}}{10} & \frac{2^{61} + 2^{31}}{10} \\ \frac{2^{61} + 2^{31}}{10} & 0 & \frac{2^{61} + 2^{31}}{10} \end{bmatrix} + \begin{bmatrix} 0 & 0 & \frac{2^{62} + 2^{32}}{10} \\ \frac{2^{62} + 2^{32}}{20} & 0 & \frac{2^{62} + 2^{32}}{20} \\ 0 & \frac{2^{62} + 2^{32}}{10} & 0 \end{bmatrix} + \begin{bmatrix} \frac{2^{62} - 2^{34}}{20} & 0 & 0 \\ 0 & \frac{2^{62} - 2^{34}}{20} & 0 \\ 0 & 0 & \frac{2^{62} - 2^{34}}{20} \end{bmatrix} =$$

$$\Rightarrow \begin{bmatrix} \frac{2^{62} - 2^{34}}{20} & \frac{2^{61} + 2^{31}}{5} & \frac{2^{62} + 2^{32}}{10} \\ \frac{2^{62} + 2^{32}}{20} & \frac{2^{63} + 2^{32} + 2^{34}}{20} & \frac{2^{63} + 2^{33}}{20} \\ \frac{2^{61} + 2^{31}}{10} & \frac{2^{62} + 2^{32}}{10} & \frac{2^{63} + 2^{32} + 2^{34}}{20} \end{bmatrix} \cdot \begin{bmatrix} 5 \\ 0 \\ 0 \end{bmatrix} =$$

$$= \begin{bmatrix} \frac{2^{62} - 2^{34}}{4} \\ \frac{2^{62} + 2^{32}}{4} \\ \frac{2^{61} + 2^{31}}{2} \end{bmatrix}$$

blue

green

red

That's a lot of
fucking pokemons.