

Stickler syndrome

Overview

Stickler syndrome is a genetic disorder that can cause serious vision, hearing and joint problems. Also known as hereditary progressive arthro-ophthalmopathy, Stickler syndrome is usually diagnosed during infancy or childhood.

Children who have Stickler syndrome often have distinctive facial features — prominent eyes, a small nose with a scooped-out facial appearance and a receding chin. They are often born with an opening in the roof of the mouth (cleft palate).

While there is no cure for Stickler syndrome, treatments can help control symptoms and prevent complications. In some cases, surgery may be needed to correct some of the physical abnormalities associated with Stickler syndrome.

Symptoms

The signs and symptoms of Stickler syndrome — and the severity of those signs and symptoms — can vary widely from person to person, even within the same family.

- **Eye problems.** In addition to having severe nearsightedness, children who have Stickler syndrome often experience cataracts, glaucoma and retinal detachments.
- **Hearing difficulties.** The extent of hearing loss varies among people who have Stickler syndrome. It usually affects the ability to hear high frequencies.
- Bone and joint abnormalities. Children who have Stickler syndrome often have overly flexible joints and are
 more likely to develop abnormal curvatures of the spine, such as scoliosis. Osteoarthritis can begin in
 adolescence.

When to see a doctor

Regular follow-up visits, as well as yearly visits to doctors specializing in eye disorders, are crucial to monitor any progression of symptoms. Early treatment can help prevent life-altering complications. Hearing should be checked every six months in children through age 5 and then yearly thereafter.

Causes

Stickler syndrome is caused by mutations in certain genes involved in the formation of collagen — one of the building blocks of many types of connective tissues. The type of collagen most commonly affected is that used to produce joint cartilage and the jellylike material (vitreous) found within the eyes.

Risk factors

Your child is more likely to be born with Stickler syndrome if either you or your partner has the disorder.

Complications

Potential complications of Stickler syndrome include:

- **Difficulty breathing or feeding.** Breathing or feeding difficulties may occur in babies born with an opening in the roof of the mouth (cleft palate), a small lower jaw and a tendency for the tongue to drop back toward the throat.
- Blindness. Blindness can occur if retinal detachments aren't repaired promptly.
- Ear infections. Children with facial structure abnormalities associated with Stickler syndrome are more likely to develop ear infections than are children with normal facial structure.
- Deafness. Hearing loss associated with Sticker syndrome may worsen with time.
- Heart problems. Some people with Stickler syndrome may be at higher risk of heart valve problems.
- **Dental problems.** Most children who have Stickler syndrome have abnormally small jaws, so there often isn't enough room for the full complement of adult teeth. Braces or, in some cases, dental surgeries may be necessary.

By Mayo Clinic Staff

Any use of this site constitutes your agreement to the Terms and Conditions and Privacy Policy linked below.

Terms and Conditions

Privacy Policy

Notice of Privacy Practices

Notice of Nondiscrimination

Mayo Clinic is a nonprofit organization and proceeds from Web advertising help support our mission. Mayo Clinic does not endorse any of the third party products and services advertised.

Advertising and sponsorship policy

Advertising and sponsorship opportunities

A single copy of these materials may be reprinted for noncommercial personal use only. "Mayo," "Mayo Clinic," "MayoClinic.org," "Mayo Clinic Healthy Living," and the triple-shield Mayo Clinic logo are trademarks of Mayo Foundation for Medical Education and Research.

© 1998-2020 Mayo Foundation for Medical Education and Research (MFMER). All rights reserved.