

What Makes a Wine Good? Chemical Structure, Wine Type, and Quality in Portuguese Wines

Avi Gobrin

Abstract

Understanding the chemical factors associated with perceived wine quality and the structural differences between red and white wines requires multivariate statistical analysis. This study examines a dataset of 6,497 Portuguese wines described by physicochemical measurements, wine type, and ordinal sensory quality. Exploratory visualization, correlation analysis, principal component analysis, assessment of multivariate normality, and comparison of covariance and correlation based principal component methods are used to construct a coherent statistical explanation of wine quality and wine type. Alcohol content is moderately positively associated with quality, while density exhibits negative association, indicating that fermentation strength contributes to perceived quality. Standardized principal component analysis shows that approximately one half of total chemical variability is governed by two latent dimensions corresponding to preservation chemistry and fermentation driven alcohol structure. Projection into this low dimensional linear subspace reveals clear separation between red and white wines, demonstrating that wine type differences arise from continuous chemical composition. Departures from multivariate normality reflect realistic chemical heterogeneity rather than modeling failure. Together, these results provide a quantitative explanation of what distinguishes higher quality wines and how red and white wines differ chemically.

Introduction

Perceived wine quality is the outcome of complex biochemical and physical processes involving fermentation, preservation, acidity balance, and sugar metabolism. Measurements such as sulfur dioxide concentration, density, alcohol content, and residual sugar therefore arise from interacting chemical mechanisms rather than independent variation. Because of this dependence, identifying the chemical properties associated with high quality wine and understanding how red and white wines differ requires a multivariate statistical perspective.

The present analysis investigates a dataset of Portuguese wines containing physicochemical measurements, categorical wine type, and ordinal sensory quality scores originally introduced by Cortez et al. [1]. The central objective is to determine which measurable chemical properties are associated with higher quality evaluations and whether the apparent complexity of wine chemistry can be summarized through a smaller number of latent dimensions. Particular emphasis is placed on interpreting principal component structure in scientifically meaningful terms and relating this structure to both quality and wine type.

Chemical Composition and Distributional Structure

The dataset contains 6,497 wine observations measured across 13 variables, including continuous physicochemical measurements, wine type classification, and an ordinal quality score ranging from 3 to 9. White wines constitute the majority of observations, while red wines form a smaller but substantial subset. Quality scores are concentrated at moderate values rather than extreme ratings, indicating that the dataset reflects realistic commercial production conditions.

Several physicochemical variables display strong right skewness and extreme but chemically plausible observations. Chlorides exhibit particularly pronounced skewness, and sulphates, fixed acidity, volatile acidity, residual sugar, and free sulfur dioxide also show clear asymmetry. These distributional properties provide evidence against multivariate normality and suggest that the chemical environment of wine production includes occasional extreme conditions rather than symmetric variability. Formal comparison of Mahalanobis distances with theoretical chi square quantiles confirms systematic deviation in the upper tail, demonstrating heavier than Gaussian behavior in multivariate space. This finding clarifies that principal component analysis should be interpreted as a descriptive representation of chemical structure rather than a consequence of normal theory inference.




Figure 1: Distributions of selected physicochemical variables by wine type. Strong skewness and extreme observations indicate realistic chemical heterogeneity and departure from multivariate normality.

Associations Between Chemical Measurements and Wine Quality

Correlation analysis provides direct quantitative evidence regarding which physicochemical characteristics are associated with higher quality scores. Alcohol content exhibits a moderate positive relationship with quality, while density shows a negative association. These relationships are chemically consistent with fermentation processes in which conversion of sugar into ethanol reduces density and increases alcohol concentration. Wines that undergo more complete fermentation therefore tend, on average, to receive higher sensory evaluations. The statistical evidence indicates that fermentation related properties, particularly higher alcohol concentration and lower density, are the physicochemical characteristics most consistently associated with improved quality scores. However, the moderate strength of these relationships demonstrates that chemical composition alone does not fully determine perceived wine quality. Sensory evaluation depends on additional structural, aromatic, and experiential factors not captured by the measured variables. Principal component analysis therefore clarifies which chemical mechanisms contribute to quality while simultaneously revealing the limits of prediction based solely on physicochemical measurements.

Sulfur dioxide measurements show strong mutual dependence, reflecting shared preservation chemistry rather than direct influence on quality. Residual sugar and density are positively related, again consistent with incomplete fermentation. Taken together, these relationships suggest that fermentation completeness and alcohol formation play a more prominent role in perceived quality than preservation chemistry alone.




Figure 2: Correlation structure among physicochemical variables and quality. Alcohol and density display the strongest relationships with quality, consistent with fermentation driven mechanisms.

Low Dimensional Chemical Structure

Principal component analysis conducted on standardized physicochemical variables provides quantitative evidence that the apparent complexity of wine chemistry can be substantially reduced without severe information loss. The first principal component explains approximately 27.5 percent of total standardized variance, and the second explains approximately 22.7 percent. Together these two components account for roughly 50.2 percent of overall variability. This result indicates that one half of the variation present across all measured chemical properties can be represented within a two dimensional linear subspace. This low dimensional representation is optimal in the sense that the first (k) principal components provide the minimum error rank- (k) linear reconstruction of the centered data matrix, a fundamental property of principal component analysis described in standard multivariate texts.

Additional dimensionality criteria further clarify the extent to which the original measurement space may be compressed. The Kaiser criterion suggests preservation of three principal components.

From the perspective of cumulative explained variance, approximately six components are required to account for at least 80 percent of total variability, while seven components are required to exceed 90 percent. These thresholds imply that the full thirteen dimensional chemical description contains substantial redundancy, since a representation using roughly half as many dimensions preserves the vast majority of information contained in the data.

Interpretation of these results depends on the analytical objective. When visualization and qualitative understanding are the primary goals, the first two principal components provide a meaningful representation that captures dominant biochemical variation. When higher fidelity reconstruction is required, retaining between six and seven components ensures that most variability is preserved. In either case, the eigenvalue structure demonstrates that wine chemistry lies near a lower dimensional manifold determined by underlying fermentation and preservation processes.




Figure 3: Eigenvalue decay across principal components supporting dimensional reduction.




Figure 4: Cumulative variance explained by principal components. Approximately one half of standardized variability is captured by two components.

Chemical Differences Between Red and White Wines

Projection of wine observations onto the first two principal components reveals clear separation between red and white wines. This separation demonstrates that wine type differences arise primarily from continuous chemical composition rather than categorical labeling alone. Red and white wines occupy distinct regions of the preservation and fermentation structure defined by the principal components, indicating systematic biochemical differences in production methods and composition.




Figure 5: Projection of wines into the two dimensional principal component space. Distinct clustering indicates chemically driven separation between red and white wines.

Role of Standardization in Revealing Structure

Principal component analysis based on the covariance matrix produces a first component that explains nearly all variance due to dominance of high variance variables such as density and sulfur dioxide. This dominance obscures meaningful multivariate structure and prevents interpretation of latent chemical processes. Standardization equalizes measurement scales and reveals interpretable dimensions related to fermentation and preservation. The comparison demonstrates that correlation based principal component analysis is essential for meaningful analysis of heterogeneous chemical data.

Conclusion

Multivariate statistical analysis of Portuguese wine chemistry reveals that perceived wine quality is most strongly associated with fermentation related properties, particularly alcohol concentration and density. Principal component analysis shows that the apparent complexity of physicochemical measurements can be summarized through a small number of latent biochemical dimensions corresponding to preservation chemistry and fermentation structure. Differences between red and white wines emerge naturally within this low dimensional representation, indicating that wine type

reflects continuous chemical variation rather than purely categorical distinction. Departures from multivariate normality reflect realistic chemical heterogeneity and do not undermine the descriptive value of principal component methods. More broadly, the analysis demonstrates that the effective dimensionality of wine chemistry is substantially lower than the number of measured variables, allowing complex biochemical variation to be summarized through a small number of interpretable latent processes.

References

- [1] Cortez, P., Cerdeira, A., Almeida, F., Matos, T., and Reis, J. Modeling wine preferences by data mining from physicochemical properties. *Decision Support Systems*, 47(4), 547–553, 2009.
- [2] STA437 Course Notes. Weeks 3–5: Multivariate normality and principal component analysis. University of Toronto, 2026.

AI Attribution

Generative artificial intelligence tools such as CHATGPT were used in a limited and supportive capacity during the preparation of this report. Specifically, AI assistance was used to help refine written language, improve clarity of academic phrasing, and organize the overall structure of the document. All statistical analyses, numerical computations, graphical outputs, and substantive interpretations were performed independently by the author using the provided dataset and course methods. No analytical results, code outputs, or mathematical conclusions were generated solely by AI tools. Every statement of methodology and interpretation was reviewed and verified for correctness against the course lecture materials and the observed data. The use of AI therefore functioned only as an editorial and organizational aid rather than as a source of analytical content.