

Existence of Solutions for a Stationary Mean Field Game System

Avi Wine

March 11, 2026

Abstract

This report studies a stationary mean field game system given by a Hamilton–Jacobi–Bellman equation coupled with a Kolmogorov–Fokker–Planck equation. Under suitable assumptions, we establish the existence of a weak solution pair in $H_0^1(\Omega) \times H_0^1(\Omega)$, and we extend the argument to a variant with a non-linear source term in the KFP equation.

Contents

1	Introduction	2
2	Method	4
2.1	Independent unique solutions	4
2.1.1	Solving the Kolmogorov–Fokker–Planck Equation	4
2.1.2	Solving Hamilton–Jacobi–Bellman	6
2.2	Solving the Coupled System of Equations	9
2.2.1	Continuity of the Fixed Point Map	10
2.2.2	Compactness of the Fixed Point Map	11
2.2.3	Boundedness of the Fixed Point Set	13
2.3	Swapping Composition Order of T and S	14
3	Extension to Non-linear KFP	15
4	Conclusion	19

1 Introduction

Context and Motivation

Mean field games is a branch of game theory devoted to the analysis of a special subset of games. These systems are distinguished by having a very large number of players, each of whom is considered almost insignificant, with minimal influence on the system as a whole. In this framework the discrete population of agents is replaced by a continuum distribution, allowing the system to be described using density functions and weak (distributional) solutions of partial differential equations.

Furthermore, each player is a rational observer and the general trend around them informs their decision on how to act, with the aim of achieving an optimal outcome. The famous example in economics of such a game, introduced by Aumann, is the behaviour of stock traders whom we treat as a continuum [1]. In this paper we consider the stationary equilibrium problem characterised by a coupled pair of equations: a Hamilton–Jacobi–Bellman (HJB) equation and a Kolmogorov–Fokker–Planck (KFP) equation.

$$\begin{aligned} -\Delta u + H(x, \nabla u) + \lambda u &= f(x, m) \quad \text{in } \Omega \text{ HJB,} \\ -\Delta m - \operatorname{div}\left(m \frac{\partial H}{\partial p}(x, \nabla u)\right) + \lambda m &= G(x) \quad \text{in } \Omega \text{ KFP.} \end{aligned} \tag{1.1}$$

This is supplemented with homogeneous Dirichlet boundary conditions $u = 0$ and $m = 0$ on $\partial\Omega$.

Interpretation of Equations

Here $m(x)$ represents the player density and $u(x)$ the value function of a representative agent in the mean field game. In other words, $u(x)$ is a player’s optimal solution achieved by minimising the expected cost at position x . The term $\frac{\partial H}{\partial p}$ denotes the partial derivative of H with respect to the momentum variable (i.e. the ∇u argument), and $\lambda > 0$. HJB describes a representative player’s optimal control behaviour for a given player density m .

On the other hand, KFP gives the evolution of $m(\cdot)$ when every player satisfies the optimal solution u . Each player tries to minimise their cost function $f(x, m)$ (in HJB), and the optimal control for an individual player is given by $\alpha^* = -\frac{\partial H}{\partial p}(x, \nabla u)$, which makes up the drift term in KFP. The Laplacian terms are diffusion terms accounting for randomness (e.g. Brownian motion) and preventing perfect alignment of players even under a shared strategy. Finally, $G(\cdot)$ is an exogenous term controlling entry/exit of players and therefore directly affects m in KFP. [4, 2]

Mathematical Contribution

In this paper we analyse the coupled system equation (1.1) and investigate the existence of a weak solution pair $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$. Establishing the existence of a solution pair (u, m) is important because such solutions correspond to Nash equilibria of the mean field

game. The function u determines the optimal strategy of a representative agent, while m describes the distribution of agents when all players adopt this strategy. A solution of the coupled system therefore represents a self-consistent equilibrium of the game. In addition, we consider a more general formulation of the system and prove an existence result in this extended setting, which constitutes an original contribution of this work.

Assumptions

1. Ω is bounded and $\partial\Omega$ is smooth.
2. $H(x, \nabla u) : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}$ is
 - Lipschitz continuous with respect to ∇u with Lipschitz constant $\tilde{C}$,
 - bounded by $C(|\nabla u| + 1)$ for a constant C , for all $x \in \Omega$ and $\nabla u \in \mathbb{R}^n$.
3. The partial derivative $\frac{\partial H}{\partial p}(x, \nabla u)$ is L^2 continuous and essentially bounded by some constant L .
4. λ is sufficiently large (in particular $\lambda > \max(\frac{L^2}{4}, \frac{C^2}{2}, \frac{\tilde{C}^2}{4})$).
5. $f(x, m(x)) : \Omega \times L^2(\Omega) \rightarrow L^2(\Omega)$ is
 - continuous with respect to m .
 - bounded by $D(|m| + 1)$ for a constant D , for all $x \in \Omega$ and $m \in L^2(\Omega)$.
6. $G(x) \in L^2(\Omega)$.

Example 1.1. A Hamiltonian satisfying Assumption 2 is

$$H(x, p) = \sqrt{1 + |p|^2} + V(x) \quad \text{on } \Omega = B(0, 1) \text{ in } \mathbb{R}^n,$$

where $V \in L^\infty(\Omega)$. In this case, we get the associated drift

$$\frac{\partial H}{\partial p} = \frac{p}{\sqrt{1 + |p|^2}}.$$

Example 1.2. For a coupling term satisfying Assumption 5, one may consider

$$f(x, m) = a(x) \phi(m),$$

where $a(x) \in L^\infty(\Omega)$ and $\phi : L^2(\Omega) \rightarrow L^2(\Omega)$ is continuous.

Main Result

Theorem 1.3 (Existence of the MFG solution). *Let the Mean Field Game system be characterised by equation (1.1) and suppose the assumptions above hold. Then a solution $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$ exists.*

We prove this theorem throughout the rest of this paper.

2 Method

Our proof consists of two stages. First, we demonstrate solvability for each equation separately, while fixing the other variable (HJB with fixed m , and KFP with fixed u). Second, we show the existence of a solution pair solving both equations simultaneously. Before starting our proof it is important to introduce the following inequalities, which will be used repeatedly in this paper.

Definition 2.1. Parameters $p, q \in [1, \infty]$ are *conjugate* if $\frac{1}{p} + \frac{1}{q} = 1$.

Theorem 2.2 (Hölder's inequality). *Let $f \in L^p(\Omega)$ and $g \in L^q(\Omega)$ where p, q are conjugate. Then*

$$\|fg\|_1 \leq \|f\|_p \|g\|_q.$$

Remark 2.3. In particular, when $p = q = 2$ we obtain the Cauchy–Schwarz inequality

$$\|fg\|_1 \leq \|f\|_2 \|g\|_2.$$

Theorem 2.4 (Young's inequality). *Let $a, b \in \mathbb{R}$. Then*

$$2ab \leq \frac{a^2}{\varepsilon^2} + \varepsilon^2 b^2 \quad \forall \varepsilon > 0.$$

2.1 Independent unique solutions

2.1.1 Solving the Kolmogorov–Fokker–Planck Equation

To solve

$$-\Delta m - \operatorname{div}\left(m \frac{\partial H}{\partial p}(x, \nabla u)\right) + \lambda m = G(x)$$

with fixed u , we note that the equation is linear in $m(x)$, so we appeal to the Lax–Milgram theorem [5].

Theorem 2.5 (Lax–Milgram). *Let V be a Hilbert space and let $a : V \times V \rightarrow \mathbb{R}$ be a bilinear form such that for all $u, v \in V$,*

$$(i) \text{ (boundedness) } |a(u, v)| \leq c_1 \|u\| \|v\|,$$

$$(ii) \text{ (coercivity) } a(v, v) \geq c_0 \|v\|^2.$$

Let $l : V \rightarrow \mathbb{R}$ be a bounded linear functional. Then there exists a unique $u \in V$ such that

$$a(u, v) = l(v) \quad \text{for all } v \in V.$$

Moreover, $\|u\|_V \leq \frac{1}{c_0} \|l\|_{V^}$.*

In our case,

$$a(m, v) = \int_{\Omega} \nabla m \cdot \nabla v + m \nabla v \cdot \frac{\partial H}{\partial p}(x, \nabla u) + \lambda m v \, dx, \quad l(v) = \int_{\Omega} G(x) v \, dx. \quad (2.1)$$

We now show that KFP meets the conditions of Theorem 2.5.

Proposition 2.6 (Boundedness of KFP). *The form $a(m, v)$ is bounded on $H_0^1(\Omega) \times H_0^1(\Omega)$.*

Proof. For arbitrary $m, v \in H_0^1(\Omega)$,

$$\begin{aligned} |a(m, v)| &\leq \int_{\Omega} |\nabla m| |\nabla v| + |\nabla v| |m| \left| \frac{\partial H}{\partial p} \right| + \lambda |m| |v| dx \\ &\leq \|\nabla m\|_{L^2} \|\nabla v\|_{L^2} + L \|m\|_{L^2} \|\nabla v\|_{L^2} + \lambda \|m\|_{L^2} \|v\|_{L^2}. \end{aligned}$$

By Cauchy–Schwarz and Assumption 3, there exists $c_1 > 0$ such that

$$|a(m, v)| \leq c_1 \|m\|_{H^1} \|v\|_{H^1}.$$

Thus $a(m, v)$ is bounded. $\square$

Proposition 2.7 (Coercivity of KFP). *If $\lambda > 0$ is sufficiently large, then $a(\cdot, \cdot)$ is coercive.*

Proof. Consider

$$\begin{aligned} a(v, v) &= \int_{\Omega} |\nabla v|^2 + v \nabla v \cdot \frac{\partial H}{\partial p}(x, \nabla u) + \lambda v^2 dx \\ &\geq \|\nabla v\|_{L^2}^2 - L \|v\|_{L^2} \|\nabla v\|_{L^2} + \lambda \|v\|_{L^2}^2. \end{aligned}$$

Applying Young’s inequality yields

$$a(v, v) \geq \left(1 - \frac{\varepsilon^2}{2}\right) \|\nabla v\|_{L^2}^2 + \left(\lambda - \frac{L^2}{2\varepsilon^2}\right) \|v\|_{L^2}^2, \quad \forall \varepsilon \in \mathbb{R}.$$

To ensure $1 - \varepsilon^2/2 > 0$, impose $\varepsilon^2 < 2$. Choosing $\varepsilon^2 = 2$ (as a limiting choice) gives the weakest requirement

$$\lambda > \frac{L^2}{4},$$

which ensures coercivity. $\square$

We state the main result of this chapter.

Theorem 2.8 (Existence of unique KFP solution). *Let the Kolmogorov–Fokker–Planck equation be as in equation (1.1) and suppose that $\lambda > 0$ is sufficiently large. Then for any fixed $u \in H_0^1(\Omega)$, there exists a unique $m \in H_0^1(\Omega)$ solving the equation.*

Proof. Let $a(m, v) = l(v)$ be the weak formulation as in equation (2.1). By Propositions 2.6 and 2.7, $a(\cdot, \cdot)$ is bounded and coercive. It remains to show l is bounded. By Cauchy–Schwarz,

$$\|l\| = \sup_{\|v\|_{H^1} \leq 1} \left| \int_{\Omega} G v dx \right| \leq \sup_{\|v\|_{L^2} \leq 1} \|v\|_{L^2} \|G\|_{L^2} \leq \|G\|_{L^2(\Omega)}.$$

Thus l is bounded. By Theorem 2.5, there exists a unique $m \in H_0^1(\Omega)$ solving the KFP equation. $\square$

2.1.2 Solving Hamilton–Jacobi–Bellman

Lax–Milgram does not apply directly to

$$-\Delta u + H(x, \nabla u) + \lambda u = f(x, m)$$

with fixed m , because the equation is non-linear so we resort to a fixed point method instead. The idea, is to create a map from the HJB equation with the right properties such that it has a fixed point and for that very fixed point to be our solution. Remarkably, this fixed point method is general enough that we can use it to solve for existence in the system as whole. Indeed, this will be the concern of the next subsection. Throughout this section, X denotes a real Banach space. We now introduce the fixed point theorems that rely on suitable properties for the HJB context.

Definition 2.9. A mapping $A : X \rightarrow X$ is called *compact* if for every bounded sequence $(u_k)_{k=1}^\infty$, the sequence $(A[u_k])_{k=1}^\infty$ is precompact in X .

Theorem 2.10 (Schaefer’s fixed point theorem). *Suppose $A : X \rightarrow X$ is continuous and compact, and assume the set*

$$\{u \in X : u = \mu A[u] \text{ for some } 0 \leq \mu \leq 1\}$$

*is bounded. Then A has a fixed point.*¹

Let us now design the map whose fixed point is a solution to the HJB equation. Consider

$$A : H_0^1(\Omega) \rightarrow H_0^1(\Omega), \quad A[u] = w \quad \text{where} \quad -\Delta w + \lambda w = f(x, m) - H(x, \nabla u).$$

We know w exists by Lax–Milgram (Theorem 2.5), so A is well defined.² Furthermore by regularity theory³ we have

$$\begin{aligned} \|w\|_{H^2(\Omega)} &\leq C \|f(x, m) - H(x, \nabla u)\|_{L^2(\Omega)} \\ &\leq C \|f(x, m)\|_{L^2(\Omega)} + C \|H(x, \nabla u)\|_{L^2(\Omega)} \\ &\leq C \|f(x, m)\|_{L^2(\Omega)} + \hat{C} \|\nabla u\| + 1 \|1\|_{L^2(\Omega)} \quad (\text{Assumption 2}) \end{aligned}$$

Where $C, \hat{C}$ are constants - note that these constants are not the same as the ones appearing in our assumptions above. We now prove the existence of an HJB solution by way of our fixed point map.

Theorem 2.11 (Existence of HJB Solution). *If $\lambda > 0$ is sufficiently large, then for any fixed $m \in H_0^1(\Omega)$ there exists $u \in H^2(\Omega) \cap H_0^1(\Omega)$ solving HJB.*

¹See [3, Section 9.2].

²LHS is bounded and coercive and RHS is bounded by linear growth condition on H .

³See [3, Section 6.3]

Proof. ⁴ In other words, for sufficiently large $\lambda > 0$, A has a fixed point. We show first that A meets the right criteria for Schaefer's Theorem. We assert that A is continuous and compact.

If $u_k \rightarrow u$ in $H_0^1(\Omega)$ and we denote $w_k = A[u_k]$ then higher regularity tells us

$$\sup_k \|w_k\|_{H^2(\Omega)} < \infty$$

(i.e. (w_k) is a bounded sequence in $H^2(\Omega) \cap H_0^1(\Omega)$) and there exists a convergent subsequence $(w_{k_j})_{j=1}^\infty \rightarrow w$ in $H_0^1(\Omega)$ ⁵.

Now the weak form of the HJB equation with w_{k_j} is

$$\int_{\Omega} \nabla w_{k_j} \cdot \nabla v + \lambda w_{k_j} \cdot v \, dx = \int_{\Omega} (f(x, m) - H(x, \nabla u_{k_j})) \cdot v \, dx, \text{ for any } v \in H_0^1(\Omega)$$

and consequently

$$\int_{\Omega} \nabla w \cdot \nabla v + \lambda w \cdot v \, dx = \int_{\Omega} (f(x, m) - H(x, \nabla u)) \cdot v \, dx, \text{ for any } v \in H_0^1(\Omega)$$

by Lipschitz continuity of H (Assumption 2). Thus, $w = A[u]$. Hence, $A[u_k] \rightarrow A[u]$ in $H_0^1(\Omega)$, so A is continuous. A is compact by similar reasoning since a bounded sequence $(u_k)_{k=1}^\infty$ yields a bounded sequence $(A[u_k])_{k=1}^\infty$ in $H^2(\Omega)$ which has a strongly convergent subsequence in $H_0^1(\Omega)$.

It remains to show

$$\{u \in H_0^1(\Omega) : u = \mu A[u] \text{ for some } 0 \leq \mu \leq 1\}$$

is bounded.

Substituting $w = u/\mu$ ⁶ and setting $v = u$, we obtain

$$\begin{aligned} \int_{\Omega} |\nabla u|^2 + \lambda |u|^2 \, dx &= \mu \int_{\Omega} f u - H(x, \nabla u) u \, dx \\ &\leq \int_{\Omega} |f u| + |H(x, \nabla u) u| \, dx, & \text{since } 0 < \mu \leq 1 \\ &\leq \int_{\Omega} |f u| + C(|\nabla u| + 1)|u| \, dx & \text{(Assumption 2)} \end{aligned}$$

To isolate $|\nabla u|^2$ and u^2 terms on the RHS, we apply Young's inequality (Theorem 2.4) once

⁴Proof is adapted from [3, Section 9.2], with several modifications specific to the present setting.

⁵See Corollary 2.26

⁶ We ignore $\mu = 0$ since then $w = 0$

on $|fu|$ and twice over on $C(|\nabla u| + 1)|u|$, to obtain

$$|fu| \leq \frac{1}{2} \left(\frac{f^2}{\varepsilon^2} + \varepsilon^2 u^2 \right) \quad \varepsilon \in (0, \infty)$$

and

$$\begin{aligned} C(|\nabla u| + 1)|u| &\leq \frac{1}{2} \left(\frac{1}{\delta^2} |Cu|^2 + \delta^2 (|\nabla u| + 1)^2 \right), \quad \delta \in (0, 1) \\ &\leq \frac{1}{2} \left(\frac{1}{\delta^2} |Cu|^2 + \delta^2 (1 + \gamma^2) |\nabla u|^2 + \delta^2 (1 + \frac{1}{\gamma^2}) \right) \\ &= \frac{1}{2} \left(\frac{1}{\delta^2} |Cu|^2 + 2\delta^2 |\nabla u|^2 + 2\delta^2 \right) \quad \text{WLOG setting } \gamma = 1. \end{aligned}$$

Returning to the weak form inequality we get

$$\int_{\Omega} |\nabla u|^2 + \lambda |u|^2 dx \leq \frac{1}{2} \int_{\Omega} \frac{f^2}{\varepsilon^2} + \left(\varepsilon^2 + \frac{C^2}{\delta^2} \right) u^2 + 2\delta^2 |\nabla u|^2 + 2\delta^2 dx.$$

Provided $\lambda > \frac{C^2}{2}$ there exist sufficiently small $\varepsilon > 0$ and sufficiently large $\delta < 1$ such that

$$\lambda > \frac{1}{2} \left(\varepsilon^2 + \frac{C^2}{\delta^2} \right) \text{ and } 1 > \delta^2$$

hold simultaneously. Taking the difference of both sides and denoting the constant factor as $K^2 := \int_{\Omega} \delta^2 dx$, yields

$$\alpha^2 \int_{\Omega} |\nabla u|^2 + |u|^2 dx \leq \frac{1}{2\varepsilon^2} \int_{\Omega} f(x, m)^2 dx + K^2,$$

where α^2 is the minimum of both factors of u^2 and $|\nabla u|^2$. Consequently,

$$\alpha \|u\|_{H_0^1(\Omega)} \leq \frac{1}{\varepsilon\sqrt{2}} \|f(x, m)\|_{L^2(\Omega)} + K.$$

Thus if $\lambda > \frac{C^2}{2}$, the set

$$\{u \in H_0^1(\Omega) : u = \mu A[u] \text{ for some } 0 \leq \mu \leq 1\}$$

is bounded independently of μ . By Theorem 2.10, A has a fixed point $u \in H_0^1(\Omega)$ and thus the HJB equation has a solution. $\square$

Remark 2.12. We can always tune ε and δ for given $\lambda > \frac{C^2}{2}$ to satisfy boundedness. This utilises the strictness of the inequality in the condition on λ .

Proposition 2.13. *If $\lambda > 0$ is sufficiently large, then the fixed point u is unique.*

Proof. Suppose $u_1, u_2 \in H_0^1(\Omega)$ both solve the HJB equation for a given $m \in H_0^1(\Omega)$. Taking the weak difference and setting $v = u_1 - u_2$ gives

$$\begin{aligned} \|\nabla(u_1 - u_2)\|_{L^2}^2 + \lambda \|u_1 - u_2\|_{L^2}^2 &\leq \int_{\Omega} |u_1 - u_2| |H_1 - H_2| dx \\ &\leq \tilde{C} \|u_1 - u_2\|_{L^2} \|\nabla(u_1 - u_2)\|_{L^2} \end{aligned}$$

using Assumption 2 and Cauchy–Schwarz. Hence, by Young’s inequality,

$$\|\nabla(u_1 - u_2)\|_{L^2}^2 + \lambda \|u_1 - u_2\|_{L^2}^2 \leq \frac{1}{2} \left(\frac{\tilde{C}^2}{\varepsilon^2} \|u_1 - u_2\|_{L^2}^2 + \varepsilon^2 \|\nabla(u_1 - u_2)\|_{L^2}^2 \right) \quad \varepsilon \in (0, \sqrt{2}).$$

By the same reasoning as when we proved existence (Theorem 2.11),

$$\lambda > \frac{\tilde{C}^2}{4} \implies u_1 = u_2 \text{ in } L^2(\Omega) \iff u_1 = u_2 \text{ in } H_0^1(\Omega).$$

□

The following theorem summarises our results.

Theorem 2.14 (Existence of unique HJB solution). *Let the Hamilton-Jacobi equation be as in equation (1.1) and suppose that $\lambda > 0$ is sufficiently large. Then for any fixed $m \in H_0^1(\Omega)$, there exists a unique $u \in H_0^1(\Omega)$ solving the equation.*

Proof. There exists a solution by Theorem 2.11 and it is unique by Proposition 2.13. □

2.2 Solving the Coupled System of Equations

Having solved the two equations in 1.1 in isolation, we now look to solve them as a system. As stated above, the fixed point theorems will allow us to prove existence of a solution pair for the system as a whole. The challenge, is to define and prove the existence of a fixed point for a suitable map. Our work in the previous subsection allows us to define the following maps.

Definition 2.15. Define $S : H_0^1(\Omega) \rightarrow H_0^1(\Omega)$ by $S[u] = m$, where m solves KFP for given u .

Definition 2.16. Define $T : H_0^1(\Omega) \rightarrow H_0^1(\Omega)$ by $T[m] = u$, where u solves HJB for given m .

Definition 2.17. Let T and S be as in definitions definition 2.15 and definition 2.16. Define

$$A : H_0^1(\Omega) \rightarrow H_0^1(\Omega), \quad A := S \circ T.$$

That is, A takes $m \in H_0^1(\Omega)$, converts it to $u = T[m]$ solving the HJB equation, and then produces $\tilde{m} = S[u]$ solving the KFP equation. If A has a fixed point, then $m = \tilde{m}$, and the coupled system has a solution pair.

We verify the hypotheses of Schaefer's theorem.

2.2.1 Continuity of the Fixed Point Map

Lemma 2.18. *Let A be as in definition 2.17 and suppose that $\lambda > 0$ as it appears in equation (1.1) is sufficiently large. Then, A is continuous in $H_0^1(\Omega)$.*

Since A is simply a composition of S and T , it suffices to show they are both continuous.

Proposition 2.19. *If $\lambda > 0$ is sufficiently large, T is continuous in $H_0^1(\Omega)$.*

Proof. Let (m_k) converge to $m \in H_0^1(\Omega)$. Denote $u = T[m]$, $u_k = T[m_k]$, $H = H(x, \nabla u)$, $H_k = H(x, \nabla u_k)$ and $f_k = f(x, m_k)$. Taking the weak formulation of the difference in HJB gives

$$\begin{aligned} \|\nabla(u - u_k)\|_{L^2}^2 + \lambda \|u - u_k\|_{L^2}^2 &\leq \int_{\Omega} |u - u_k| |H - H_k| + |u - u_k| |f - f_k| dx \\ &\leq \tilde{C} \|u - u_k\|_{L^2} \|\nabla(u - u_k)\|_{L^2} + \|u - u_k\|_{L^2} \|f - f_k\|_{L^2}. \end{aligned} \tag{1}$$

By Cauchy-Schwarz and Assumption 2. Since f is $L^2(\Omega)$ -continuous (Assumption 5) and

$$m_k \rightarrow m \text{ in } H_0^1(\Omega) \implies m_k \rightarrow m \text{ in } L^2(\Omega),$$

$\|f - f_k\|_{L^2} \rightarrow 0$. Also, from the energy estimate

$$\|\nabla u_k\|_{L^2}^2 + \lambda \|u_k\|_{L^2}^2 \leq \int_{\Omega} |u_k| |f_k| + |u_k| |H_k| dx.$$

Adopting the same argument for boundedness as in Theorem 2.11 gives the inequality

$$\alpha \|u_k\|_{H_0^1(\Omega)} \leq \frac{1}{\varepsilon \sqrt{2}} \|f_k\|_{L^2(\Omega)} + K.$$

where K , α and ε are independent (positive) constants. By the fact that $f_k \rightarrow f$ we know that $\|f_k\|_{L^2(\Omega)}$ is bounded and so therefore (u_k) is bounded in $H_0^1(\Omega)$. Thus, $\|u - u_k\|_{L^2} \|f - f_k\|_{L^2} \rightarrow 0$. Thus, in the limit of k , (1) becomes

$$\|\nabla(u - u_k)\|_{L^2}^2 + \lambda \|u - u_k\|_{L^2}^2 \leq \tilde{C} \|u - u_k\|_{L^2} \|\nabla(u - u_k)\|_{L^2}$$

for which $\lambda > 0$ being sufficiently large implies $u_k \rightarrow u$ i.e T is continuous on $H_0^1(\Omega)$. ⁷ $\square$

⁷See the uniqueness argument above (Proposition 2.13).

Proposition 2.20. *If $\lambda > 0$ is sufficiently large, S is continuous in $H_0^1(\Omega)$.*

Proof. Let (u_k) converge to u in $H_0^1(\Omega)$. Denote $m = S[u]$, $m_k = S[u_k]$, $b = \frac{\partial H}{\partial p}(x, \nabla u)$ and $b_k = \frac{\partial H}{\partial p}(x, \nabla u_k)$. Taking the weak formulation of the difference in KFP yields

$$\|\nabla(m - m_k)\|_{L^2}^2 + \lambda \|m - m_k\|_{L^2}^2 \leq \int_{\Omega} |\nabla(m - m_k)| |m b - m_k b_k| dx.$$

Define the correction vector field $\beta_k := b - b_k$. Note that since b_k is L^2 continuous with respect to u_k we have $\|\beta_k\|_{L^2(\Omega)} \rightarrow 0$ (Assumption 3). Then,

$$\begin{aligned} \|\nabla(m - m_k)\|_{L^2}^2 + \lambda \|m - m_k\|_{L^2}^2 &\leq \int_{\Omega} |\nabla(m - m_k)| |m b - m_k b| + |\nabla(m - m_k)| |m_k| |\beta_k| dx \\ &\leq L \int_{\Omega} |\nabla(m - m_k)| |m - m_k| dx + \int_{\Omega} |\nabla(m - m_k)| |m_k| |\beta_k| dx. \end{aligned}$$

m and m_k are bounded in $H_0^1(\Omega)$ so as k gets large

$$\int_{\Omega} |\nabla(m - m_k)| |m_k| |\beta_k| dx \rightarrow 0.$$

Therefore, by the largeness of λ (see Proposition 2.7), we get $m_k \rightarrow m$ i.e S is continuous in $H_0^1(\Omega)$. $\square$

Remark 2.21. The image of S is always bounded as a direct consequence of Lax-Milgram (Theorem 2.5).

Proof of Lemma 2.18. T is continuous (Proposition 2.19) and S is continuous (Proposition 2.20) so their composition is continuous in $H_0^1(\Omega)$. $\square$

2.2.2 Compactness of the Fixed Point Map

Lemma 2.22. *Let A be as in definition 2.17 and suppose that $\lambda > 0$ as it appears in equation (1.1) is sufficiently large. Then, A is compact in $H_0^1(\Omega)$.*

Since S is continuous, it suffices to show that T is compact in $H_0^1(\Omega)$:

$$T \text{ compact} \implies A = S \circ T \text{ compact}.$$

We introduce the following theorems. These will allow us to conclude convergence of a subsequence in a bigger space if we are given a bounded sequence in a smaller space. The particular relationship that concerns us is the one between $H^2(\Omega) \cap H_0^1(\Omega)$ and $H_0^1(\Omega)$.

Definition 2.23. Let X and Y be Banach spaces with $X \subset Y$. We say that X is compactly embedded in Y , written $X \subset\subset Y$, provided:

- (i) $\|x\|_Y \leq C \|x\|_X$ for some constant C ,
- (ii) each bounded sequence in X is precompact in Y .

Theorem 2.24 (Rellich–Kondrachov compactness theorem). *Assume U is a bounded open subset of $\mathbb{R}^n$ and ∂U is C^1 . Suppose $1 \leq p < n$. Then*

$$W^{1,p}(U) \subset\subset L^q(U) \quad \text{for each } 1 \leq q < p^*, \quad p^* = \frac{np}{n-p}.$$

Remark 2.25. Observe $p^* > p$. In particular,

$$W^{1,p}(U) \subset\subset L^p(U) \quad \text{for all } 1 \leq p < n.$$

Corollary 2.26.

$$H^2(\Omega) \cap H_0^1(\Omega) \subset\subset H_0^1(\Omega).$$

Proof. Firstly, we have

$$\|u\|_{H_0^1(\Omega)} \leq \|u\|_{H^2} \quad \text{for all } u \in H^2(\Omega) \cap H_0^1(\Omega).$$

Now let (u_k) be a bounded sequence in $H^2(\Omega) \cap H_0^1(\Omega)$. Then (u_k) and $(\partial_i u_k)$ are bounded in $H^1(\Omega)$ for each $i = 1, \dots, n$.

By the Rellich–Kondrachov compactness theorem, the embedding $H^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact. Hence there exists a subsequence, still denoted (u_k) , such that

$$u_k \rightarrow u \quad \text{in } L^2(\Omega),$$

and for each i ,

$$\partial_i u_k \rightarrow v_i \quad \text{in } L^2(\Omega).$$

Since weak derivatives are closed under L^2 convergence, it follows that $v_i = \partial_i u$. Therefore, by a diagonal argument there exists a common subsequence still denoted (u_k)

$$u_k \rightarrow u \quad \text{strongly in } H^1(\Omega).$$

Furthermore, $H_0^1(\Omega)$ is a closed subspace of $H^1(\Omega)$ so the limit lies in $H_0^1(\Omega)$. $\square$

We now have enough to show T is compact.

Proposition 2.27. *Let $\lambda > 0$ be sufficiently large. Then*

$$|f(x, m)| \leq D(|m| + 1) \implies T \text{ is compact in } H_0^1(\Omega).$$

Proof. Let (m_k) be a bounded sequence in $H_0^1(\Omega)$. Denote $u_k = T[m_k]$. We must show that (u_k) admits a subsequence converging strongly in $H_0^1(\Omega)$. From Theorem 2.11 we have the inequality

$$\alpha \|u_k\|_{H_0^1(\Omega)} \leq \frac{1}{\varepsilon\sqrt{2}} \|f(x, m_k)\|_{L^2(\Omega)} + K$$

where α, ε and K are independent (positive) constants. By the growth condition on f we

have

$$\begin{aligned}
\alpha \|u_k\|_{H_0^1(\Omega)} &\leq \frac{1}{\varepsilon\sqrt{2}} \|f(x, m_k)\|_{L^2(\Omega)} + K \\
&\leq \frac{1}{\varepsilon\sqrt{2}} (\|m_k\|_{L^2(\Omega)} + |\Omega|^{\frac{1}{2}}) + K \\
&\leq \frac{1}{\varepsilon\sqrt{2}} (\|m_k\|_{H_0^1(\Omega)} + |\Omega|^{\frac{1}{2}}) + K
\end{aligned}$$

which is uniformly bounded by our choice of a bounded sequence (m_k) in $H_0^1(\Omega)$. We also have that

$$\|H(x, \nabla u_k)\|_{L^2(\Omega)} \leq C(\|\nabla u_k\|_{L^2(\Omega)} + |\Omega|^{\frac{1}{2}})$$

which is uniformly bounded by Assumption 2 and the boundedness of (u_k) in $H_0^1(\Omega)$. This means that both $f_k = f(x, m_k)$ and $H_k = H(x, \nabla u_k)$ are bounded in $L^2(\Omega)$. Recall that u_k weakly solves

$$-\Delta u_k + \lambda u_k = g_k, \quad g_k := f_k - H_k.$$

By elliptic regularity for the Dirichlet problem, there exists $C > 0$ such that

$$\|u_k\|_{H^2(\Omega)} \leq C \|g_k\|_{L^2(\Omega)}$$

so (u_k) is bounded in $H^2(\Omega) \cap H_0^1(\Omega)$. By Corollary 2.26 there exists a subsequence (u_{k_j}) converging to $u \in H_0^1(\Omega)$. Thus T is compact, and hence A is compact in $H_0^1(\Omega)$. $\square$

We have shown that A is continuous and compact in $H_0^1(\Omega)$. The last step before we can apply Schauder's theorem is to prove that if there were to exist a fixed point, it would be bounded.

2.2.3 Boundedness of the Fixed Point Set

Lemma 2.28. *Let A be as in definition 2.17 then*

$$\{m \in H_0^1(\Omega) : m = \mu A[m] \text{ for some } \mu \in [0, 1]\}$$

is bounded in $H_0^1(\Omega)$.

Proof. Consider

$$\{m \in H_0^1(\Omega) : m = \mu A[m] \text{ for some } \mu \in [0, 1]\}.$$

The corresponding system, with $A[m]$ substituted for m/μ ⁸, is

$$-\Delta u + H(x, \nabla u) + \lambda u = f(x, m), \quad \frac{1}{\mu} \left(-\Delta m - \operatorname{div} \left(m \frac{\partial H}{\partial p}(x, \nabla u) \right) + \lambda m \right) = G(x).$$

⁸See 6

Rearranging the latter and using the Lax–Milgram bound implies

$$\|m\|_{H_0^1(\Omega)} \leq \frac{1}{c_0} \mu \|G\|_{L^2} \leq \frac{1}{c_0} \|G\|_{L^2},$$

which is bounded in $L^2(\Omega)$ (Assumption 6). $\square$

By Schaefer’s theorem, A has a fixed point. We may conclude our finding with the proof of Theorem 1.3.

Proof of the Main Result

Proof of Theorem 1.3. Let A be as in definition 2.17. A is compact and continuous and has bounded fixed points (if they were to exist) by Lemma 2.28, Lemma 2.18 and Lemma 2.22. By Schaefer’s theorem (Theorem 2.10), A has a fixed point. Thus, a solution pair $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$ to the stationary Mean Field Game system exists. $\square$

2.3 Swapping Composition Order of T and S

In hindsight, we can see that the composition of T (definition 2.16) followed by S (definition 2.15) was convenient because it provided a natural bound for the fixed point. We relied on the fact that the image of S is always bounded (remark 2.21) independently of m . An equally valid construction for A however, is the reverse composition

$$A := T \circ S.$$

Here too the existence of a fixed point proves Theorem 1.3 yet crucially, recall that we have not shown the image of T to have any constant bound in $H_0^1(\Omega)$ rather it depends on the boundedness of its arguments. We note that bounded sets under S remain bounded so A retains its continuity and compactness.⁹ Therefore, we need only demonstrate boundedness of the fixed point set for us to use Schaefer’s Theorem. Under this new construction consider

$$\{u \in H_0^1(\Omega) : u = \mu A[u] \text{ for some } \mu \in [0, 1]\}.$$

The corresponding system, with $A[u]$ substituted for u/μ ¹⁰, is

$$-\frac{\Delta u}{\mu} + H(x, \frac{\nabla u}{\mu}) + \frac{\lambda u}{\mu} = f(x, m), \quad -\Delta m - \operatorname{div}(m \frac{\partial H}{\partial p}(x, \nabla u)) + \lambda m = G(x).$$

We obtain the weak form of HJB with $v = u$

$$\begin{aligned} \int_{\Omega} |\nabla u|^2 + \lambda |u|^2 dx &= \mu \int_{\Omega} f u - H(x, \frac{\nabla u}{\mu}) u dx \\ &\leq \int_{\Omega} |f u| + \mu |H(x, \frac{\nabla u}{\mu}) u| dx \quad \text{since } 0 < \mu \leq 1 \end{aligned}$$

⁹Maps S and T themselves are unchanged. We have already shown that S is continuous (Proposition 2.20) and that T is continuous and compact (Propositions 2.19 and 2.27).

¹⁰See 6

$$\begin{aligned}
&\leq \int_{\Omega} |fu| + \mu C(|\frac{\nabla u}{\mu}| + 1)|u| dx \quad (\text{Assumption 2}) \\
&= \int_{\Omega} |fu| + C(|\nabla u| + 1)|u| dx.
\end{aligned}$$

This is the same form as the inequality in the proof of Theorem 2.11. There we arrived at the following bound

$$\alpha \|u\|_{H_0^1(\Omega)} \leq \frac{1}{\varepsilon\sqrt{2}} \|f(x, m)\|_{L^2(\Omega)} + K$$

with α , ε and K independent (positive) constants. The bound of u must be independent of m so we use the growth condition on f .

Proposition 2.29. *If $A = T \circ S$, $\{u \in H_0^1(\Omega) : u = \mu A[u] \text{ for some } \mu \in [0, 1]\}$ is bounded.*

Proof. We have

$$\begin{aligned}
\alpha \|u\|_{H_0^1(\Omega)} &\leq \frac{1}{\varepsilon\sqrt{2}} \|f(x, m)\|_{L^2(\Omega)} + K \\
&\leq \frac{1}{\varepsilon\sqrt{2}} (\|m\|_{L^2(\Omega)} + |\Omega|^{\frac{1}{2}}) + K \\
&\leq \frac{1}{\varepsilon\sqrt{2}} (\|m\|_{H_0^1(\Omega)} + |\Omega|^{\frac{1}{2}}) + K \\
&\leq \frac{1}{\varepsilon\sqrt{2}} \left(\frac{1}{c_0} \|G\|_{L^2(\Omega)} + |\Omega|^{\frac{1}{2}} \right) + K
\end{aligned}$$

by the Lax-Milgram bound (Theorem 2.5) since $m \in \text{Im}(S)$. Thus, the set is bounded. $\square$

Remark 2.30. Notice the similarity in the proofs of Proposition 2.27 and Proposition 2.29. They both illustrate that the image of T is bounded on a bounded domain. The difference, when we consider the fixed point of the map as a whole, arises in the domains of T . In the first composition T is acting on all $m \in H_0^1(\Omega)$ whereas in the latter it acts on the image of S .

We summarise the analysis of this subsection with a lemma.

Lemma 2.31. *Let $A = T \circ S$ and suppose that $\lambda > 0$ as it appears in equation (1.1) is sufficiently large. Then, A has a fixed point.*

Proof. A is compact and continuous since T is compact and continuous (Proposition 2.27, Proposition 2.19) and S is bounded continuous (Proposition 2.20). The set of fixed points is bounded by Proposition 2.29. Thus, there exists a fixed point by Schaefer's Theorem. $\square$

3 Extension to Non-linear KFP

In certain applications it is natural to allow for the entry or exit of agents from the system. This can be represented by introducing a source term $G(x, m)$ in place of $G(x)$ in equation (1.1). Such terms may model, for example, external recruitment of agents into

the population, departures from the system, or density-dependent entry and exit effects. In this section we consider a class of nonlinear functions $G(x, m)$ satisfying the structural conditions stated in Assumption 7. Our new system of equations reads

$$\begin{aligned} -\Delta u + H(x, \nabla u) + \lambda u &= f(x, m) \quad \text{in } \Omega, \\ -\Delta m - \operatorname{div}\left(m \frac{\partial H}{\partial p}(x, \nabla u)\right) + \lambda m &= G(x, m) \quad \text{in } \Omega. \end{aligned} \tag{3.1}$$

In this scenario we cannot apply the Lax-Milgram Theorem to the KFP equation. We will therefore adopt fixed point theorems to find a solution but as with $H(x, \nabla u)$ we impose assumptions on $G(x, m)$.

7. $G(x, m) : \Omega \times L^2(\Omega) \rightarrow L^2(\Omega)$ is

- Lipschitz continuous with respect to m with Lipschitz constant $\tilde{B}$,
- bounded by $B(|m| + 1)$ for a given constant B , for all $x \in \Omega$ and $m \in L^2(\Omega)$.

8. In *addition* to our previous constraint on λ , we have $\lambda > \max(B, \tilde{B}) + \frac{L^2}{4}$

Example 3.1. For a source term G satisfying Assumption 7, one may consider

$$G(x, m) = a(x) \tanh(m),$$

where $a \in L^\infty(\Omega)$. From a modelling perspective, this choice corresponds to density-dependent entry or exit with saturation: for small m the response is approximately linear, while for large m the source term remains bounded.

Theorem 3.2 (Existence of Non-linear KFP Solution). *If $\lambda > 0$ is sufficiently large, there exists $m \in H^2(\Omega) \cap H_0^1(\Omega)$ solving KFP with fixed u .*

Proof. We design the map whose fixed point is a solution to the non-linear equation.

$$A : H_0^1(\Omega) \rightarrow H_0^1(\Omega), \quad A[m] = w \quad \text{where} \quad -\Delta w - \operatorname{div}\left(w \frac{\partial H}{\partial p}(x, \nabla u)\right) + \lambda w = G(x, m).$$

We know w exists by Lax-Milgram Theorem 2.5, so A is well defined. Continuity and compactness are the same as in Theorem 2.11 so we omit them. The only difference is in the threshold of λ which we'll now derive. As in Theorem 2.11 we want to show

$$\{m \in H_0^1(\Omega) : m = \mu A[m] \text{ for some } \mu \in [0, 1]\}$$

is bounded. Substituting $w = m/\mu$ ¹¹ and taking $v = m$, we obtain the weak form

$$\int_{\Omega} |\nabla m|^2 + m \nabla m \frac{\partial H}{\partial p} + \lambda |m|^2 dx = \mu \int_{\Omega} G(x, m) m dx$$

¹¹See 6

Moving the $\frac{\partial H}{\partial p}$ term over and Assumption 3 give

$$\begin{aligned} \int_{\Omega} |\nabla m|^2 + \lambda |m|^2 dx &\leq \int_{\Omega} |G(x, m)| |m| + L |m| |\nabla m| dx && \text{since } 0 < \mu \leq 1 \\ &\leq \int_{\Omega} B(|m| + 1) |m| + L |m| |\nabla m| dx && (\text{Assumption 7}) \end{aligned}$$

To isolate $|\nabla m|^2$ and m^2 terms on the RHS, we apply Young's inequality 2.4 to give

$$\int_{\Omega} |\nabla m|^2 + \lambda |m|^2 dx \leq \int_{\Omega} \frac{1}{2} \delta^2 |\nabla m|^2 + \left(B + \frac{1}{2} \varepsilon^2 + \frac{L^2}{2\delta^2}\right) |m|^2 dx + K^2 \quad \varepsilon, \delta \in (0, \sqrt{2})$$

where K^2 is the sum of terms independent of $m, \nabla m$. If $\lambda > B + \frac{L^2}{4}$ then we can find suitable ε, δ such that $\|m\|_{H_0^1(\Omega)}$ is bounded.¹² After taking the difference of both sides, the inequality simplifies to

$$\tilde{\alpha} \|m\|_{H_0^1(\Omega)} \leq K$$

where $\tilde{\alpha}$ is the minimum of both factors of m^2 and ∇m^2 . □

Remark 3.3. This result has a very important implication. It tells us that any solution $m \in H_0^1(\Omega)$ is still bounded by a constant despite $G(\cdot)$ being non-linear.

Proposition 3.4. *If $\lambda > 0$ is sufficiently large, then the fixed point m is unique.*

Proof. Suppose m_1, m_2 both solve the non-linear KFP equation. Taking the difference and setting $v = m_1 - m_2$ gives

$$\begin{aligned} \|\nabla(m_1 - m_2)\|_{L^2}^2 + \lambda \|m_1 - m_2\|_{L^2}^2 &\leq \int_{\Omega} \left| \frac{\partial H}{\partial p} \right| |m_1 - m_2| |\nabla(m_1 - m_2)| + |G_1 - G_2| |m_1 - m_2| dx \\ &\leq L \|m_1 - m_2\|_{L^2} \|\nabla(m_1 - m_2)\|_{L^2} + \tilde{B} \|m_1 - m_2\|_{L^2}^2 \end{aligned}$$

using Assumptions 3, 7 and Cauchy–Schwarz. Hence, by our previous use of Young's inequality in Proposition 2.7,

$$\lambda - \tilde{B} > \frac{L^2}{4} \implies m_1 = m_2 \text{ in } H_0^1(\Omega).$$

□

¹²See Theorem 2.11 for further elaboration on this point.

As with the linear version of the KFP equation we define a map $\tilde{S}$ to take the place of S in definition 2.15.

Definition 3.5. Define $\tilde{S} : H_0^1(\Omega) \rightarrow H_0^1(\Omega)$ by $\tilde{S}(u) = m$, where m solves non-linear KFP for given u .

Since the form of HJB in equation (3.1) is the same as that in equation (1.1), we can investigate the existence of a fixed point of the map $\tilde{A}$.

Definition 3.6. Let T and $\tilde{S}$ be as in definitions 2.15 and 2.16. Define

$$\tilde{A} : H_0^1(\Omega) \rightarrow H_0^1(\Omega), \quad A := \tilde{S} \circ T.$$

A fixed point of $\tilde{A}$ means a solution to the new system of equations exists.

Main Result for Non-linear KFP

Theorem 3.7 (Existence of MFG Solution with Non-linear KFP). *Let the Mean Field Game system be characterised by equation (3.1) and suppose the assumptions above are true. Then a solution $(u, m) \in H_0^1(\Omega) \times H_0^1(\Omega)$ exists.*

Proof. We prove the theorem via showing $\tilde{A}$ has a fixed point. Once again, we wish to use Schaefer's theorem. We need to show that $\tilde{S}$ is continuous. Let (u_k) converge to $u \in H_0^1(\Omega)$. Denote $m = \tilde{S}[u]$, $m_k = \tilde{S}[u_k]$, $b = \frac{\partial H}{\partial p}(x, \nabla u)$, $b_k = \frac{\partial H}{\partial p}(x, \nabla u_k)$ and $G_k = G(x, m_k)$. Taking the weak formulation of the difference in KFP and setting $v = m - m_k$ yields

$$\|\nabla(m - m_k)\|_{L^2}^2 + \lambda \|m - m_k\|_{L^2}^2 \leq \int_{\Omega} |\nabla(m - m_k)| |m b - m_k b_k| + |G - G_k| |m - m_k| dx.$$

Applying the Lipschitz condition (Assumption 7) to G, G_k gives

$$\|\nabla(m - m_k)\|_{L^2}^2 + \lambda \|m - m_k\|_{L^2}^2 \leq \int_{\Omega} |\nabla(m - m_k)| |m b - m_k b_k| + \tilde{B} |m - m_k|^2 dx.$$

By λ being sufficiently large to account for the $\tilde{B}$ term (Assumption 8) and recalling the same line of reasoning in Proposition 2.20 we conclude that $\tilde{S}$ is continuous in $H_0^1(\Omega)$. We also have that $\tilde{S}$ is bounded (Theorem 3.2) and that T is both compact in $H_0^1(\Omega)$ (Proposition 2.27) and continuous in $H_0^1(\Omega)$ (Proposition 2.19). $\tilde{A}$ is therefore compact in $H_0^1(\Omega)$.

To meet the final condition of Schaefer's theorem, we must show

$$\{m \in H_0^1(\Omega) : m = \mu \tilde{A}[m] \text{ for some } \mu \in [0, 1]\}$$

is bounded. The corresponding system, with $\tilde{A}[m]$ substituted for m/μ ¹³, is

$$-\Delta u + H(x, \nabla u) + \lambda u = f(x, m), \quad \frac{1}{\mu} \left(-\Delta m - \operatorname{div} \left(m \frac{\partial H}{\partial p}(x, \nabla u) \right) + \lambda m \right) = G(x, \frac{m}{\mu}).$$

¹³See 6

Taking the weak form of the latter and setting $v = m$ implies

$$\begin{aligned} \int_{\Omega} |\nabla m|^2 + \lambda |m|^2 dx &\leq \int_{\Omega} \mu |G(x, \frac{m}{\mu})| |m| + L |m| |\nabla m| dx && \text{See Theorem 3.2} \\ &\leq \int_{\Omega} B(|m| + 1) |m| + L |m| |\nabla m| dx && (\text{Assumption 7}) \end{aligned}$$

In Theorem 3.2 we showed that when $\lambda > 0$ is sufficiently large, the inequality simplifies to

$$\tilde{\alpha} \|m\|_{H_0^1(\Omega)} \leq K$$

Thus, the set

$$\{m \in H_0^1(\Omega) : m = \mu \tilde{A}[m] \text{ for some } \mu \in [0, 1]\}$$

is bounded independently of μ . By Schaefer's theorem (2.10), $\tilde{A}$ has a fixed point. $\square$

4 Conclusion

We briefly summarise the strategy used to establish the existence of steady state solutions to the coupled system. Our approach proceeds by analysing the two equations separately and then combining the resulting solution operators. For a fixed density $m \in H_0^1(\Omega)$, the Hamilton–Jacobi–Bellman equation is solved to obtain $u \in H_0^1(\Omega)$, while for a fixed $u \in H_0^1(\Omega)$ the Kolmogorov–Fokker–Planck equation determines the corresponding density $m \in H_0^1(\Omega)$. These steps define two operators between the relevant function spaces. By composing these operators we obtain a map whose fixed points correspond to solution pairs (u, m) of the coupled system.

The fixed point framework therefore provides a flexible mechanism for passing from solvability of the individual equations to the existence of a solution for the full mean field game system. Finally, we further extend this analysis by incorporating nonlinear source terms $G(x, m)$ in the KFP equation and show that, under suitable structural assumptions, the same argument yields existence of steady state solutions in this more general setting.

References

- [1] Robert J. Aumann. Markets with a continuum of traders. *Econometrica*, 32(1-2), 1964.
- [2] Pierre Cardaliaguet. Notes on mean field games. 2013.
- [3] Lawrence C. Evans. *Partial Differential Equations*. American Mathematical Society, 2 edition, 2010.
- [4] Pierre-Louis Lions Jean-Michel Lasry. Mean field games. *Japanese Journal of Mathematics*, 2007.

- [5] Peter D. Lax and Arthur N. Milgram. Parabolic equations. *Contributions to the Theory of Partial Differential Equations*, 1954.