

Calculus Notes

Avid David

July 3, 2025

Contents

I	Integral Calculus in One Variable	2
1	Indefinite Integrals	2
1.1	Integral of a function	2
1.2	Ways to calculate indefinite integrals	2
1.3	Integral of rational functions	4
1.4	Integral of trigonometric functions	6

Part I

Integral Calculus in One Variable

1 Indefinite Integrals

1.1 Integral of a function

The integral is the inverse of the derivative operator. Given a function $f(x)$, we can find a function $F(x)$ such that $F'(x) = f(x)$. The integral of a function is not unique, as for each $f(x)$, $F(x) + C$ is also an integral of that function.

Concept

Both the derivative and the integral are linear operators, so we have:

$$\int [\alpha f(x) + \beta g(x)] dx = \alpha \int f(x) dx + \beta \int g(x) dx$$

1.2 Ways to calculate indefinite integrals

1. Expansion

We use the linearity rule to turn a complicated integral into the sum of many simpler ones, then calculate one by one.

2. Changing the differential expression (u -substitution)

If $\int f(x) dx = F(x) + C$ then $\int f(u) dx = F(u) + C$, where $u = u(x)$ is a continuously differentiable function. Then, we can change the integrand $g(x) dx$ into:

$$g(x) = f(u(x))u'(x)dx$$

Then the integral turns into:

$$\int g(x) dx = \int f(u(x))u'(x) dx = \int f(u(x)) du = F(u) + C$$

Insight

In the simple case $u = ax + b$, we have $du = a dx$. Then if $\int f(x) dx = F(x) + C$ then:

$$\int f(ax + b) dx = \frac{1}{a} F(ax + b) + C$$

3. **Change of variables** Consider the integral $I = \int f(x)dx$, where $f(x)$ is a continuous function. We can change $f(x)$ such that we work with functions with known or easier antiderivatives:

(a) **Change of variables type 1:**

Let $x = \varphi(t)$, where $\varphi(t)$ is a monotonic and continuously differentiable function. Then:

$$I = \int f(x)dx = \int f[\varphi(t)]\varphi'(t)dt$$

Denote the antiderivative of $g(t) = f[\varphi(t)]\varphi'(t)$ as $G(t)$ and $h(x)$ as the inverse of $x = \varphi(t)$, we then have:

$$\int g(t)dt = G(t) + C \Rightarrow I = G[h(x)] + C$$

(b) **Change of variables type 2:**

Let $t = \psi(x)$, where $\psi(x)$ is a continuously differentiable function and we can write $f(x) = g[\psi(x)]\psi'(x)$. Then:

$$I = \int f(x)dx = \int g[\psi(x)]\psi'(x)dx$$

Denote the antiderivative of $g(t)$ as $G(t)$, then:

$$I = G[\psi(x)] + C$$

Important

Remember to change back to the original variable!

4. **Integration by parts** Let $u = u(x)$ and $v = v(x)$ be continuously differentiable functions. We know:

$$d(uv) = u dv + v du \Rightarrow \int d(uv) = \int u dv + v du$$

Then we have the following formula:

$$\int u dv = uv - \int v du$$

Consider the integral $I = \int f(x)dx$ We need to express:

$$f(x)dx = [g(x)h(x)]dx = g(x)[h(x)dx] = u dv$$

then apply the integration by parts formula to $u = g(x)$, $v = h(x)dx$

1.3 Integral of rational functions

A rational function is one with the form $f(x) = \frac{P(x)}{Q(x)}$, where $P(x)$ and $Q(x)$ are polynomials in x . If $\deg P(x) < \deg Q(x)$ then it's called a *true rational function*.

Using polynomial division, we can rewrite any rational function:

$$f(x) = H(x) + \frac{r(x)}{Q(x)}$$

The integral of $H(x)$ can easily be computed. As for the true rational function $\frac{r(x)}{Q(x)}$, we will use partial fractions to decompose it into four simpler types of functions. First, using the method of Undetermined Coefficients, we can write $Q(x)$ as the product of linear polynomials and quadratic polynomials with no real roots:

$$Q(x) = \prod_{k=1}^n (x - a_k)^{\alpha_k} \prod_{k=1}^n (x^2 + p_k x + q_k)^{\beta_k}$$

Note that this can always be achieved as per the Fundamental Theorem of Algebra. Then, we just need to compute the following four types of integrals:

Insight

1. $\int \frac{A}{x - a} dx$
2. $\int \frac{A}{(x - a)^k} dx \quad (k \geq 2)$
3. $\int \frac{Mx + N}{x^2 + px + q} dx$
4. $\int \frac{Mx + N}{(x^2 + px + q)^m} dx \quad (m \geq 2)$

These integrals cover all cases because any partial fraction decomposition over \mathbb{R} involves repeated linear factors and irreducible quadratic factors, possibly raised to powers. The first two ones are simple:

1. $\int \frac{A}{x - a} dx = A \ln |x - a| + C$
2. $\int \frac{A}{(x - a)^k} dx = A \int (x - a)^{-k} = \frac{-A}{(k - 1)(x - a)^{k-1}} \quad (k \geq 2)$

The third one is a bit more complicated: we need to use a clever substitution, inspired by completing the square, to bring it back to familiar forms:

Let $a = \sqrt{q - \frac{p^2}{4}}$ and $t = x + \frac{p}{2}$, the integral becomes:

$$\begin{aligned}\int \frac{Mx + N}{x^2 + px + q} dx &= \int \frac{Mt + (N - \frac{Mp}{2})}{t^2 + a^2} dt \\ &= M \int \frac{t}{t^2 + a^2} dt + \left(N - \frac{Mp}{2}\right) \int \frac{1}{t^2 + a^2} dt\end{aligned}$$

For the first integral, substitute in $u = t^2$ and we have a logarithm. For the second one, divide both the numerator and the denominator by a^2 then substitute $u = \frac{t}{a}$, we yield the inverse tangent. Then the integral is simply:

$$\int \frac{Mx + N}{x^2 + px + q} dx = \frac{M}{2} \ln(x^2 + px + q) + \frac{2N - Mp}{\sqrt{4q - p^2}} \arctan \frac{2x + p}{\sqrt{4q - p^2}} + C$$

Before moving on to solve the last integral, let's review a bit of complex numbers:

Concept

Some important (and beautiful) formulas::

Euler's Formula: $e^{ix} = \cos x + i \sin x$

De Moivre's Formula: $(\cos x + i \sin x)^n = \cos nx + i \sin nx$

We will prove it using ODEs. Let $f(x) = e^x$ and $g(x) = \cos x + i \sin x$. We know:

$$f'(x) = ie^{ix} = if(x)$$

$$g'(x) = -\sin x + i \cos x = ig(x)$$

And also the initial values:

$$f(0) = g(0) = 1$$

By uniqueness of solutions to first-order linear ODEs, Euler's formula is proven. By substituting $f(nx) = g(nx)$, De Moivre's formula is proven. Then we have:

Insight

- $\cos x = \Re(e^{ix}) = \frac{e^{ix} + e^{-ix}}{2}$

- $\sin x = \Im(e^{ix}) = \frac{e^{ix} - e^{-ix}}{2i}$

From this, we can derive a general formula for powers of the sine and cosine:

$$\cos^n x = \left(\frac{e^{ix} + e^{-ix}}{2} \right)^n = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} e^{i(n-2k)x}$$

Now if we group k and $n - k$ in pairs to get $\cos(n - 2k)$, we get:

$$\cos^n x = \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos((n - 2k)x)$$

Analogously, we also have:

$$\sin^n x = \frac{1}{(2i)^n} \sum_{k=0}^n \binom{n}{k} (-1)^k \cos\left(\frac{\pi n}{2} - (n - 2k)x\right)$$

Now, back to the fourth integral, we first do the same thing: change variables $a = \sqrt{q - \frac{p^2}{4}}$ and $t = x + \frac{p}{2}$. This yields:

$$\begin{aligned} \int \frac{Mx + N}{(x^2 + px + q)^m} dt &= \int \frac{Mt + (N - \frac{Mp}{2})}{(t^2 + a^2)^m} dt \\ &= M \int \frac{t}{(t^2 + a^2)^m} dt + \left(N - \frac{Mp}{2}\right) \int \frac{1}{(t^2 + a^2)^m} dt \end{aligned}$$

Again, to compute the first integral, we use the substitution $u = t^2$. Then it is:

$$M \int \frac{t}{(t^2 + a^2)^m} dt = \frac{-M}{2(m-1)(t^2 + a^2)^{m-1}} + C$$

For the second integral however, this time we need a different substitution. Let $t = a \tan z$. Then: $t^2 + a^2 = a^2 \sec^2 z$, and $dt = a \sec^2 z dz$. Then the integral turns into:

$$\int \frac{1}{(t^2 + a^2)^m} dt = a^{2m-1} \int \cos^{2m-2} z dz$$

Let $I(z) = \int \cos^{2m-2} z dz$ and now, we can take advantage of the linearity rule and the formula for powers of cosine we just derived. The integral turns into:

$$\begin{aligned} I(z) &= \int \frac{1}{2^{2m-2}} \sum_{k=0}^{2m-2} \binom{2m-2}{k} \cos(2(m-k-1)z) dz \\ &= \frac{1}{2^{2m-2}} \sum_{k=0}^{2m-2} \binom{2m-2}{k} \frac{1}{2m-2-2k} \sin(2(m-k-1)z) + C \\ &= \frac{1}{2^{2m-2}} \left[\binom{2m-2}{m-1} z + \sum_{k=0, k \neq m-1}^{2m-2} \binom{2m-2}{k} \frac{1}{2m-2-2k} \sin(2(m-k-1)z) \right] + C \end{aligned}$$

Then the integral, as a whole, evaluates to:

$$\int \frac{Mx + N}{(x^2 + px + q)^m} dx = \frac{-M}{2(m-1)(x^2 + px + q)^{m-1}} + \left(N - \frac{Mp}{2}\right) \left(\sqrt{q - \frac{p^2}{4}}\right)^{2m-1} \cdot I(z) + C$$

1.4 Integral of trigonometric functions