

9.A.1 BIG O NOTATION – EXAMPLES

Example 1

Let $f(n) = 3n^3$ for an algorithm. Prove that $f(n)$ of the algorithm is in $O(n^3)$.

Solution

The definition of the Big -Oh notation is that $f(n) \leq c \times g(n)$.

In order to prove that we know:

$$f(n) \in O(g(n))$$

$$\text{or, } f(n) = O(g(n))$$

Where $g(n)$ is in n^3 .

Hence, we can show that:

$3n^3 \leq c \times n^3$, holds good for a positive number c and sufficiently large values of n .

$f(n) = 3n^3$ can also be written as:

$$f(n) = 3n^3 + 0$$

We can write it as:

$$3n^3 + 0 \leq 3n^3 + n^3$$

$$\approx 3n^3 + 0 \leq 4n^3 \text{ [Note } 3n^3 = 3n^3 \text{ but } 3n^3 \text{ is always } \leq 4n^3]$$

[As highest degree of polynomial n is 3]

$$\approx 0 \leq 4n^3 - 3n^3$$

$$\approx 0 \leq n^3$$

Or, Divide n^2 in both side:

$$\approx \frac{0}{n^2} \leq \frac{n^3}{n^2}$$

$$\approx 0 \leq n$$

$$\text{or, } n \geq 0$$

Therefore $n_0 = 0$.

We can again write it as:

$$3n^3 + 0 \leq 4 \times n^3$$

Hence $c \geq 4$.

Therefore, $f = O(g)$

or in other words, the algorithm is $O(n^3)$.

Hence proved.

Example 2

Let $f(n) = 3n + 8$ for an algorithm. Prove that $f(n)$ of the algorithm is in $O(n)$.

Solution

The definition of the Big -Oh notation is that $f(n) \leq c \times g(n)$.

In order to prove that we know:

$$f(n) \in O(g(n))$$

$$\text{or, } f(n) = O(g(n))$$

Where $g(n)$ is in n .

Hence, we can show that:

$3n + 8 \leq c \times n$, holds good for a positive number c and sufficiently large values of n .

$$3n + 8 \leq 3n + n$$

$$\approx 3n + 8 \leq 4n \text{ [} 3n \text{ is always } \leq 4n \text{]}$$

[As highest degree of polynomial n is 1]

$$\therefore 3n + 8 \leq 4n$$

$$\approx 8 \leq 4n - 3n$$

$$\approx 8 \leq n$$

$$\text{or, } n \geq 8$$

$\therefore n_0 = 8$

$$3n + 8 \leq 4n$$

Can be written as:

$$3n + 8 \leq 4 \times n$$

$$\text{Hence, } c \geq 4.$$

Therefore, f is $O(g)$

or in other words $O(n)$.

Hence proved.

Example 3

Let $f(n) = n^2 + 1$ for an algorithm. Prove that $f(n)$ of the algorithm is in $O(n^2)$.

Solution

The definition of the Big -Oh notation is that $f(n) \leq c \times g(n)$.

In order to prove that we know:

$$f(n) \in O(g(n))$$

$$\text{or, } f(n) = O(g(n))$$

Where $g(n)$ is in n^2 .

Hence, we can show that:

$n^2 + 1 \leq c \times n^2$, holds good for a positive number c and sufficiently large values of n .

$$n^2 + 1 \leq n^2 + n^2$$

$$\approx n^2 + 1 \leq 2n^2 [n^2 \text{ is always } \leq 2n^2]$$

[As highest degree of polynomial n is 2]

$$\therefore n^2 + 1 \leq 2n^2$$

$$\approx 1 \leq 2n^2 - n^2$$

$$\approx 1 \leq n^2$$

$$\approx -n^2 + 1 \leq 0$$

$$\approx -(n^2 + 1) \leq 0$$

$$\approx -(n^2 - (-1)^2) \leq 0$$

$$\text{As we know: } x^2 - y^2 = (x + y)(x - y)$$

$$\approx -1 \times ((n + (-1)) (n - (-1))) \leq 0$$

$$\approx -1 \times ((n - 1) (n + 1)) \leq 0$$

$$\approx (n - 1)(n + 1) \leq \frac{0}{-1}$$

$$\approx (n - 1)(n + 1) \leq 0$$

[From Quadratic Inequalities]

Say $n = 1$

$$\approx (1 - 1) \times (1 + 1) \leq 0$$

$$\approx 0 \times 2 \leq 0$$

$$\approx 0 \leq 0[True]$$

Also,

$$\approx 0 \geq 0[True]$$

Say $n = 2$

$$\approx (2 - 1) \times (2 + 1) \leq 0$$

$$\approx 1 \times 3 \leq 0$$

$$\approx 3 \leq 0[False]$$

But,

$$\approx 3 \geq 0[True]$$

Hence, we can say $n \geq 1$

Say $n = 0$

$$\approx (0 - 1) \times (0 + 1) \leq 0$$

$$\approx -1 \times 1 \leq 0$$

$$\approx -1 \leq 0[True]$$

Say $n = -1$


$$\approx (-1 - 1) \times (-1 + 1) \leq 0$$

$$\approx -2 \times 0 \leq 0$$

$$\approx 0 \leq 0[True]$$

That implies $n \leq -1$

$$\therefore -1 \geq n \geq 1$$



$(-\infty, -1] \cup [1, \infty)$ and they are in Sem – open and semi closed intervals.

Hence now we get:

$$n^4 + 1 \leq 2n^2 \text{ for all } n \geq 1$$

As from the definition of Big - O notation, all should be positive numbers but not negative numbers.

Therefore , we get $n_0 = 1$.

And,

$$n^4 + 1 \leq 2 \times n^2 \text{ in terms of } f(n) \leq c \times g(n)$$

Hence:

$$c \geq 2 .$$

Therefore, f is $O(g)$

or in other words $O(n^2)$.

Hence proved.

Example 4

Let $f(n) = n^4 + 100n^2 + 50$ for an algorithm. Prove that $f(n)$ of the algorithm is in $O(n^4)$.

Solution

The definition of the Big -Oh notation is that $f(n) \leq c \times g(n)$.

In order to prove that we know:

$$f(n) \in O(g(n))$$

$$\text{or, } f(n) = O(g(n))$$

Where $g(n)$ is in n^4 .

Hence, we can show that:

$$\begin{aligned}
n^4 + 100n^2 + 50 &\leq c \times n^4 \\
&\approx n^4 + 100n^2 + 50 \leq n^4 + n^4 \\
&\approx n^4 + 100n^2 + 50 \leq 2n^4 \text{ [} n^4 \text{ is always } \leq 2n^4 \text{]} \\
&\text{[As highest degree of polynomial } n \text{ is 4]}
\end{aligned}$$

Now,

$$\begin{aligned}
&\approx -2n^4 + n^4 + 100n^2 + 50 \leq 0 \\
&\approx -n^4 + 100n^2 + 50 \leq 0
\end{aligned}$$

We can write the above equation as:

$$\approx -n^4 + 100n^2 + 50 = 0 \text{ and } -n^4 + 100n^2 + 50 < 0$$

Taking the equation:

$$\approx -n^4 + 100n^2 + 50 = 0$$

Rewriting the equation, $u = n^2$ and $u^2 = n^4$:

$$\approx -u^2 + 100u + 50 = 0$$

Solving with quadratic equation formula:

Quadratic Equation of the form $ax^2 + bx + c = 0$:

$$x_{1,2} = \frac{(-b \pm \sqrt{b^2 - 4ac})}{2a}$$

For $a = -1$, $b = 100$, $c = 50$

$$u_{1,2} = \frac{(-100 \pm \sqrt{100^2 - 4(-1)(50)})}{2(-1)}$$

$$u_{1,2} = \frac{(-100 \pm \sqrt{100^2 - 4(-1)(50)})}{-2}$$

$$u_{1,2} = \frac{(-100 \pm \sqrt{100^2 + 200})}{-2}$$

$$u_{1,2} = \frac{(-100 \pm \sqrt{10000 + 200})}{-2}$$

$$u_{1,2} = \frac{(-100 \pm \sqrt{10200})}{-2}$$

Using prime factorization of $10200 = 2^3 \times 3 \times 5^2 \times 17$

$$u_{1,2} = \frac{(-100 \pm \sqrt{2^3 \times 3 \times 5^2 \times 17})}{-2}$$

Applying exponent rule: $a^{b+c} = a^b \times a^c$

$$u_{1,2} = \frac{(-100 \pm \sqrt{2^2 \times 2 \times 3 \times 5^2 \times 17})}{-2}$$

Applying radical rule: $\sqrt[n]{ab} = \sqrt[n]{a} \times \sqrt[n]{b}$

$$u_{1,2} = \frac{(-100 \pm \sqrt{2^2} \times \sqrt{5^2} \times \sqrt{2 \times 3 \times 17})}{-2}$$

Applying radical rule: $\sqrt[n]{a^n} = a$

$$u_{1,2} = \frac{(-100 \pm 2 \times 5 \times \sqrt{2 \times 3 \times 17})}{-2}$$

$$u_{1,2} = \frac{(-100 \pm 10 \times \sqrt{2 \times 3 \times 17})}{-2}$$

$$u_{1,2} = \frac{(-100 \pm 10 \sqrt{102})}{-2}$$

$$\begin{aligned} u &= \frac{(-100 + 10 \sqrt{102})}{-2} \\ &= \frac{10(-10 + \sqrt{102})}{-2} \\ &= -5(-10 + \sqrt{102}) - -i \end{aligned}$$

$$\begin{aligned} u &= \frac{(-100 - 10 \sqrt{102})}{-2} \\ &= \frac{-10(10 + \sqrt{102})}{-2} \\ &= 5(10 + \sqrt{102}) - -ii \end{aligned}$$

Substituting back $u = n^2$ and solving for n,

$$n^2 = -5(-10 + \sqrt{102})$$

$(g(x))^2$ cannot be negative for $x \in R$, hence no solution.

$$n^2 = 5(10 + \sqrt{102})$$

We know $(g(x))^2 = f(a)$ the solutions are $\sqrt{f(a)}$, $-\sqrt{f(a)}$

$$\therefore n = \sqrt{5(10 + \sqrt{102})} \quad \text{and} \quad n = -\sqrt{5(10 + \sqrt{102})}$$

Now we can easily understand,

$$\approx -n^4 + 100n^2 + 50 = 0$$

putting $n = -\sqrt{5(10 + \sqrt{102})}$ in the above equation we will get a negative value while putting $n = \sqrt{5(10 + \sqrt{102})}$ in the above equation we will get a positive value,

Hence:

$$n \leq -\sqrt{5(10 + \sqrt{102})}$$

or

$$n \geq \sqrt{5(10 + \sqrt{102})}$$

Now what does $\sqrt{5(10 + \sqrt{102})}$ stand for : 10.028484537

As by definition: the function f and g should be set of natural numbers and it should grow by time (growth rate) we take $n \geq 11$.

Or by analysis:

$$\approx -n^4 + 100n^2 + 50 \leq 0$$

$$\approx 100n^2 + 50 \leq n^4$$

$$\text{or, } n^4 \geq 100n^2 + 50$$

if we take $n = 10$

$$\approx 10^4 \geq 100 \times 10^2 + 50$$

$$\approx 10000 \geq 10000 + 50$$

$$\approx 10000 \geq 10050 [\text{Not True}]$$

if we take $n = 11$

$$\approx 11^4 \geq 100 \times 11^2 + 50$$

$$\approx 14641 \geq 1210 + 50$$

$$\approx 14641 \geq 1260 \text{ [True]}$$

Hence, we confirm in both the ways that:

$$n^4 + 100n^2 + 50 \leq 2n^4 \text{ for all } n \geq 11$$

Therefore, we get $n_0 = 11$.

And,

$$n^4 + 100n^2 + 50 \leq 2 \times n^4 \text{ in terms of } f(n) \leq c \times g(n)$$

Hence:

$$c \geq 2.$$

Therefore, f is $O(g)$

or in other words $O(n^4)$.

Hence proved.

Example 5

Let $f(n) = n$ for an algorithm. Let $g(n) = n$. Prove that $f(n)$ of this algorithm is in $O(n)$.

Solution

$$f(n) \leq c \times g(n)$$

$$\Rightarrow n \leq 1 \times n, \text{ for all } n \geq 1$$

$$\Rightarrow n = O(n), c \geq 1 \text{ and } n_0 = 1$$

Example 6

Let $f(n) = 410$ for an algorithm. Let $g(n) = 410$. Prove that $f(n)$ of this algorithm is in $O(1)$.

Solution

$$f(n) \leq c \times g(n)$$

$$\Rightarrow 410 \leq 1 \times 410, \text{ for all } n \geq 1$$

$$\Rightarrow 410 = O(1), c \geq 1 \text{ and } n_0 = 1$$

Example 7

Find upper bound for $f(n) = 2n^3 - 2n^2$. Prove that $f(n)$ of this algorithm is in $O(n^3)$.

Solution

The definition of Big-Oh notation is that $f(n) \leq c \times g(n)$.

$$\Rightarrow f(n) \leq (2n^3 + 2n^3) - 2n^3 \text{ [As growth rate is doubled also } 4n^3 \geq 2n^3]$$

$$\Rightarrow f(n) \leq (4n^3) - 2n^3$$

$$\Rightarrow f(n) \leq 2n^3$$

And we can say:

$$2n^3 - 2n^2 \leq 2n^3$$

Deducing it to $f(n) \leq c \times g(n)$, we get:

$$\Rightarrow 2n^3 - 2n^2 \leq 2 \times n^3$$

Hence $c = 2$ and $g(n) = n^3$

By Inequality deduction:

$$\Rightarrow -2n^2 \leq 2n^3 - 2n^3$$

$$\Rightarrow -2n^2 \leq 0$$

$$\Rightarrow n^2 \leq 0$$

$$\Rightarrow n \leq 0$$

More specifically:

N	$2n^3 - 2n^2$	$2n^3$
0	0	0
1	0	1
2	8	16

Hence it starts from 1, hence $n \geq 1$

Therefore $n_0 = 1$.

Hence $f = O(g)$

or $f(n) = O(n^3)$

NO UNIQUENESS in Above Method

There is no unique set of values for n_0 and c in proving the asymptotic bounds.

Let us consider, $100n + 5 = O(n)$. For this function there are multiple n_0 and c values possible.

Solution1:

$$100n+5 \leq 100n+n$$

$\approx 100n+5 \leq 101n$, for all $n \geq 5$, $n_0 = 5$ and $c \geq 101$ is a solution.

Solution2:

$$100n+5 \leq 100n+5n$$

$\approx 100n+5 \leq 105n$, for all $n \geq 1$, $n_0 = 1$ and $c \geq 105$ is also a solution.

Example 8

Let $f(n) = 3n^3 + 2n^2 + 3$ for an algorithm. Let $g(n) = n^3$. Prove that $f(n)$ of this algorithm is in $O(n^3)$.

Solution

The definition of Big-Oh notation is that $f(n) \leq c \times g(n)$. Therefore, one must show that $3n^3 + 2n^2 + 3 \leq cn^3$ holds good for a positive number c and for sufficiently large values of n .

$$f(n) = 3n^3 + 2n^2 + 3$$

$$f(n) \leq 3n^3 + 2n^3 + 3 \text{ (as growth of functions } n^2 \text{ to } n^3)$$

$$f(n) \leq 3n^3 + 2n^3 + 3n^3 \text{ (3 is less than } n^3)$$

$$f(n) \leq 8n^3$$

It can be observed that $c = 3 + 2 + 3 = 8$ (one can approximate $2n^2$ and 3 to $2n^3$ and $3n^3$ respectively). This condition holds good for any values of $c \geq 8$.

Let the polynomial be

$$f(n) = \sum_{i=0}^m a_i n^i$$

whose degree is m . Then one can show that $f(n) = O(n^m)$.

$$|f(n)| \leq |a_m|n^m + |a_{m-1}|n^{m-1} + \dots + |a_1|n + |a_0|$$

$$\approx |f(n)| \leq |a_m|n^m + |a_{m-1}|n^m + \dots + |a_1|n^m + |a_0|n^m \text{ for all } n \geq 1$$

$$\approx \left(\sum_{i=0}^m |a_i| \right) n^m$$

$$\approx c \times n^m$$

$$\approx O(n^m)$$

Hence the above algorithm has $O(n^3)$.

This is another way we can prove the algorithm has *the complexity*.

Example 9

Let $f(n) = \frac{(2x^3 + 13 \log_2 x)}{7n^2}$ for an algorithm A. Prove that $f(n)$ of algorithm A is $O(n)$.

Solution

It can be observed that $\log x < x$ is always true. Therefore, one can argue that $13 \log_2 x \leq 13x$ and as $13x \leq 13x^3$ always, one can rewrite $f(n)$ as follows:

$$f(n) \leq \frac{2x^3 + 13x^3}{7n^2}$$

$$f(n) \leq \frac{15x^3}{7n^2}$$

$$\cong 2n^{3-2}$$

$$\cong 2n \text{ for all } n > 1$$

$$\therefore f(n) = O(n)$$