

4.1 INTRODUCTION TO VECTOR SPACES

Definition of a Vector Space

In Theorems 1.3 and 1.11, we proved eight properties of addition and scalar multiplication in \mathbb{R}^n and \mathcal{M}_{mn} . These properties are important because all other results involving these operations can be derived from them. We now introduce the general class of sets called **vector spaces**,¹ having operations of addition and scalar multiplication with these same eight properties, as well as two **closure properties**.

Definition A **vector space** is a set \mathcal{V} together with an operation called **vector addition** (a rule for adding two elements of \mathcal{V} to obtain a third element of \mathcal{V}) and another operation called **scalar multiplication** (a rule for multiplying a real number times an element of \mathcal{V} to obtain a second element of \mathcal{V}) on which the following ten properties hold:

For every \mathbf{u}, \mathbf{v} , and \mathbf{w} in \mathcal{V} , and for every a and b in \mathbb{R} ,

- | | |
|--|---|
| (A) $\mathbf{u} + \mathbf{v} \in \mathcal{V}$ | Closure Property of Addition |
| (B) $a\mathbf{u} \in \mathcal{V}$ | Closure Property of Scalar Multiplication |
| (1) $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ | Commutative Law of Addition |
| (2) $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$ | Associative Law of Addition |
| (3) There is an element $\mathbf{0}$ of \mathcal{V} so that for every \mathbf{y} in \mathcal{V} we have $\mathbf{0} + \mathbf{y} = \mathbf{y} = \mathbf{y} + \mathbf{0}$. | Existence of Identity Element for Addition |
| (4) There is an element $-\mathbf{u}$ in \mathcal{V} such that $\mathbf{u} + (-\mathbf{u}) = \mathbf{0} = (-\mathbf{u}) + \mathbf{u}$. | Existence of Additive Inverse |
| (5) $a(\mathbf{u} + \mathbf{v}) = (a\mathbf{u}) + (a\mathbf{v})$ | Distributive Laws for Scalar Multiplication over Addition |
| (6) $(a + b)\mathbf{u} = (a\mathbf{u}) + (b\mathbf{u})$ | Associativity of Scalar Multiplication |
| (7) $(ab)\mathbf{u} = a(b\mathbf{u})$ | Identity Property for Scalar Multiplication |
| (8) $1\mathbf{u} = \mathbf{u}$ | |

The elements of a vector space \mathcal{V} are called **vectors**.

The two closure properties require that both the operations of vector addition and scalar multiplication always produce an element of the vector space as a result.

¹ We actually define what are called *real vector spaces*, rather than just vector spaces. The word *real* implies that the scalars involved in the scalar multiplication are real numbers. In Chapter 7, we consider *complex vector spaces*, where the scalars are complex numbers. Other types of vector spaces involving more general sets of scalars are not considered in this book.

The standard plus sign, “+,” is used to indicate both vector addition and the sum of real numbers, two different operations. All sums in properties (1), (2), (3), (4), and (5) are vector sums. In property (6), the “+” on the left side of the equation represents addition of real numbers; the “+” on the right side stands for the sum of two vectors. In property (7), the left side of the equation contains one product of real numbers, ab , and one instance of scalar multiplication, (ab) times \mathbf{u} . The right side of property (7) involves two scalar multiplications — first, b times \mathbf{u} , then, a times the vector $(b\mathbf{u})$. Usually we can tell from the context which type of operation is being used.

In any vector space, the additive identity element in property (3) is unique, and the additive inverse (property (4)) of each vector is unique (see the proof of part (3) of Theorem 4.1 and Exercise 12).

Examples of Vector Spaces

Example 1

Let $\mathcal{V} = \mathbb{R}^n$, with addition and scalar multiplication of n -vectors as defined in Section 1.1. Since these operations always produce vectors in \mathbb{R}^n , the closure properties certainly hold for \mathbb{R}^n . By Theorem 1.3, the remaining eight properties hold as well. Thus, $\mathcal{V} = \mathbb{R}^n$ is a vector space with these operations.

Similarly, consider \mathcal{M}_{mn} , the set of $m \times n$ matrices. The usual operations of matrix addition and scalar multiplication on \mathcal{M}_{mn} always produce $m \times n$ matrices, and so the closure properties certainly hold for \mathcal{M}_{mn} . By Theorem 1.11, the remaining eight properties hold as well. Hence, \mathcal{M}_{mn} is a vector space with these operations.

\mathbb{R}^n and \mathcal{M}_{mn} (with the usual operations of addition and scalar multiplication) are representative of most of the vector spaces we consider here. Keep \mathbb{R}^n and \mathcal{M}_{mn} in mind as examples later, as we consider theorems involving general vector spaces.

Some vector spaces can have additional operations. For example, \mathbb{R}^n has the dot product, and \mathcal{M}_{nn} has matrix multiplication and the transpose. But these additional structures are not shared by all vector spaces because they are not included in the definition. We cannot assume the existence of any additional operations in a general discussion of vector spaces. In particular, there is no such operation as multiplication or division of one vector by another in general vector spaces. The only general vector space operation that combines two *vectors* is vector addition.

Example 2

The set $\mathcal{V} = \{\mathbf{0}\}$ is a vector space with the rules for addition and multiplication given by $\mathbf{0} + \mathbf{0} = \mathbf{0}$ and $a\mathbf{0} = \mathbf{0}$ for every scalar (real number) a . Since $\mathbf{0}$ is the only possible result of either operation, \mathcal{V} must be closed under both addition and scalar multiplication. A quick check verifies that the remaining eight properties also hold for \mathcal{V} . This vector space is called the **trivial vector space**, and no smaller vector space is possible (why?).

Example 3


Consider \mathbb{R}^3 as the set of 3-vectors in three-dimensional space, all with initial points at the origin. Let \mathcal{W} be any plane containing the origin. \mathcal{W} can also be considered as the set of all 3-vectors whose terminal point lies in this plane (that is, \mathcal{W} is the set of all 3-vectors that lie entirely in the plane when drawn on a graph, since both the initial point and terminal point of each vector lie in the plane). For example, in Figure 4.1, \mathcal{W} is the plane containing the vectors \mathbf{u} and \mathbf{v} (elements of \mathcal{W}); \mathbf{q} is not in \mathcal{W} because its terminal point does not lie in the plane. We will prove that \mathcal{W} is a vector space.

To check the closure properties, we must show that the sum of any two vectors in \mathcal{W} is a vector in \mathcal{W} and that any scalar multiple of a vector in \mathcal{W} also lies in \mathcal{W} .

If \mathbf{x} and \mathbf{y} are elements of \mathcal{W} , then the parallelogram they form lies entirely in the plane, because \mathbf{x} and \mathbf{y} do. Hence, the diagonal $\mathbf{x} + \mathbf{y}$ of this parallelogram also lies in the plane, so $\mathbf{x} + \mathbf{y}$ is in \mathcal{W} . This observation verifies that \mathcal{W} is closed under vector addition (that is, the closure property holds for vector addition). Notice that it is not enough to know that the sum of two 3-vectors in \mathcal{W} produces another 3-vector. We have to show that the sum they produce is actually in the set \mathcal{W} .

Next consider scalar multiplication. If \mathbf{x} is a vector in \mathcal{W} , then any scalar multiple of \mathbf{x} , $a\mathbf{x}$, is either parallel to \mathbf{x} or equal to $\mathbf{0}$. Therefore, $a\mathbf{x}$ lies in any plane through the origin that contains \mathbf{x} (in particular, \mathcal{W}). Hence, $a\mathbf{x}$ is in \mathcal{W} , and \mathcal{W} is closed under scalar multiplication.

We now check that the remaining eight vector space properties hold. Properties (1), (2), (5), (6), (7), and (8) are true for all vectors in \mathcal{W} by Theorem 1.3, since $\mathcal{W} \subseteq \mathbb{R}^3$. However, properties (3) and (4) must be checked separately for \mathcal{W} because they are *existence* properties. We know that the zero vector and additive inverses exist in \mathbb{R}^3 , but are they in \mathcal{W} ? Now, $\mathbf{0} = [0, 0, 0]$ is in \mathcal{W} , because the plane \mathcal{W} passes through the origin, thus proving property (3). Also, the opposite (additive inverse) of any vector lying in the plane \mathcal{W} also lies in \mathcal{W} , thus proving property (4). Hence, all eight properties and the closure properties are true, so \mathcal{W} is a vector space.

**FIGURE 4.1**

A plane \mathcal{W} in \mathbb{R}^3 containing the origin

Example 4

Let \mathcal{P}_n be the set of polynomials of degree $\leq n$, with real coefficients. The vectors in \mathcal{P}_n have the form $\mathbf{p} = a_n x^n + \cdots + a_1 x + a_0$ for some real numbers a_0, a_1, \dots, a_n . We define addition of polynomials in the usual manner — that is, by adding corresponding coefficients. Then the sum of any two polynomials of degree $\leq n$ also has degree $\leq n$ and so is in \mathcal{P}_n . Thus, the closure property of addition holds. Similarly, if b is a real number and $\mathbf{p} = a_n x^n + \cdots + a_1 x + a_0$ is in \mathcal{P}_n , we define $b\mathbf{p}$ to be the polynomial $(ba_n)x^n + \cdots + (ba_1)x + ba_0$, which is also in \mathcal{P}_n . Hence, the closure property of scalar multiplication holds. Then, if the remaining eight vector space properties hold, \mathcal{P}_n is a vector space under these operations. We verify properties (1), (3), and (4) of the definition and leave the others for you to check.

(1) Commutative Law of Addition: We must show that the order in which two vectors (polynomials) are added makes no difference. Now, by the commutative law of addition for real numbers,

$$\begin{aligned} (a_n x^n + \cdots + a_1 x + a_0) + (b_n x^n + \cdots + b_1 x + b_0) \\ &= (a_n + b_n)x^n + \cdots + (a_1 + b_1)x + (a_0 + b_0) \\ &= (b_n + a_n)x^n + \cdots + (b_1 + a_1)x + (b_0 + a_0) \\ &= (b_n x^n + \cdots + b_1 x + b_0) + (a_n x^n + \cdots + a_1 x + a_0). \end{aligned}$$

(3) Existence of Identity Element for Addition: The zero-degree polynomial $\mathbf{z} = 0x^n + \cdots + 0x + 0$ acts as the additive identity element $\mathbf{0}$. That is, adding \mathbf{z} to any vector $\mathbf{p} = a_n x^n + \cdots + a_1 x + a_0$ does not change the vector:

$$\mathbf{z} + \mathbf{p} = (0 + a_n)x^n + \cdots + (0 + a_1)x + (0 + a_0) = \mathbf{p}.$$

(4) Existence of Additive Inverse: We must show that each vector $\mathbf{p} = a_n x^n + \cdots + a_1 x + a_0$ in \mathcal{P}_n has an additive inverse in \mathcal{P}_n . But, the vector $-\mathbf{p} = -(a_n x^n + \cdots + a_1 x + a_0) = (-a_n)x^n + \cdots + (-a_1)x + (-a_0)$ has the property that $\mathbf{p} + [-\mathbf{p}] = \mathbf{z}$, the zero vector, and so $-\mathbf{p}$ acts as the additive inverse of \mathbf{p} . Because $-\mathbf{p}$ is also in \mathcal{P}_n , we are done. ■

The vector space in Example 4 is similar to our prototype \mathbb{R}^n . For any polynomial in \mathcal{P}_n , consider the sequence of its $n + 1$ coefficients. This sequence completely describes that polynomial and can be thought of as an $(n + 1)$ -vector. For example, a polynomial $a_2 x^2 + a_1 x + a_0$ in \mathcal{P}_2 can be described by the 3-vector $[a_2, a_1, a_0]$. In this way, the vector space \mathcal{P}_2 “resembles” the vector space \mathbb{R}^3 , and in general, \mathcal{P}_n “resembles” \mathbb{R}^{n+1} . We will frequently capitalize on this “resemblance” in an informal way throughout the chapter. We will formalize this relationship between \mathcal{P}_n and \mathbb{R}^{n+1} in Section 5.5.

Example 5

The set \mathcal{P} of all polynomials (of all degrees) is a vector space under the usual (term-by-term) operations of addition and scalar multiplication (see Exercise 15). ■

Example 6

Let \mathcal{V} be the set of all real-valued functions defined on \mathbb{R} . For example, $\mathbf{f}(x) = \arctan(x)$ is in \mathcal{V} . We define addition of functions as usual: $\mathbf{h} = \mathbf{f} + \mathbf{g}$ is the function such that $\mathbf{h}(x) = \mathbf{f}(x) + \mathbf{g}(x)$, for every $x \in \mathbb{R}$. Similarly, if $a \in \mathbb{R}$ and \mathbf{f} is in \mathcal{V} , we define the scalar multiple $\mathbf{h} = a\mathbf{f}$ to be the function such that $\mathbf{h}(x) = a\mathbf{f}(x)$, for every $x \in \mathbb{R}$. Now, the closure properties hold for \mathcal{V} because sums and scalar multiples of real-valued functions produce real-valued functions. To finish verifying that \mathcal{V} is a vector space, we must check that the remaining eight vector space properties hold.

Suppose that \mathbf{f}, \mathbf{g} , and \mathbf{h} are in \mathcal{V} , and a and b are real numbers.

Property (1): For every x in \mathbb{R} , $\mathbf{f}(x)$ and $\mathbf{g}(x)$ are both real numbers. Hence, $\mathbf{f}(x) + \mathbf{g}(x) = \mathbf{g}(x) + \mathbf{f}(x)$ for all $x \in \mathbb{R}$, by the commutative law of addition for real numbers, so each represents the same function of x . Hence, $\mathbf{f} + \mathbf{g} = \mathbf{g} + \mathbf{f}$.

Property (2): For every $x \in \mathbb{R}$, $\mathbf{f}(x) + (\mathbf{g}(x) + \mathbf{h}(x)) = (\mathbf{f}(x) + \mathbf{g}(x)) + \mathbf{h}(x)$, by the associative law of addition for real numbers. Thus, $\mathbf{f} + (\mathbf{g} + \mathbf{h}) = (\mathbf{f} + \mathbf{g}) + \mathbf{h}$.

Property (3): Let \mathbf{z} be the function given by $\mathbf{z}(x) = 0$ for every $x \in \mathbb{R}$. Then, for each x , $\mathbf{z}(x) + \mathbf{f}(x) = 0 + \mathbf{f}(x) = \mathbf{f}(x)$. Hence, $\mathbf{z} + \mathbf{f} = \mathbf{f}$.

Property (4): Given \mathbf{f} in \mathcal{V} , define $-\mathbf{f}$ by $[-\mathbf{f}](x) = -(\mathbf{f}(x))$ for every $x \in \mathbb{R}$. Then, for all x , $[-\mathbf{f}](x) + \mathbf{f}(x) = -(\mathbf{f}(x)) + \mathbf{f}(x) = 0$. Therefore, $[-\mathbf{f}] + \mathbf{f} = \mathbf{z}$, the zero vector, and so the additive inverse of \mathbf{f} is also in \mathcal{V} .

Properties (5) and (6): For every $x \in \mathbb{R}$, $a(\mathbf{f}(x) + \mathbf{g}(x)) = a\mathbf{f}(x) + a\mathbf{g}(x)$ and $(a + b)\mathbf{f}(x) = a\mathbf{f}(x) + b\mathbf{f}(x)$ by the distributive laws for real numbers of multiplication over addition. Hence, $a(\mathbf{f} + \mathbf{g}) = a\mathbf{f} + a\mathbf{g}$, and $(a + b)\mathbf{f} = a\mathbf{f} + b\mathbf{f}$.

Property (7): For every $x \in \mathbb{R}$, $(ab)\mathbf{f}(x) = a(b\mathbf{f}(x))$ follows from the associative law of multiplication for real numbers. Hence, $(ab)\mathbf{f} = a(b\mathbf{f})$.

Property (8): Since $1 \cdot \mathbf{f}(x) = \mathbf{f}(x)$ for every real number x , we have $1 \cdot \mathbf{f} = \mathbf{f}$ in \mathcal{V} . ■

Two Unusual Vector Spaces

The next two examples place unusual operations on familiar sets to create new vector spaces. In such cases, regardless of how the operations are defined, we sometimes use the symbols \oplus and \odot to denote addition and scalar multiplication, respectively, in order to remind ourselves that these operations are not the “regular” ones. Note that \oplus is defined differently in Examples 7 and 8 (and similarly for \odot).

Example 7

Let \mathcal{V} be the set \mathbb{R}^+ of positive real numbers. This set is not a vector space under the usual operations of addition and scalar multiplication (why?). However, we can define new rules for these operations to make \mathcal{V} a vector space. In what follows, we sometimes think of elements of \mathbb{R}^+ as abstract vectors (in which case we use boldface type, such as \mathbf{v}) or as the values on the positive real number line they represent (in which case we use italics, such as v).

To define “addition” on \mathcal{V} , we use *multiplication* of real numbers. That is,

$$\mathbf{v}_1 \oplus \mathbf{v}_2 = v_1 \cdot v_2$$

for every \mathbf{v}_1 and \mathbf{v}_2 in \mathcal{V} , where we use the symbol \oplus for the “addition” operation on \mathcal{V} to emphasize that this is not addition of real numbers. The definition of a vector space states only that vector addition must be a rule for combining two vectors to yield a third vector so that properties (1) through (8) hold. There is no stipulation that vector addition must be at all similar to ordinary addition of real numbers.²

We next define “scalar multiplication,” \odot , on \mathcal{V} by

$$a \odot \mathbf{v} = v^a$$

for every $a \in \mathbb{R}$ and $\mathbf{v} \in \mathcal{V}$.

From the given definitions, we see that if \mathbf{v}_1 and \mathbf{v}_2 are in \mathcal{V} and a is in \mathbb{R} , then both $\mathbf{v}_1 \oplus \mathbf{v}_2$ and $a \odot \mathbf{v}_1$ are in \mathcal{V} , thus verifying the two closure properties. To prove the other eight properties, we assume that $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 \in \mathcal{V}$ and that $a, b \in \mathbb{R}$. We then have the following:

Property (1): $\mathbf{v}_1 \oplus \mathbf{v}_2 = v_1 \cdot v_2 = v_2 \cdot v_1$ (by the commutative law of multiplication for real numbers) $= \mathbf{v}_2 \oplus \mathbf{v}_1$.

Property (2): $\mathbf{v}_1 \oplus (\mathbf{v}_2 \oplus \mathbf{v}_3) = \mathbf{v}_1 \oplus (v_2 \cdot v_3) = v_1 \cdot (v_2 \cdot v_3) = (v_1 \cdot v_2) \cdot v_3$ (by the associative law of multiplication for real numbers) $= (\mathbf{v}_1 \oplus \mathbf{v}_2) \cdot v_3 = (\mathbf{v}_1 \oplus \mathbf{v}_2) \oplus \mathbf{v}_3$.

Property (3): The number 1 in \mathbb{R}^+ acts as the zero vector $\mathbf{0}$ in \mathcal{V} (why?).

Property (4): The additive inverse of \mathbf{v} in \mathcal{V} is the positive real number $(1/v)$, because $\mathbf{v} \oplus (1/v) = v \cdot (1/v) = 1$, the zero vector in \mathcal{V} .

Property (5): $a \odot (\mathbf{v}_1 \oplus \mathbf{v}_2) = a \odot (v_1 \cdot v_2) = (v_1 \cdot v_2)^a = v_1^a \cdot v_2^a = (a \odot \mathbf{v}_1) \cdot (a \odot \mathbf{v}_2) = (a \odot \mathbf{v}_1) \oplus (a \odot \mathbf{v}_2)$.

Property (6): $(a + b) \odot \mathbf{v} = v^{a+b} = v^a \cdot v^b = (a \odot \mathbf{v}) \cdot (b \odot \mathbf{v}) = (a \odot \mathbf{v}) \oplus (b \odot \mathbf{v})$.

Property (7): $(ab) \odot \mathbf{v} = v^{ab} = (v^b)^a = (b \odot \mathbf{v})^a = a \odot (b \odot \mathbf{v})$.

Property (8): $1 \odot \mathbf{v} = v^1 = \mathbf{v}$. ■

Example 8

Let $\mathcal{V} = \mathbb{R}^2$, with addition defined by

$$[x, y] \oplus [w, z] = [x + w + 1, y + z - 2]$$

and scalar multiplication defined by

$$a \odot [x, y] = [ax + a - 1, ay - 2a + 2].$$

The closure properties hold for these operations (why?). In fact, \mathcal{V} forms a vector space because the eight vector properties also hold. We verify properties (2), (3), (4), and (6) and leave the others for you to check.

² You might expect the operation \oplus to be called something other than “addition.” However, most of our vector space terminology comes from the motivating example of \mathbb{R}^n , so the word *addition* is a natural choice for the name of the operation.

Property (2): $[x, y] \oplus ([u, v] \oplus [w, z]) = [x, y] \oplus [u + w + 1, v + z - 2]$

$$= [x + u + w + 2, y + v + z - 4]$$

$$= [x + u + 1, y + v - 2] \oplus [w, z]$$

$$= ([x, y] \oplus [u, v]) \oplus [w, z].$$

Property (3): The vector $[-1, 2]$ acts as the zero vector, since

$$[x, y] \oplus [-1, 2] = [x + (-1) + 1, y + 2 - 2] = [x, y].$$

Property (4): The additive inverse of $[x, y]$ is $[-x - 2, -y + 4]$, because

$$[x, y] \oplus [-x - 2, -y + 4] = [x - x - 2 + 1, y - y + 4 - 2] = [-1, 2],$$

the zero vector in \mathcal{V} .

Property (6):

$$(a + b) \odot [x, y] = [(a + b)x + (a + b) - 1, (a + b)y - 2(a + b) + 2]$$

$$= [(ax + a - 1) + (bx + b - 1) + 1, (ay - 2a + 2) + (by - 2b + 2) - 2]$$

$$= [ax + a - 1, ay - 2a + 2] \oplus [bx + b - 1, by - 2b + 2]$$

$$= (a \odot [x, y]) \oplus (b \odot [x, y]).$$

Some Elementary Properties of Vector Spaces

The next theorem contains several simple results regarding vector spaces. Although these are obviously true in the most familiar examples, we must prove them in general before we know they hold in every possible vector space.

Theorem 4.1 Let \mathcal{V} be a vector space. Then, for every vector \mathbf{v} in \mathcal{V} and every real number a , we have

- | | |
|---|---|
| (1) $a\mathbf{0} = \mathbf{0}$ | Any scalar multiple of the zero vector yields the zero vector. |
| (2) $0\mathbf{v} = \mathbf{0}$ | The scalar zero multiplied by any vector yields the zero vector. |
| (3) $(-1)\mathbf{v} = -\mathbf{v}$ | The scalar -1 multiplied by any vector yields the additive inverse of that vector. |
| (4) If $a\mathbf{v} = \mathbf{0}$, then $a = 0$ or $\mathbf{v} = \mathbf{0}$. | If a scalar multiplication yields the zero vector, then either the scalar is zero, or the vector is the zero vector, or both. |

Part (3) justifies the notation for the additive inverse in property (4) of the definition of a vector space and shows we do not need to distinguish between $-\mathbf{v}$ and $(-1)\mathbf{v}$.

This theorem must be proved directly from the properties in the definition of a vector space because at this point we have no other known facts about general vector spaces. We prove parts (1), (3), and (4). The proof of part (2) is similar to the proof of part (1) and is left as Exercise 18.

Proof. (Abridged):

Part (1): By direct proof,

$$\begin{aligned}
 a\mathbf{0} &= a\mathbf{0} + \mathbf{0} && \text{by property (3)} \\
 &= a\mathbf{0} + (a\mathbf{0} + (-[a\mathbf{0}])) && \text{by property (4)} \\
 &= (a\mathbf{0} + a\mathbf{0}) + (-[a\mathbf{0}]) && \text{by property (2)} \\
 &= a(\mathbf{0} + \mathbf{0}) + (-[a\mathbf{0}]) && \text{by property (5)} \\
 &= a\mathbf{0} + (-[a\mathbf{0}]) && \text{by property (3)} \\
 &= \mathbf{0}. && \text{by property (4)}
 \end{aligned}$$

Part (3): First, note that $\mathbf{v} + (-1)\mathbf{v} = 1\mathbf{v} + (-1)\mathbf{v}$ (by property (8)) $= (1 + (-1))\mathbf{v}$ (by property (6)) $= 0\mathbf{v} = \mathbf{0}$ (by part (2) of Theorem 4.1). Therefore, $(-1)\mathbf{v}$ acts as an additive inverse for \mathbf{v} . We will finish the proof by showing that the additive inverse for \mathbf{v} is unique. Hence, $(-1)\mathbf{v}$ will be *the* additive inverse of \mathbf{v} .

Suppose that \mathbf{x} and \mathbf{y} are both additive inverses for \mathbf{v} . Thus, $\mathbf{x} + \mathbf{v} = \mathbf{0}$ and $\mathbf{v} + \mathbf{y} = \mathbf{0}$. Hence,

$$\mathbf{x} = \mathbf{x} + \mathbf{0} = \mathbf{x} + (\mathbf{v} + \mathbf{y}) = (\mathbf{x} + \mathbf{v}) + \mathbf{y} = \mathbf{0} + \mathbf{y} = \mathbf{y}.$$

Therefore, any two additive inverses of \mathbf{v} are equal. (Note that this is, in essence, the same proof we gave for Theorem 2.10, the uniqueness of inverse for matrix multiplication. You should compare these proofs.)

Part (4): This is an “If A then B or C ” statement. Therefore, we assume that $a\mathbf{v} = \mathbf{0}$ and $a \neq 0$ and show that $\mathbf{v} = \mathbf{0}$. Now,

$$\begin{aligned}
 \mathbf{v} &= 1\mathbf{v} && \text{by property (8)} \\
 &= \left(\frac{1}{a} \cdot a\right)\mathbf{v} && \text{because } a \neq 0 \\
 &= \left(\frac{1}{a}\right)(a\mathbf{v}) && \text{by property (7)} \\
 &= \left(\frac{1}{a}\right)\mathbf{0} && \text{because } a\mathbf{v} = \mathbf{0} \\
 &= \mathbf{0}. && \text{by part (1) of Theorem 4.1} \quad \square
 \end{aligned}$$

Theorem 4.1 is valid even for unusual vector spaces, such as those in Examples 7 and 8. For instance, part (4) of the theorem claims that, in general, $a\mathbf{v} = \mathbf{0}$ implies $a = 0$ or $\mathbf{v} = \mathbf{0}$. This statement can quickly be verified for the vector space $\mathcal{V} = \mathbb{R}^+$ with operations \oplus and \odot from Example 7. In this case, $a \odot \mathbf{v} = v^a$, and the zero vector $\mathbf{0}$ is the real number 1. Then, part (4) is equivalent here to the true statement that $v^a = 1$ implies $a = 0$ or $v = 1$.

Applying parts (2) and (3) of Theorem 4.1 to an unusual vector space \mathcal{V} gives a quick way of finding the zero vector $\mathbf{0}$ of \mathcal{V} and the additive inverse $-\mathbf{v}$ for any vector \mathbf{v} in \mathcal{V} . For instance, in Example 8, we have $\mathcal{V} = \mathbb{R}^2$ with scalar multiplication $a \odot [x, y] = [ax + a - 1, ay - 2a + 2]$. To find the zero vector $\mathbf{0}$ in \mathcal{V} , we simply multiply the scalar 0 by any general vector $[x, y]$ in \mathcal{V} :

$$\mathbf{0} = 0 \odot [x, y] = [0x + 0 - 1, 0y - 2(0) + 2] = [-1, 2].$$

Similarly, if $[x, y] \in \mathcal{V}$, then $-1 \odot [x, y]$ gives the additive inverse of $[x, y]$.

$$\begin{aligned} -[x, y] &= -1 \odot [x, y] = [-1x + (-1) - 1, -1y - 2(-1) + 2] \\ &= [-x - 2, -y + 4]. \end{aligned}$$

Failure of the Vector Space Conditions

We conclude this section by considering some sets that are not vector spaces to see what can go wrong.

Example 9

The set Φ of real-valued functions, f , defined on the interval $[0, 1]$ such that $f(\frac{1}{2}) = 1$, is not a vector space under the usual operations of function addition and scalar multiplication because the closure properties do not hold. If f and g are in Φ , then

$$(f + g)\left(\frac{1}{2}\right) = f\left(\frac{1}{2}\right) + g\left(\frac{1}{2}\right) = 1 + 1 = 2 \neq 1,$$

so $f + g$ is not in Φ . Therefore, Φ is not closed under addition and cannot be a vector space. (Is Φ closed under scalar multiplication?)

Example 10

Let \mathcal{Y} be the set \mathbb{R}^2 with operations

$$\mathbf{v}_1 \oplus \mathbf{v}_2 = \mathbf{v}_1 + \mathbf{v}_2 \quad \text{and} \quad c \odot \mathbf{v} = c(\mathbf{A}\mathbf{v}), \quad \text{where } \mathbf{A} = \begin{bmatrix} -3 & 1 \\ 5 & -2 \end{bmatrix}.$$

With these operations, \mathcal{Y} is not a vector space. You can verify that \mathcal{Y} is closed under \oplus and \odot , but properties (7) and (8) of the definition are not satisfied. For example, property (8) fails since

$$1 \odot \begin{bmatrix} 2 \\ 7 \end{bmatrix} = 1 \left(\begin{bmatrix} -3 & 1 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} 2 \\ 7 \end{bmatrix} \right) = 1 \begin{bmatrix} 1 \\ -4 \end{bmatrix} = \begin{bmatrix} 1 \\ -4 \end{bmatrix} \neq \begin{bmatrix} 2 \\ 7 \end{bmatrix}.$$

New Vocabulary

closure properties	vector addition (in a general vector space)
scalar multiplication (in a general vector space)	vector space
trivial vector space	vectors (in a general vector space)

Highlights

- Vector spaces have two specified operations: vector addition (+) and scalar multiplication (\cdot). A vector space is closed under these operations and possesses eight additional fundamental properties (as stated in the definition).
- The smallest possible vector space is the trivial vector space.
- Familiar vector spaces (under natural operations) include \mathbb{R}^n , \mathcal{M}_{mn} , \mathcal{P}_n , \mathcal{P} , a line through the origin, a plane through the origin, all real-valued functions.
- Any scalar multiple of the zero vector equals the zero vector.
- The scalar 0 times any vector equals the zero vector.
- The scalar -1 times any vector gives the additive inverse of the vector.
- If a scalar multiple of a vector equals the zero vector, then either the scalar is zero or the vector is zero.

EXERCISES FOR SECTION 4.1

Remember: To verify that a given set with its operations is a vector space, you must prove the two closure properties as well as the remaining eight properties in the definition. To show that a set with operations is *not* a vector space, you need only find an example showing that one of the closure properties or one of the remaining eight properties is not satisfied.

1. Rewrite properties (2), (5), (6), and (7) in the definition of a vector space using the symbols \oplus for vector addition and \odot for scalar multiplication. (The notations for real number addition and multiplication should not be changed.)
2. Prove that the set of all scalar multiples of the vector $[1, 3, 2]$ in \mathbb{R}^3 forms a vector space with the usual operations on 3-vectors.
3. Verify that the set of polynomials \mathbf{f} in \mathcal{P}_3 such that $\mathbf{f}(2) = 0$ forms a vector space with the standard operations.
4. Prove that \mathbb{R} is a vector space using the operations \oplus and \odot given by $\mathbf{x} \oplus \mathbf{y} = (x^3 + y^3)^{1/3}$ and $a \odot \mathbf{x} = (\sqrt[3]{a})x$.

- ★5. Show that the set of singular 2×2 matrices under the usual operations is *not* a vector space.
6. Prove that the set of nonsingular $n \times n$ matrices under the usual operations is *not* a vector space.
7. Show that \mathbb{R} , with ordinary addition but with scalar multiplication replaced by $a \odot \mathbf{x} = \mathbf{0}$ for every real number a , is *not* a vector space.
- ★8. Show that the set \mathbb{R} , with the usual scalar multiplication but with addition given by $x \oplus y = 2(x + y)$, is *not* a vector space.
9. Show that the set \mathbb{R}^2 , with the usual scalar multiplication but with vector addition replaced by $[x, y] \oplus [w, z] = [x + w, 0]$, does *not* form a vector space.
10. Let $\mathcal{A} = \mathbb{R}$, with the operations \oplus and \odot given by $\mathbf{x} \oplus \mathbf{y} = (x^5 + y^5)^{1/5}$ and $a \odot \mathbf{x} = ax$. Determine whether \mathcal{A} is a vector space. Prove your answer.
11. Let \mathbf{A} be a fixed $m \times n$ matrix, and let \mathbf{B} be a fixed m -vector (in \mathbb{R}^m). Let \mathcal{V} be the set of solutions \mathbf{X} (in \mathbb{R}^n) to the matrix equation $\mathbf{AX} = \mathbf{B}$. Endow \mathcal{V} with the usual n -vector operations.
 - (a) Assume \mathcal{V} is nonempty. Show that the closure properties are satisfied in \mathcal{V} if and only if $\mathbf{B} = \mathbf{0}$.
 - (b) Explain why properties (1), (2), (5), (6), (7), and (8) in the definition of a vector space have already been proved for \mathcal{V} in Theorem 1.3.
 - (c) Prove that property (3) in the definition of a vector space is satisfied if and only if $\mathbf{B} = \mathbf{0}$.
 - (d) Explain why property (4) in the definition makes no sense unless property (3) is satisfied. Prove property (4) when $\mathbf{B} = \mathbf{0}$.
 - (e) Use parts (a) through (d) of this exercise to determine necessary and sufficient conditions for \mathcal{V} to be a vector space.
12. Let \mathcal{V} be a vector space. Prove that the identity element for vector addition in \mathcal{V} is unique. (Hint: Use a proof by contradiction.)
13. The set \mathbb{R}^2 with operations $[x, y] \oplus [w, z] = [x + w - 2, y + z + 3]$ and $a \odot [x, y] = [ax - 2a + 2, ay + 3a - 3]$ is a vector space. Use parts (2) and (3) of Theorem 4.1 to find the zero vector $\mathbf{0}$ and the additive inverse of each vector $\mathbf{v} = [x, y]$ for this vector space. Then check your answers.
14. Let \mathcal{V} be a vector space. Prove the following **cancellation laws**:
 - (a) If \mathbf{u}, \mathbf{v} , and \mathbf{w} are vectors in \mathcal{V} for which $\mathbf{u} + \mathbf{v} = \mathbf{w} + \mathbf{v}$, then $\mathbf{u} = \mathbf{w}$.
 - (b) If a and b are scalars and $\mathbf{v} \neq \mathbf{0}$ is a vector in \mathcal{V} with $a\mathbf{v} = b\mathbf{v}$, then $a = b$.
 - (c) If $a \neq 0$ is a scalar and $\mathbf{v}, \mathbf{w} \in \mathcal{V}$ with $a\mathbf{v} = a\mathbf{w}$, then $\mathbf{v} = \mathbf{w}$.

15. Prove that the set \mathcal{P} of all polynomials with real coefficients forms a vector space under the usual operations of polynomial (term-by-term) addition and scalar multiplication.
16. Let X be any set, and let $\mathcal{V} = \{\text{all real-valued functions with domain } X\}$. Prove that \mathcal{V} is a vector space using ordinary addition and scalar multiplication of real-valued functions. (Hint: Alter the proof in Example 6 appropriately.)
17. Let $\mathbf{v}_1, \dots, \mathbf{v}_n$ be vectors in a vector space \mathcal{V} , and let a_1, \dots, a_n be any real numbers. Use induction to prove that $\sum_{i=1}^n a_i \mathbf{v}_i$ is in \mathcal{V} .
18. Prove part (2) of Theorem 4.1.
19. Prove that every nontrivial vector space has an infinite number of distinct elements.
- ★20. True or False:
 - (a) The set \mathbb{R}^n under any operations of “addition” and “scalar multiplication” is a vector space.
 - (b) The set of all polynomials of degree 7 is a vector space under the usual operations of addition and scalar multiplication.
 - (c) The set of all polynomials of degree ≤ 7 is a vector space under the usual operations of addition and scalar multiplication.
 - (d) If \mathbf{x} is a vector in a vector space \mathcal{V} , and c is a nonzero scalar, then $c\mathbf{x} = \mathbf{0}$ implies $\mathbf{x} = \mathbf{0}$.
 - (e) In a vector space, scalar multiplication by the zero scalar always results in the zero scalar.
 - (f) In a vector space, scalar multiplication of a vector \mathbf{x} by -1 always results in the additive inverse of \mathbf{x} .
 - (g) The set of all real-valued functions f on \mathbb{R} such that $f(1) = 0$ is a vector space under the usual operations of addition and scalar multiplication.

4.2 SUBSPACES

Section 4.1 presented several examples in which two vector spaces share the same addition and scalar multiplication operations, with one as a subset of the other. In fact, most of these examples involve subsets of either \mathbb{R}^n , \mathcal{M}_{mn} , or the vector space of real-valued functions defined on some set (see Exercise 16 in Section 4.1). As we will see, when a vector space is a subset of a known vector space and has the same operations, it becomes easier to handle. These subsets, called **subspaces**, also provide additional information about the larger vector space.

Definition of a Subspace and Examples

Definition Let \mathcal{V} be a vector space. Then \mathcal{W} is a **subspace** of \mathcal{V} if and only if \mathcal{W} is a subset of \mathcal{V} , and \mathcal{W} is itself a vector space with the same operations as \mathcal{V} .

That is, \mathcal{W} is a subspace of \mathcal{V} if and only if \mathcal{W} is a vector space inside \mathcal{V} such that for every a in \mathbb{R} and every \mathbf{v} and \mathbf{w} in \mathcal{W} , $\mathbf{v} + \mathbf{w}$ and $a\mathbf{v}$ yield the same vectors when the operations are performed in \mathcal{W} as when they are performed in \mathcal{V} .

Example 1

Example 3 of Section 4.1 showed that the set of points lying on a plane \mathcal{W} through the origin in \mathbb{R}^3 forms a vector space under the usual addition and scalar multiplication in \mathbb{R}^3 . \mathcal{W} is certainly a subset of \mathbb{R}^3 . Hence, the vector space \mathcal{W} is a subspace of \mathbb{R}^3 .

Example 2

The set \mathcal{S} of scalar multiples of the vector $[1, 3, 2]$ in \mathbb{R}^3 forms a vector space under the usual addition and scalar multiplication in \mathbb{R}^3 (see Exercise 2 in Section 4.1). \mathcal{S} is certainly a subset of \mathbb{R}^3 . Hence, \mathcal{S} is a subspace of \mathbb{R}^3 . Notice that \mathcal{S} corresponds geometrically to the set of points lying on the line through the origin in \mathbb{R}^3 in the direction of the vector $[1, 3, 2]$ (see Figure 4.2). In the same manner, every line through the origin determines a subspace of \mathbb{R}^3 — namely, the set of scalar multiples of a nonzero vector in the direction of that line.




FIGURE 4.2

Line containing all scalar multiples of $[1, 3, 2]$

Example 3

Let \mathcal{V} be any vector space. Then \mathcal{V} is a subspace of itself (why?). Also, if \mathcal{W} is the subset $\{\mathbf{0}\}$ of \mathcal{V} , then \mathcal{W} is a vector space under the same operations as \mathcal{V} (see Example 2 of Section 4.1). Therefore, $\mathcal{W} = \{\mathbf{0}\}$ is a subspace of \mathcal{V} .

Although the subspaces \mathcal{V} and $\{\mathbf{0}\}$ of a vector space \mathcal{V} are important, they occasionally complicate matters because they must be considered as special cases in proofs. The subspace $\mathcal{W} = \{\mathbf{0}\}$ is called the **trivial subspace** of \mathcal{V} . A vector space containing at least one nonzero vector has at least two distinct subspaces, the trivial subspace and the vector space itself. In fact, under the usual operations, \mathbb{R} has only these two subspaces (see Exercise 16).

All subspaces of \mathcal{V} other than \mathcal{V} itself are called **proper subspaces** of \mathcal{V} . If we consider Examples 1 to 3 in the context of \mathbb{R}^3 , we find at least four different types of subspaces of \mathbb{R}^3 . These are the trivial subspace $\{[0, 0, 0]\} = \{\mathbf{0}\}$, subspaces like Example 2 that can be geometrically represented as a line (thus “resembling” \mathbb{R}), subspaces like Example 1 that can be represented as a plane (thus “resembling” \mathbb{R}^2), and the subspace \mathbb{R}^3 itself.³ All but the last are proper subspaces. Later we will show that each subspace of \mathbb{R}^3 is, in fact, one of these four types. Similarly, we will show later that all subspaces of \mathbb{R}^n “resemble” $\{\mathbf{0}\}$, \mathbb{R} , \mathbb{R}^2 , \mathbb{R}^3 , \dots , \mathbb{R}^{n-1} , or \mathbb{R}^n .

Example 4

Consider the vector spaces (using ordinary function addition and scalar multiplication) in the following chain:

$$\begin{aligned} \mathcal{P}_0 \subset \mathcal{P}_1 \subset \mathcal{P}_2 \subset \dots &\subset \mathcal{P} \\ &\subset \{\text{differentiable real-valued functions on } \mathbb{R}\} \\ &\subset \{\text{continuous real-valued functions on } \mathbb{R}\} \\ &\subset \{\text{all real-valued functions on } \mathbb{R}\}. \end{aligned}$$

Some of these we encountered in Section 4.1, and the rest are discussed in Exercise 7 of this section. Each of these vector spaces is a proper subspace of every vector space after it in the chain (why?).

When Is a Subset a Subspace?

It is important to note that not every subset of a vector space is a subspace. A subset \mathcal{S} of a vector space \mathcal{V} fails to be a subspace of \mathcal{V} if \mathcal{S} does not satisfy the properties of a vector space in its own right or if \mathcal{S} does not use the same operations as \mathcal{V} .

³ Although some subspaces of \mathbb{R}^3 “resemble” \mathbb{R} and \mathbb{R}^2 geometrically, note that \mathbb{R} and \mathbb{R}^2 are not actually subspaces of \mathbb{R}^3 because they are not subsets of \mathbb{R}^3 .

Example 5

Consider the first quadrant in \mathbb{R}^2 — that is, the set Ω of all 2-vectors of the form $[x, y]$ where $x \geq 0$ and $y \geq 0$. This subset Ω of \mathbb{R}^2 is not a vector space under the normal operations of \mathbb{R}^2 because it is not closed under scalar multiplication. (For example, $[3, 4]$ is in Ω , but $-2 \cdot [3, 4] = [-6, -8]$ is not in Ω .) Therefore, Ω cannot be a subspace of \mathbb{R}^2 .

Example 6

Consider the vector space \mathbb{R} under the usual operations. Let \mathcal{W} be the subset \mathbb{R}^+ . By Example 7 of Section 4.1, we know that \mathcal{W} is a vector space under the unusual operations \oplus and \odot , where \oplus represents multiplication and \odot represents exponentiation. Although \mathcal{W} is a nonempty subset of \mathbb{R} and is itself a vector space, \mathcal{W} is not a subspace of \mathbb{R} because \mathcal{W} and \mathbb{R} do not share the same operations.

The following theorem provides a shortcut for verifying that a (nonempty) subset \mathcal{W} of a vector space is a subspace; if the closure properties hold for \mathcal{W} , then the remaining eight vector space properties automatically follow as well.

Theorem 4.2 Let \mathcal{V} be a vector space, and let \mathcal{W} be a nonempty subset of \mathcal{V} using the same operations. Then \mathcal{W} is a subspace of \mathcal{V} if and only if \mathcal{W} is closed under vector addition and scalar multiplication in \mathcal{V} .

Notice that this theorem applies only to *nonempty subsets* of a vector space. Even though the empty set is a subset of every vector space, it is not a subspace of any vector space because it does not contain an additive identity.

Proof. Since this is an “if and only if” statement, the proof has two parts. First we must show that if \mathcal{W} is a subspace of \mathcal{V} , then it is closed under the two operations. Now, as a subspace, \mathcal{W} is itself a vector space. Hence, the closure properties hold for \mathcal{W} as they do for any vector space.

For the other part of the proof, we must show that if the closure properties hold for a nonempty subset \mathcal{W} of \mathcal{V} , then \mathcal{W} is itself a vector space under the operations in \mathcal{V} . That is, we must prove the remaining eight vector space properties hold for \mathcal{W} .

Properties (1), (2), (5), (6), (7), and (8) are all true in \mathcal{W} because they are true in \mathcal{V} , a known vector space. That is, since these properties hold for all vectors in \mathcal{V} , they must be true for all vectors in its subset, \mathcal{W} . For example, to prove property (1) for \mathcal{W} , let $\mathbf{u}, \mathbf{v} \in \mathcal{W}$. Then,

$$\begin{aligned}
 \underbrace{\mathbf{u} + \mathbf{v}}_{\text{addition in } \mathcal{W}} &= \underbrace{\mathbf{u} + \mathbf{v}}_{\text{addition in } \mathcal{V}} && \text{because } \mathcal{W} \text{ and } \mathcal{V} \text{ share the same operations} \\
 &= \underbrace{\mathbf{v} + \mathbf{u}}_{\text{addition in } \mathcal{V}} && \text{because } \mathcal{V} \text{ is a vector space and property (1) holds} \\
 &= \underbrace{\mathbf{v} + \mathbf{u}}_{\text{addition in } \mathcal{W}} && \text{because } \mathcal{W} \text{ and } \mathcal{V} \text{ share the same operations}
 \end{aligned}$$

Next we prove property (3), the existence of an additive identity in \mathcal{W} . Because \mathcal{W} is nonempty, we can choose an element \mathbf{w}_1 from \mathcal{W} . Now \mathcal{W} is closed under scalar

multiplication, so $0\mathbf{w}_1$ is in \mathcal{W} . However, since this is the same operation as in \mathcal{V} , a known vector space, part (2) of Theorem 4.1 implies that $0\mathbf{w}_1 = \mathbf{0}$. Hence, $\mathbf{0}$ is in \mathcal{W} . Because $\mathbf{0} + \mathbf{v} = \mathbf{v}$ for all \mathbf{v} in \mathcal{V} , it follows that $\mathbf{0} + \mathbf{w} = \mathbf{w}$ for all \mathbf{w} in \mathcal{W} . Therefore, \mathcal{W} contains the same additive identity that \mathcal{V} has.

Finally, we must prove that property (4), the existence of additive inverses, holds for \mathcal{W} . Let $\mathbf{w} \in \mathcal{W}$. Then $\mathbf{w} \in \mathcal{V}$. Part (3) of Theorem 4.1 shows $(-1)\mathbf{w}$ is the additive inverse of \mathbf{w} in \mathcal{V} . If we can show that this additive inverse is also in \mathcal{W} , we will be done. But since \mathcal{W} is closed under scalar multiplication, $(-1)\mathbf{w} \in \mathcal{W}$. \square

Checking for Subspaces in \mathcal{M}_{nn} and \mathbb{R}^n

In the next three examples, we apply Theorem 4.2 to determine whether several subsets of \mathcal{M}_{nn} and \mathbb{R}^n are subspaces. Assume that \mathcal{M}_{nn} and \mathbb{R}^n have the usual operations.

Example 7

Consider \mathcal{U}_n , the set of upper triangular $n \times n$ matrices. Since \mathcal{U}_n is nonempty, we may apply Theorem 4.2 to see whether \mathcal{U}_n is a subspace of \mathcal{M}_{nn} . Closure of \mathcal{U}_n under vector addition holds because the sum of any two $n \times n$ upper triangular matrices is again upper triangular. The closure property in \mathcal{U}_n for scalar multiplication also holds, since any scalar multiple of an upper triangular matrix is again upper triangular. Hence, \mathcal{U}_n is a subspace of \mathcal{M}_{nn} .

Similar arguments show that \mathcal{L}_n (lower triangular $n \times n$ matrices) and \mathcal{D}_n (diagonal $n \times n$ matrices) are also subspaces of \mathcal{M}_{nn} . \blacksquare

The subspace \mathcal{D}_n of \mathcal{M}_{nn} in Example 7 is the intersection of the subspaces \mathcal{U}_n and \mathcal{L}_n . In fact, the intersection of subspaces of a vector space always produces a subspace under the same operations (see Exercise 18).

If either closure property fails to hold for a subset, it cannot be a subspace. For this reason, none of the following subsets of \mathcal{M}_{nn} , $n \geq 2$, is a subspace:

- (1) the set of nonsingular $n \times n$ matrices
- (2) the set of singular $n \times n$ matrices
- (3) the set of $n \times n$ matrices in reduced row echelon form.

You should check that the closure property for vector addition fails in each case and that the closure property for scalar multiplication fails in (1) and (3).

Example 8

Let \mathcal{Y} be the set of vectors in \mathbb{R}^4 of the form $[a, 0, b, 0]$, that is, 4-vectors whose second and fourth coordinates are zero. We prove that \mathcal{Y} is a subspace of \mathbb{R}^4 by checking the closure properties.

To prove closure under vector addition, we must add two arbitrary elements of \mathcal{Y} and check that the result has the correct form for a vector in \mathcal{Y} . Now, $[a, 0, b, 0] + [c, 0, d, 0] = [(a+c), 0, (b+d), 0]$. The second and fourth coordinates of the sum are zero, so \mathcal{Y} is closed under addition. Similarly, we must prove closure under scalar multiplication. Now,

$k[a, 0, b, 0] = [ka, 0, kb, 0]$. Since the second and fourth coordinates of the product are zero, \mathcal{V} is closed under scalar multiplication. Hence, by Theorem 4.2, \mathcal{V} is a subspace of \mathbb{R}^4 . ■

Example 9

Let \mathcal{W} be the set of vectors in \mathbb{R}^3 of the form $[a, b, \frac{1}{2}a - 2b]$, that is, 3-vectors whose third coordinate is half the first coordinate minus twice the second coordinate. We show that \mathcal{W} is a subspace of \mathbb{R}^3 by checking the closure properties.

Checking closure under vector addition, we have

$$\begin{aligned} \left[a, b, \frac{1}{2}a - 2b \right] + \left[c, d, \frac{1}{2}c - 2d \right] &= \left[a + c, b + d, \frac{1}{2}a - 2b + \frac{1}{2}c - 2d \right] \\ &= \left[a + c, b + d, \frac{1}{2}(a + c) - 2(b + d) \right], \end{aligned}$$

which has the required form, since it equals $[A, B, \frac{1}{2}A - 2B]$, where $A = a + c$ and $B = b + d$.

Checking closure under scalar multiplication, we get

$$k \left[a, b, \frac{1}{2}a - 2b \right] = \left[ka, kb, k \left(\frac{1}{2}a - 2b \right) \right] = \left[ka, kb, \frac{1}{2}(ka) - 2(kb) \right],$$

which has the required form (why?).

Note that

$$\left[a, b, \frac{1}{2}a - 2b \right] = a \left[1, 0, \frac{1}{2} \right] + b[0, 1, -2],$$

and so \mathcal{W} consists of the set of all linear combinations of $[1, 0, \frac{1}{2}]$ and $[0, 1, -2]$. Geometrically, \mathcal{W} is the plane in \mathbb{R}^3 through the origin containing the vectors $[1, 0, \frac{1}{2}]$ and $[0, 1, -2]$, shown in Figure 4.3. This plane is the set of all possible “destinations” using these two directions (starting from the origin). This is the type of subspace of \mathbb{R}^3 discussed in Example 1.




FIGURE 4.3

The plane through the origin containing $[1, 0, \frac{1}{2}]$ and $[0, 1, -2]$

The following subsets of \mathbb{R}^n are not subspaces. In each case, at least one of the two closure properties fails. (Can you determine which properties?)

- (1) The set of n -vectors whose first coordinate is nonnegative (in \mathbb{R}^2 , this set is a half-plane)
- (2) The set of unit n -vectors (in \mathbb{R}^3 , this set is a sphere)
- (3) For $n \geq 2$, the set of n -vectors with a zero in at least one coordinate (in \mathbb{R}^3 , this set is the union of three planes)
- (4) The set of n -vectors having all integer coordinates
- (5) For $n \geq 2$, the set of all n -vectors whose first two coordinates add up to 3 (in \mathbb{R}^2 , this is the line $x + y = 3$)

The subsets (2) and (5), which do not contain the additive identity $\mathbf{0}$ of \mathbb{R}^n , can quickly be disqualified as subspaces. In general,

If a subset \mathcal{S} of a vector space \mathcal{V} does not contain the zero vector $\mathbf{0}$ of \mathcal{V} , then \mathcal{S} is not a subspace of \mathcal{V} .

Checking for the presence of the additive identity is usually easy and thus is a fast way to show that certain subsets are not subspaces.

Linear Combinations Remain in a Subspace

As in Chapter 1, we define a **linear combination** of vectors in a general vector space to be a sum of scalar multiples of the vectors. The next theorem asserts that if a finite set of vectors is in a given subspace of a vector space, then so is any linear combination of those vectors.

Theorem 4.3 Let \mathcal{W} be a subspace of a vector space \mathcal{V} , and let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be vectors in \mathcal{W} . Then, for any scalars a_1, a_2, \dots, a_n , we have $a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n \in \mathcal{W}$.

Essentially, this theorem points out that a subspace is “closed under linear combinations.” That is, when the vectors of a subspace are used to form linear combinations, all possible “destination vectors” remain in the subspace.

Proof. Suppose that \mathcal{W} is a subspace of a vector space \mathcal{V} . We give a proof by induction on n .

Base Step: If $n = 1$, then we must show that if $\mathbf{v}_1 \in \mathcal{W}$ and a_1 is a scalar, then $a_1\mathbf{v}_1 \in \mathcal{W}$. But this is certainly true since the subspace \mathcal{W} is closed under scalar multiplication.

Inductive Step: Assume that the theorem is true for any linear combination of n vectors in \mathcal{W} . We must prove the theorem holds for a linear combination of $n + 1$ vectors. Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n, \mathbf{v}_{n+1}$ are vectors in \mathcal{W} , and $a_1, a_2, \dots, a_n, a_{n+1}$ are scalars. We must show

that $a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_n\mathbf{v}_n + a_{n+1}\mathbf{v}_{n+1} \in \mathcal{W}$. However, by the inductive hypothesis, we know that $a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_n\mathbf{v}_n \in \mathcal{W}$. Also, $a_{n+1}\mathbf{v}_{n+1} \in \mathcal{W}$, since \mathcal{W} is closed under scalar multiplication. But since \mathcal{W} is also closed under addition, the sum of any two vectors in \mathcal{W} is again in \mathcal{W} , so $(a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \cdots + a_n\mathbf{v}_n) + (a_{n+1}\mathbf{v}_{n+1}) \in \mathcal{W}$. \square

Example 10

In Example 9, we found that the set \mathcal{W} of all vectors of the form $[a, b, \frac{1}{2}a - 2b]$ is a subspace of \mathbb{R}^3 . In particular, $[1, 0, \frac{1}{2}]$ and $[0, 1, -2]$ are in \mathcal{W} . By Theorem 4.3, any linear combination of these vectors is also in \mathcal{W} . For example, $6[1, 0, \frac{1}{2}] - 5[0, 1, -2] = [6, -5, 13]$ and $-4[1, 0, \frac{1}{2}] + 2[0, 1, -2] = [-4, 2, -6]$ are both in \mathcal{W} . Of course, this makes sense geometrically, since \mathcal{W} is a plane through the origin, and any linear combination of vectors in such a plane remains in that plane. \blacksquare

An Eigenspace Is a Subspace

We conclude this section by noting that any eigenspace of an $n \times n$ matrix is a subspace of \mathbb{R}^n . (In fact, this is why the word “space” appears in the term “eigenspace.”)

Theorem 4.4 Let \mathbf{A} be an $n \times n$ matrix, and let λ be an eigenvalue for \mathbf{A} , having eigenspace E_λ . Then E_λ is a subspace of \mathbb{R}^n .

Proof. Let λ be an eigenvalue for an $n \times n$ matrix \mathbf{A} . By definition, the eigenspace E_λ of λ is the set of all n -vectors \mathbf{X} having the property that $\mathbf{A}\mathbf{X} = \lambda\mathbf{X}$, including the zero n -vector. We will use Theorem 4.2 to show that E_λ is a subspace of \mathbb{R}^n .

Since $\mathbf{0} \in E_\lambda$, E_λ is a nonempty subset of \mathbb{R}^n . We must show that E_λ is closed under addition and scalar multiplication.

Let $\mathbf{X}_1, \mathbf{X}_2$ be any two vectors in E_λ . To show that $\mathbf{X}_1 + \mathbf{X}_2 \in E_\lambda$, we need to verify that $\mathbf{A}(\mathbf{X}_1 + \mathbf{X}_2) = \lambda(\mathbf{X}_1 + \mathbf{X}_2)$. But, $\mathbf{A}(\mathbf{X}_1 + \mathbf{X}_2) = \mathbf{A}\mathbf{X}_1 + \mathbf{A}\mathbf{X}_2 = \lambda\mathbf{X}_1 + \lambda\mathbf{X}_2 = \lambda(\mathbf{X}_1 + \mathbf{X}_2)$.

Similarly, let \mathbf{X} be a vector in E_λ , and let c be a scalar. We must show that $c\mathbf{X} \in E_\lambda$. But, $\mathbf{A}(c\mathbf{X}) = c(\mathbf{A}\mathbf{X}) = c(\lambda\mathbf{X}) = \lambda(c\mathbf{X})$, and so $c\mathbf{X} \in E_\lambda$. Hence, E_λ is a subspace of \mathbb{R}^n . \square

Example 11

Consider

$$\mathbf{A} = \begin{bmatrix} 16 & -4 & -2 \\ 3 & 3 & -6 \\ 2 & -8 & 11 \end{bmatrix}.$$

Computing $|\lambda\mathbf{I}_3 - \mathbf{A}|$ produces $p_{\mathbf{A}}(x) = x^3 - 30x^2 + 225x = x(x - 15)^2$. Solving $(0\mathbf{I}_3 - \mathbf{A})\mathbf{X} = \mathbf{0}$ yields $E_0 = \{c[1, 3, 2] \mid c \in \mathbb{R}\}$. Thus, the eigenspace for $\lambda_1 = 0$ is the subspace of \mathbb{R}^3 from Example 2. Similarly, solving $(15\mathbf{I}_3 - \mathbf{A})\mathbf{X} = \mathbf{0}$ gives $E_{15} = \{a[4, 1, 0] + b[2, 0, 1] \mid a, b \in \mathbb{R}\}$. By Theorem 4.4, E_{15} is also a subspace of \mathbb{R}^3 . Although it is not obvious, E_{15} is the same subspace of \mathbb{R}^3 that we studied in Examples 9 and 10 (see Exercises 14(b) and 14(c)). \blacksquare

New Vocabulary

linear combination (of vectors in a vector space)	subspace
proper subspace(s)	trivial subspace

Highlights

- A subset of a vector space is a subspace if it is a vector space itself under the same operations.
- The subset $\{0\}$ is a trivial subspace of any vector space.
- Any subspace of a vector space \mathcal{V} other than \mathcal{V} itself is considered a proper subspace.
- Familiar proper nontrivial subspaces of \mathbb{R}^3 are any line through the origin, any plane through the origin.
- Familiar proper subspaces of the real-valued functions on \mathbb{R} are $\mathcal{P}_n, \mathcal{P}$, all differentiable real-valued functions on \mathbb{R} , all continuous real-valued functions on \mathbb{R} .
- Familiar proper subspaces of \mathcal{M}_{nn} are $\mathcal{U}_n, \mathcal{L}_n, \mathcal{D}_n$, the symmetric $n \times n$ matrices, the skew-symmetric $n \times n$ matrices.
- A nonempty subset of a vector space is a subspace if it is closed under vector addition and scalar multiplication.
- If a subset of a vector space does not contain the zero vector, it cannot be a subspace.
- If a set of vectors is in a subspace, then any (finite) linear combination of those vectors is also in the subspace.
- If λ is an eigenvalue for an $n \times n$ matrix \mathbf{A} , then E_λ (eigenspace for λ) is a subspace of \mathbb{R}^n .
- The intersection of subspaces is a subspace.

EXERCISES FOR SECTION 4.2

Note: From this point onward in the book, use a calculator or available software packages to avoid tedious calculations.

1. Prove or disprove that each given subset of \mathbb{R}^2 is a subspace of \mathbb{R}^2 under the usual vector operations. (In these problems, a and b represent arbitrary real numbers.)
 - ★(a) The set of unit 2-vectors
 - (b) The set of 2-vectors of the form $[1, a]$

- ★(c) The set of 2-vectors of the form $[a, 2a]$
 - (d) The set of 2-vectors having a zero in at least one coordinate
 - ★(e) The set $\{[1, 2]\}$
 - (f) The set of 2-vectors whose second coordinate is zero
 - ★(g) The set of 2-vectors of the form $[a, b]$, where $|a| = |b|$
 - (h) The set of points in the plane lying on the line $y = -3x$
 - (i) The set of points in the plane lying on the line $y = 7x - 5$
 - ★(j) The set of points lying on the parabola $y = x^2$
 - (k) The set of points in the plane lying above the line $y = 2x - 5$
 - ★(l) The set of points in the plane lying inside the circle of radius 1 centered at the origin
2. Prove or disprove that each given subset of \mathcal{M}_{22} is a subspace of \mathcal{M}_{22} under the usual matrix operations. (In these problems, a and b represent arbitrary real numbers.)
- ★(a) The set of matrices of the form $\begin{bmatrix} a & -a \\ b & 0 \end{bmatrix}$
 - (b) The set of 2×2 matrices that have at least one row of zeroes
 - ★(c) The set of symmetric 2×2 matrices
 - (d) The set of nonsingular 2×2 matrices
 - ★(e) The set of 2×2 matrices having the sum of all entries zero
 - (f) The set of 2×2 matrices having trace zero (Recall that the *trace* of a square matrix is the sum of the main diagonal entries.)
 - ★(g) The set of 2×2 matrices \mathbf{A} such that $\mathbf{A} \begin{bmatrix} 1 & 3 \\ -2 & -6 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
 - ★(h) The set of 2×2 matrices having the product of all entries zero
3. Prove or disprove that each given subset of \mathcal{P}_5 is a subspace of \mathcal{P}_5 under the usual operations.
- ★(a) $\{\mathbf{p} \in \mathcal{P}_5 \mid \text{the coefficient of the first-degree term of } \mathbf{p} \text{ equals the coefficient of the fifth-degree term of } \mathbf{p}\}$
 - ★(b) $\{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p}(3) = 0\}$
 - (c) $\{\mathbf{p} \in \mathcal{P}_5 \mid \text{the sum of the coefficients of } \mathbf{p} \text{ is zero}\}$
 - (d) $\{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p}(3) = \mathbf{p}(5)\}$
 - ★(e) $\{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p} \text{ is an odd-degree polynomial (highest-order nonzero term has odd degree)}\}$
 - (f) $\{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p} \text{ has a relative maximum at } x = 0\}$
 - ★(g) $\{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p}'(4) = 0, \text{ where } \mathbf{p}' \text{ is the derivative of } \mathbf{p}\}$
 - (h) $\{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p}'(4) = 1, \text{ where } \mathbf{p}' \text{ is the derivative of } \mathbf{p}\}$

4. Show that the set of vectors of the form $[a, b, 0, c, a - 2b + c]$ in \mathbb{R}^5 forms a subspace of \mathbb{R}^5 under the usual operations.
5. Show that the set of vectors of the form $[2a - 3b, a - 5c, a, 4c - b, c]$ in \mathbb{R}^5 forms a subspace of \mathbb{R}^5 under the usual operations.
6. (a) Prove that the set of all 3-vectors orthogonal to $[1, -1, 4]$ forms a subspace of \mathbb{R}^3 .
 (b) Is the subspace from part (a) all of \mathbb{R}^3 , a plane passing through the origin in \mathbb{R}^3 , or a line passing through the origin in \mathbb{R}^3 ?
7. Show that each of the following sets is a subspace of the vector space of all real-valued functions on the given domain, under the usual operations of function addition and scalar multiplication:
 - (a) The set of continuous real-valued functions with domain \mathbb{R}
 - (b) The set of differentiable real-valued functions with domain \mathbb{R}
 - (c) The set of all real-valued functions \mathbf{f} defined on the interval $[0, 1]$ such that $\mathbf{f}(\frac{1}{2}) = 0$ (Compare this vector space with the set in Example 9 of Section 4.1.)
 - (d) The set of all continuous real-valued functions \mathbf{f} defined on the interval $[0, 1]$ such that $\int_0^1 \mathbf{f}(x) dx = 0$
8. Let \mathcal{W} be the set of differentiable real-valued functions $y = \mathbf{f}(x)$ defined on \mathbb{R} that satisfy the differential equation $3(dy/dx) - 2y = 0$. Show that, under the usual function operations, \mathcal{W} is a subspace of the vector space of all differentiable real-valued functions. (Do not forget to show \mathcal{W} is nonempty.)
9. Show that the set \mathcal{W} of solutions to the differential equation $y'' + 2y' - 9y = 0$ is a subspace of the vector space of all twice-differentiable real-valued functions defined on \mathbb{R} . (Do not forget to show that \mathcal{W} is nonempty.)
10. Prove that the set of discontinuous real-valued functions defined on \mathbb{R} (for example, $\mathbf{f}(x) = \begin{cases} 0 & \text{if } x \leq 0 \\ 1 & \text{if } x > 0 \end{cases}$) with the usual function operations is not a subspace of the vector space of all real-valued functions with domain \mathbb{R} .
11. Let \mathbf{A} be a fixed $n \times n$ matrix, and let \mathcal{W} be the subset of \mathcal{M}_{nn} of all $n \times n$ matrices that commute with \mathbf{A} under multiplication (that is, $\mathbf{B} \in \mathcal{W}$ if and only if $\mathbf{AB} = \mathbf{BA}$). Show that \mathcal{W} is a subspace of \mathcal{M}_{nn} under the usual vector space operations. (Do not forget to show that \mathcal{W} is nonempty.)
12. (a) A careful reading of the proof of Theorem 4.2 reveals that only closure under scalar multiplication (not closure under addition) is sufficient to prove the remaining eight vector space properties for \mathcal{W} . Explain, nevertheless, why closure under addition is a necessary condition for \mathcal{W} to be a subspace of \mathcal{V} .
 (b) Show that the set of singular $n \times n$ matrices is closed under scalar multiplication in \mathcal{M}_{nn} .

- (c) Use parts (a) and (b) to determine which of the eight vector space properties are true for the set of singular $n \times n$ matrices.
 - (d) Show that the set of singular $n \times n$ matrices is not closed under vector addition and hence is not a subspace of \mathcal{M}_{nn} ($n \geq 2$).
 - ★(e) Is the set of nonsingular $n \times n$ matrices closed under scalar multiplication? Why or why not?
13. (a) Prove that the set of all points lying on a line passing through the origin in \mathbb{R}^2 is a subspace of \mathbb{R}^2 (under the usual operations).
 (b) Prove that the set of all points in \mathbb{R}^2 lying on a line not passing through the origin does not form a subspace of \mathbb{R}^2 (under the usual operations).
 14. Let \mathcal{W} be the subspace from Examples 9 and 10, and let \mathbf{A} and E_{15} be as given in Example 11.
 (a) Use Theorem 4.2 to prove directly that E_{15} is a subspace of \mathbb{R}^3 .
 (b) Show that $E_{15} \subseteq \mathcal{W}$ by proving that every vector in E_{15} has the form $[a, b, \frac{1}{2}a - 2b]$.
 (c) Prove that $\mathcal{W} \subseteq E_{15}$ by showing that every nonzero vector of the form $[a, b, \frac{1}{2}a - 2b]$ is an eigenvector for \mathbf{A} corresponding to $\lambda_2 = 15$.
 - ★15. Suppose \mathbf{A} is an $n \times n$ matrix and $\lambda \in \mathbb{R}$ is *not* an eigenvalue for \mathbf{A} . Determine exactly which vectors are in $S = \{\mathbf{X} \in \mathbb{R}^n \mid \mathbf{A}\mathbf{X} = \lambda\mathbf{X}\}$. Is this set a subspace of \mathbb{R}^n ? Why or why not?
 16. Prove that \mathbb{R} (under the usual operations) has no subspaces except \mathbb{R} and $\{\mathbf{0}\}$. (Hint: Let \mathcal{V} be a nontrivial subspace of \mathbb{R} , and show that $\mathcal{V} = \mathbb{R}$.)
 17. Let \mathcal{W} be a subspace of a vector space \mathcal{V} . Show that the set $\mathcal{W}' = \{\mathbf{v} \in \mathcal{V} \mid \mathbf{v} \notin \mathcal{W}\}$ is not a subspace of \mathcal{V} .
 18. Let \mathcal{V} be a vector space, and let \mathcal{W}_1 and \mathcal{W}_2 be subspaces of \mathcal{V} . Prove that $\mathcal{W}_1 \cap \mathcal{W}_2$ is a subspace of \mathcal{V} . (Do not forget to show $\mathcal{W}_1 \cap \mathcal{W}_2$ is nonempty.)
 19. Let \mathcal{V} be any vector space, and let \mathcal{W} be a nonempty subset of \mathcal{V} .
 (a) Prove that \mathcal{W} is a subspace of \mathcal{V} if and only if $a\mathbf{w}_1 + b\mathbf{w}_2$ is an element of \mathcal{W} for every $a, b \in \mathbb{R}$ and every $\mathbf{w}_1, \mathbf{w}_2 \in \mathcal{W}$. (Hint: For one half of the proof, first consider the case where $a = b = 1$ and then the case where $b = 0$ and a is arbitrary.)
 (b) Prove that \mathcal{W} is a subspace of \mathcal{V} if and only if $a\mathbf{w}_1 + \mathbf{w}_2$ is an element of \mathcal{W} for every real number a and every \mathbf{w}_1 and \mathbf{w}_2 in \mathcal{W} .
 20. Let \mathcal{W} be a nonempty subset of a vector space \mathcal{V} , and suppose every linear combination of vectors in \mathcal{W} is also in \mathcal{W} . Prove that \mathcal{W} is a subspace of \mathcal{V} . (This is the converse of Theorem 4.3.)

21. Let λ be an eigenvalue for an $n \times n$ matrix \mathbf{A} . Show that if $\mathbf{X}_1, \dots, \mathbf{X}_k$ are eigenvectors for \mathbf{A} corresponding to λ , then any linear combination of $\mathbf{X}_1, \dots, \mathbf{X}_k$ is in E_λ .
- ★22. True or False:
- (a) A nonempty subset \mathcal{W} of a vector space \mathcal{V} is always a subspace of \mathcal{V} under the same operations as those in \mathcal{V} .
 - (b) Every vector space has at least one subspace.
 - (c) Any plane \mathcal{W} in \mathbb{R}^3 is a subspace of \mathbb{R}^3 (under the usual operations).
 - (d) The set of all lower triangular 5×5 matrices is a subspace of \mathcal{M}_{55} (under the usual operations).
 - (e) The set of all vectors of the form $[0, a, b, 0]$ is a subspace of \mathbb{R}^4 (under the usual operations).
 - (f) If a subset \mathcal{W} of a vector space \mathcal{V} contains the zero vector $\mathbf{0}$ of \mathcal{V} , then \mathcal{W} must be a subspace of \mathcal{V} (under the same operations).
 - (g) Any linear combination of vectors from a subspace \mathcal{W} of a vector space \mathcal{V} must also be in \mathcal{W} .
 - (h) If λ is an eigenvalue for a 4×4 matrix \mathbf{A} , then E_λ is a subspace of \mathbb{R}^4 .

4.3 SPAN

In this section, we study the concept of linear combinations in more depth. We show that the set of all linear combinations of the vectors in a subset S of \mathcal{V} forms an important subspace of \mathcal{V} , called the span of S in \mathcal{V} .

Finite Linear Combinations

In Section 4.2, we introduced linear combinations of vectors in a general vector space. We now extend the concept of linear combination to include the possibility of forming sums of scalar multiples from infinite, as well as finite, sets.

Definition Let S be a nonempty (possibly infinite) subset of a vector space \mathcal{V} . Then a vector \mathbf{v} in \mathcal{V} is a **(finite) linear combination of the vectors in S** if and only if there exists a *finite* subset $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ of S such that $\mathbf{v} = a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n$ for some real numbers a_1, \dots, a_n .

Examples 1 and 2 below involve a finite set S , while Examples 3 and 4 use an infinite set S . In all these examples, however, only a *finite* number of vectors from S are used at any given time to form linear combinations.

Example 1

Consider the subset $S = \{[1, -1, 0], [1, 0, 2], [0, -2, 5]\}$ of \mathbb{R}^3 . The vector $[1, -2, -2]$ is a linear combination of the vectors in S according to the definition, because $[1, -2, -2] = 2[1, -1, 0] + (-1)[1, 0, 2]$. In this case, the (finite) subset of S used (from the definition) is $\{[1, -1, 0], [1, 0, 2]\}$. However, we could have used all of S to form the linear combination by placing a zero coefficient in front of the remaining vector $[0, -2, 5]$. That is, $[1, -2, -2] = 2[1, -1, 0] + (-1)[1, 0, 2] + 0[0, -2, 5]$.

We see from Example 1 that if S is a *finite* subset of a vector space \mathcal{V} , any linear combination \mathbf{v} formed using *some* of the vectors in S can always be formed using *all* the vectors in S by placing zero coefficients on the remaining vectors.

A linear combination formed from a set $\{\mathbf{v}\}$ containing a single vector is just a scalar multiple $a\mathbf{v}$ of \mathbf{v} , as we see in the next example.

Example 2

Let $S = \{[1, -2, 7]\}$, a subset of \mathbb{R}^3 containing a single element. Then the only linear combinations that can be formed from S are scalar multiples of $[1, -2, 7]$, such as $[3, -6, 21]$ and $[-4, 8, -28]$.

Example 3

Consider \mathcal{P} , the vector space of polynomials with real coefficients, and let $S = \{1, x^2, x^4, \dots\}$, the infinite subset of \mathcal{P} consisting of all nonnegative even powers of x (since $x^0 = 1$). We can form linear combinations of vectors in S using any finite subset of S . For example, $\mathbf{p}(x) = 7x^8 - (1/4)x^4 + 10$ is a linear combination formed from S because it is a sum of scalar multiples of elements of a finite subset $\{x^8, x^4, 1\}$ of S . In fact, the possible linear combinations of vectors in S are precisely the polynomials involving only even powers of x .

Notice that we cannot use all of the elements in an infinite set S when forming a linear combination because an “infinite” sum would result. This is why a linear combination is frequently called a *finite* linear combination in order to stress that only a finite number of vectors are combined at any time.

Example 4

Let $S = \mathcal{U}_2 \cup \mathcal{L}_2$, an infinite subset of \mathcal{M}_{22} . (Recall that \mathcal{U}_2 and \mathcal{L}_2 are, respectively, the sets of upper and lower triangular 2×2 matrices.) The matrix $\mathbf{A} = \begin{bmatrix} 2 & 3 \\ -1 & \frac{1}{2} \end{bmatrix}$ is a linear combination of the elements in S , because

$$\mathbf{A} = \underbrace{\frac{1}{2} \begin{bmatrix} 4 & 6 \\ 0 & 1 \end{bmatrix}}_{\text{in } \mathcal{U}_2} + (-1) \underbrace{\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}}_{\text{in } \mathcal{L}_2}.$$

But there are many other ways to express \mathbf{A} as a finite linear combination of the elements in S . We can add more elements from S with zero coefficients, as in Example 1, but in this case there are further possibilities. For example,

$$\mathbf{A} = 2 \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}}_{\text{in } \mathcal{U}_2 \text{ and } \mathcal{L}_2} + 3 \underbrace{\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}}_{\text{in } \mathcal{U}_2} + (-1) \underbrace{\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}}_{\text{in } \mathcal{L}_2} + \frac{1}{2} \underbrace{\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}}_{\text{in } \mathcal{U}_2 \text{ and } \mathcal{L}_2}.$$

Definition of the Span of a Set

Definition Let S be a nonempty subset of a vector space \mathcal{V} . Then the **span** of S in \mathcal{V} is the set of all possible (finite) linear combinations of the vectors in S . We use the notation $\text{span}(S)$ to denote the span of S in \mathcal{V} .

The span of a set S is a generalization of the row space of a matrix; each is just the set of all linear combinations of a set of vectors. In fact, from this definition:

The span of the set of rows of a matrix is precisely the row space of the matrix.

We now consider some examples of the span of a subset.

Example 5

In Example 3, we found that for $S = \{1, x^2, x^4, \dots\}$ in \mathcal{P} , $\text{span}(S)$ is the set of all polynomials containing only even-degree terms. This consists of all the “destinations” obtainable by traveling in the “directions” $1, x^2, x^4, \dots$, etc. Thus, we can visualize $\text{span}(S)$ as the set of “possible destinations” in the same sense as the row space is the set of “possible destinations” obtainable from the rows of a given matrix. Notice that we may only use a finite number of the possible “directions” to obtain a given “destination.” That is, $\text{span}(S)$ only contains polynomials, not infinite series.

Example 6

Let $S = \mathcal{U}_2 \cup \mathcal{L}_2$ in \mathcal{M}_{22} , as in Example 4. Then $\text{span}(S) = \mathcal{M}_{22}$ because every 2×2 matrix can be expressed as a finite linear combination of upper and lower triangular matrices, as follows:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} + d \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Notice that the span of a given set often (but not always) contains many more vectors than the set itself.

Example 6 shows that, when $S = \mathcal{U}_2 \cup \mathcal{L}_2$ and $\mathcal{V} = \mathcal{M}_{22}$, every vector in \mathcal{V} is a linear combination of vectors in S . That is, $\text{span}(S) = \mathcal{V}$ itself. When this happens, we say that \mathcal{V} is **spanned by** S or that S **spans** \mathcal{V} . Here, we are using *span* as a *verb* to indicate that the *span* (noun) of a set S equals \mathcal{V} . Thus, \mathcal{M}_{22} is *spanned* (verb) by $\mathcal{U}_2 \cup \mathcal{L}_2$, since the *span* (noun) of $\mathcal{U}_2 \cup \mathcal{L}_2$ is \mathcal{M}_{22} .

Example 7

Note that \mathbb{R}^3 is spanned by $S_1 = \{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$, since $\text{span}(S_1) = \mathbb{R}^3$. That is, every 3-vector can be expressed as a linear combination of \mathbf{i}, \mathbf{j} , and \mathbf{k} (why?). However, \mathbb{R}^3 is not spanned by the smaller set $S_2 = \{\mathbf{i}, \mathbf{j}\}$, since $\text{span}(S_2)$ is the xy -plane in \mathbb{R}^3 (why?). More generally, \mathbb{R}^n is spanned by the set of standard unit vectors $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$. Note that no proper subset of $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ will span \mathbb{R}^n .

Span(S) Is the Minimal Subspace Containing S

The next theorem completely characterizes the span.

Theorem 4.5 Let S be a nonempty subset of a vector space \mathcal{V} . Then:

- (1) $S \subseteq \text{span}(S)$.
- (2) $\text{Span}(S)$ is a subspace of \mathcal{V} (under the same operations as \mathcal{V}).
- (3) If \mathcal{W} is a subspace of \mathcal{V} with $S \subseteq \mathcal{W}$, then $\text{span}(S) \subseteq \mathcal{W}$.
- (4) $\text{Span}(S)$ is the smallest subspace of \mathcal{V} containing S .

Proof. Part (1): We must show that each vector $\mathbf{w} \in S$ is also in $\text{span}(S)$. But if $\mathbf{w} \in S$, then $\mathbf{w} = 1\mathbf{w}$ is a sum of scalar multiples from the subset $\{\mathbf{w}\}$ of S . Hence, $\mathbf{w} \in \text{span}(S)$.

Part (2): Since S is nonempty, part (1) shows that $\text{span}(S)$ is nonempty. Therefore, by Theorem 4.2, $\text{span}(S)$ is a subspace of \mathcal{V} if we can prove the closure properties hold for $\text{span}(S)$.

First, let us verify closure under scalar multiplication. Let \mathbf{v} be in $\text{span}(S)$, and let c be a scalar. We must show that $c\mathbf{v} \in \text{span}(S)$. Now, since $\mathbf{v} \in \text{span}(S)$, a finite subset $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ of S and real numbers a_1, \dots, a_n exist such that $\mathbf{v} = \sum_{i=1}^n a_i \mathbf{v}_i$. Then,

$$c\mathbf{v} = c(a_1 \mathbf{v}_1 + \dots + a_n \mathbf{v}_n) = (ca_1) \mathbf{v}_1 + \dots + (ca_n) \mathbf{v}_n.$$

Hence, $c\mathbf{v}$ is a linear combination of the finite subset $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ of S , and so $c\mathbf{v} \in \text{span}(S)$.

Finally, we show that $\text{span}(S)$ is closed under vector addition. First we will consider the case in which S is a finite set. Thus, we suppose that $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$. Let \mathbf{x} and \mathbf{y} be two vectors in $\text{span}(S)$. Hence, there exist real numbers a_1, \dots, a_n and b_1, \dots, b_n such that

$$\mathbf{x} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_n \mathbf{v}_n \text{ and } \mathbf{y} = b_1 \mathbf{v}_1 + b_2 \mathbf{v}_2 + \dots + b_n \mathbf{v}_n.$$

Therefore,

$$\mathbf{x} + \mathbf{y} = (a_1 + b_1) \mathbf{v}_1 + (a_2 + b_2) \mathbf{v}_2 + \dots + (a_n + b_n) \mathbf{v}_n,$$

and we have expressed $\mathbf{x} + \mathbf{y}$ as a linear combination of vectors in S . Hence, $\mathbf{x} + \mathbf{y} \in \text{span}(S)$.

The proof of closure under addition in the case in which S has an infinite number of elements is identical in concept to the finite case. However, the linear combinations for \mathbf{x} and \mathbf{y} might now be formed using two different finite subsets of vectors from S . This complication is remedied by uniting these two subsets into one common finite subset of S that we use to form the linear combinations for \mathbf{x} and \mathbf{y} . Then we place a coefficient of zero in front of any vector in the union that is unneeded when forming the desired linear combination for \mathbf{x} , and similarly for \mathbf{y} . You are asked to complete the details for this part of the proof in Exercise 28.

Part (3): This part asserts that if S is a subset of a subspace \mathcal{W} , then any (finite) linear combination from S is also in \mathcal{W} . This is merely a rewording of Theorem 4.3 using the “span” concept. The fact that $\text{span}(S)$ cannot contain vectors outside \mathcal{W} is illustrated in Figure 4.4.

Part (4): This is merely a summary of the other three parts. Parts (1) and (2) assert that $\text{span}(S)$ is a subspace of \mathcal{V} containing S . But part (3) shows that $\text{span}(S)$ is the smallest such subspace because $\text{span}(S)$ must be a subset of, and hence smaller than, any other subspace of \mathcal{V} that contains S . \square

Theorem 4.5 implies that $\text{span}(S)$ is created by appending to S precisely those vectors needed to make the closure properties hold. In fact, the whole idea behind span is to “close up” a subset of a vector space to create a subspace.

Example 8

Let \mathbf{v}_1 and \mathbf{v}_2 be any two vectors in \mathbb{R}^4 . Then, by Theorem 4.5, $\text{span}(\{\mathbf{v}_1, \mathbf{v}_2\})$ is the smallest subspace of \mathbb{R}^4 containing \mathbf{v}_1 and \mathbf{v}_2 . In particular, if $\mathbf{v}_1 = [1, 3, -2, 5]$ and $\mathbf{v}_2 = [0, -4, 3, -1]$, then $\text{span}(\{\mathbf{v}_1, \mathbf{v}_2\})$ is the subspace of \mathbb{R}^4 consisting of all vectors of the form

$$a[1, 3, -2, 5] + b[0, -4, 3, -1] = [a, 3a - 4b, -2a + 3b, 5a - b].$$

No smaller subspace of \mathbb{R}^4 contains \mathbf{v}_1 and \mathbf{v}_2 . \blacksquare

The following useful result is left for you to prove in Exercise 21.




FIGURE 4.4

(a) Situation that *must* occur if \mathcal{W} is a subspace containing S ; (b) situation that *cannot* occur if \mathcal{W} is a subspace containing S

Corollary 4.6 Let \mathcal{V} be a vector space, and let S_1 and S_2 be subsets of \mathcal{V} with $S_1 \subseteq S_2$. Then $\text{span}(S_1) \subseteq \text{span}(S_2)$.

Simplifying $\text{Span}(S)$ using Row Reduction

Our next goal is to find a simplified form for the vectors in the span of a given set S . The fact that span is a generalization of the row space concept suggests that we can use results from Chapter 2 involving row spaces to help us compute and simplify $\text{span}(S)$. If we form the matrix \mathbf{A} whose rows are the vectors in S , the rows of the reduced row echelon form of \mathbf{A} often give a simpler expression for $\text{span}(S)$, since row equivalent matrices have the same row space. Hence, we have the following:

Method for Simplifying $\text{Span}(S)$ Using Row Reduction (Simplified Span Method)

Suppose that S is a finite subset of \mathbb{R}^n containing k vectors, with $k \geq 2$.

Step 1: Form a $k \times n$ matrix \mathbf{A} by using the vectors in S as the rows of \mathbf{A} . (Thus, $\text{span}(S)$ is the row space of \mathbf{A}).

Step 2: Let \mathbf{C} be the reduced row echelon form matrix for \mathbf{A} .

Step 3: Then, a simplified form for $\text{span}(S)$ is given by the set of all linear combinations of the *nonzero* rows of \mathbf{C} .

Example 9

Let S be the subset $\{[1, 4, -1, -5], [2, 8, 5, 4], [-1, -4, 2, 7], [6, 24, -1, -20]\}$ of \mathbb{R}^4 . By definition, $\text{span}(S)$ is the set of all vectors of the form

$$a[1, 4, -1, -5] + b[2, 8, 5, 4] + c[-1, -4, 2, 7] + d[6, 24, -1, -20]$$

for $a, b, c, d \in \mathbb{R}$. We want to use the Simplified Span Method to find a simplified form for the vectors in $\text{span}(S)$. We first create

$$\mathbf{A} = \begin{bmatrix} 1 & 4 & -1 & -5 \\ 2 & 8 & 5 & 4 \\ -1 & -4 & 2 & 7 \\ 6 & 24 & -1 & -20 \end{bmatrix},$$

whose rows are the vectors in S . Then, $\text{span}(S)$ is the row space of \mathbf{A} ; that is, the set of all linear combinations of the rows of \mathbf{A} .

Next, we simplify the form of the row space of \mathbf{A} by obtaining its reduced row echelon form matrix

$$\mathbf{C} = \begin{bmatrix} 1 & 4 & 0 & -3 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

By Theorem 2.8, the row space of \mathbf{A} is the same as the row space of \mathbf{C} , which is the set of all 4-vectors of the form

$$a[1, 4, 0, -3] + b[0, 0, 1, 2] = [a, 4a, b, -3a + 2b].$$

Therefore, $\text{span}(S) = \{[a, 4a, b, -3a + 2b] \mid a, b \in \mathbb{R}\}$, a subspace of \mathbb{R}^4 . Note, for example, that the vector $[3, 12, -2, -13]$ is in $\text{span}(S)$ ($a = 3, b = -2$). However, the vector $[-2, -8, 4, 6]$ is not in $\text{span}(S)$ because the following system has no solutions:

$$\begin{cases} a = -2 \\ 4a = -8 \\ b = 4 \\ -3a + 2b = 6 \end{cases}.$$

Example 10

Recall that the eigenspace E_{15} for the matrix \mathbf{A} in Example 11 in Section 4.2 is $E_{15} = \{a[4, 1, 0] + b[2, 0, 1] \mid a, b \in \mathbb{R}\}$. Hence, E_{15} is spanned by $\{[4, 1, 0], [2, 0, 1]\}$. Although the form of E_{15} is already simple, we can obtain an alternative form by using the Simplified Span Method. Row reducing the matrix

$$\mathbf{A} = \begin{bmatrix} 4 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}, \quad \text{we obtain} \quad \mathbf{C} = \begin{bmatrix} 1 & 0 & \frac{1}{2} \\ 0 & 1 & -2 \end{bmatrix}.$$

Hence, an alternative form for the vectors in E_{15} is $\left\{a\left[1, 0, \frac{1}{2}\right] + b[0, 1, -2] \mid a, b \in \mathbb{R}\right\} = \left\{[a, b, \frac{1}{2}a - 2b] \mid a, b \in \mathbb{R}\right\}$, just as we claimed in Example 11 in Section 4.2.

The method used in Examples 9 and 10 works in vector spaces other than \mathbb{R}^n , as we see in the next example. This fact will follow from the discussion of isomorphism in Section 5.5. (However, we will not use this fact in proofs of theorems until after Section 5.5.)

Example 11

Let S be the subset $\{5x^3 + 2x^2 + 4x - 3, -x^2 + 3x - 7, 2x^3 + 4x^2 - 8x + 5, x^3 + 2x + 5\}$ of \mathcal{P}_3 . We use the Simplified Span Method to find a simplified form for the vectors in $\text{span}(S)$.

Consider the coefficients of each polynomial as the coordinates of a vector in \mathbb{R}^4 , yielding the corresponding set of vectors $T = \{[5, 2, 4, -3], [0, -1, 3, -7], [2, 4, -8, 5], [1, 0, 2, 5]\}$. Using the Simplified Span Method, we create the following matrix, whose rows are the vectors in T .

$$\mathbf{A} = \begin{bmatrix} 5 & 2 & 4 & -3 \\ 0 & -1 & 3 & -7 \\ 2 & 4 & -8 & 5 \\ 1 & 0 & 2 & 5 \end{bmatrix}$$

Then $\text{span}(T)$ is the row space of the reduced row echelon form of \mathbf{A} , which is

$$\mathbf{C} = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Taking each nonzero row of \mathbf{C} as the coefficients of a polynomial in \mathcal{P}_3 , we see that

$$\begin{aligned} \text{span}(S) &= \{a(x^3 + 2x) + b(x^2 - 3x) + c(1) \mid a, b, c \in \mathbb{R}\} \\ &= \{ax^3 + bx^2 + (2a - 3b)x + c \mid a, b, c \in \mathbb{R}\}. \end{aligned}$$

A Spanning Set for an Eigenspace

In Section 3.4, we illustrated a method for diagonalizing an $n \times n$ matrix, when possible. In fact, a set S of fundamental eigenvectors generated for a given eigenvalue λ spans the eigenspace E_λ (see Exercise 27). We illustrate this in the following example:

Example 12

Let

$$\mathbf{A} = \begin{bmatrix} 0 & -6 & 3 \\ 2 & -13 & 6 \\ 4 & -24 & 11 \end{bmatrix}.$$

A little work yields $p_{\mathbf{A}}(x) = x^3 + 2x^2 + x = x(x+1)^2$. We solve the homogeneous system $(-\mathbf{I}_3 - \mathbf{A})\mathbf{X} = \mathbf{0}$ to find the eigenspace E_{-1} for \mathbf{A} .

Row reducing $[(-\mathbf{I}_3 - \mathbf{A})|\mathbf{0}]$ produces

$$\left[\begin{array}{ccc|c} 1 & -6 & 3 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right],$$

giving the solution set

$$E_{-1} = \{[6b - 3c, b, c] \mid b, c \in \mathbb{R}\} = \{b[6, 1, 0] + c[-3, 0, 1] \mid b, c \in \mathbb{R}\}.$$

Thus, $E_{-1} = \text{span}(S)$, where $S = \{[6, 1, 0], [-3, 0, 1]\}$. The set S is precisely the set of fundamental eigenvectors that we would obtain for $\lambda = -1$ (verify!).

Special Case: The Span of the Empty Set

Until now, our results involving span have specified that the subset S of the vector space \mathcal{V} be nonempty. However, our understanding of $\text{span}(S)$ as the smallest subspace of \mathcal{V} containing S allows us to give a meaningful definition for the span of the empty set.

Definition $\text{Span}(\{\}) = \{\mathbf{0}\}$.

This definition makes sense because the trivial subspace is the smallest subspace of \mathcal{V} , hence the smallest one containing the empty set. Thus, Theorem 4.5 is also true when the set S is empty. Similarly, to maintain consistency, we *define* any linear combination of the empty set of vectors to be $\mathbf{0}$. This ensures that the span of the empty set equals the set of all linear combinations of vectors taken from this set.

New Vocabulary

finite linear combination (of vectors in a vector space)	span (of a set of vectors)
Simplified Span Method	spanned by (as in “ \mathcal{V} is spanned by S ”)
	span of the empty set

Highlights

- The span of a set is the collection of all finite linear combinations of vectors from the set.
- A set S spans a vector space \mathcal{V} (i.e., \mathcal{V} is spanned by S) if every vector in \mathcal{V} is a (finite) linear combination of vectors in S .
- The row space of a matrix is the span of the rows of the matrix.
- \mathbb{R}^3 is spanned by $\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$; \mathbb{R}^n is spanned by $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$; \mathcal{P}_n is spanned by $\{1, x, x^2, \dots, x^n\}$; \mathcal{M}_{mn} is spanned by $\{\Psi_{ij}\}$, where each Ψ_{ij} has a 1 in the (i, j) entry, and zeroes elsewhere.
- The span of a set of vectors is always a subspace of the vector space, and is, in fact, the smallest subspace containing that set.
- If $S_1 \subseteq S_2$, then $\text{span}(S_1) \subseteq \text{span}(S_2)$.
- The Simplified Span Method generally produces a more simplified form of the span of a set of vectors by calculating the reduced row echelon form of the matrix whose *rows* are the given vectors.
- The span of the empty set is $\{\mathbf{0}\}$.

EXERCISES FOR SECTION 4.3

- In each of the following cases, use the Simplified Span Method to find a simplified general form for all the vectors in $\text{span}(S)$, where S is the given subset of \mathbb{R}^n :
 - $S = \{[1, 1, 0], [2, -3, -5]\}$
 - $S = \{[3, 1, -2], [-3, -1, 2], [6, 2, -4]\}$
 - $S = \{[1, -1, 1], [2, -3, 3], [0, 1, -1]\}$
 - $S = \{[1, 1, 1], [2, 1, 1], [1, 1, 2]\}$
 - $S = \{[1, 3, 0, 1], [0, 0, 1, 1], [0, 1, 0, 1], [1, 5, 1, 4]\}$
 - $S = \{[2, -1, 3, 1], [1, -2, 0, -1], [3, -3, 3, 0], [5, -4, 6, 1], [1, -5, -3, -4]\}$
- In each case, use the Simplified Span Method to find a simplified general form for all the vectors in $\text{span}(S)$, where S is the given subset of \mathcal{P}_3 :
 - $S = \{x^3 - 1, x^2 - x, x - 1\}$
 - $S = \{x^3 + 2x^2, 1 - 4x^2, 12 - 5x^3, x^3 - x^2\}$
 - $S = \{x^3 - x + 5, 3x^3 - 3x + 10, 5x^3 - 5x - 6, 6x - 6x^3 - 13\}$
- In each case, use the Simplified Span Method to find a simplified general form for all the vectors in $\text{span}(S)$, where S is the given subset of \mathcal{M}_{22} . (Hint: Rewrite each matrix as a 4-vector.)
 - $S = \left\{ \begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \right\}$
 - $S = \left\{ \begin{bmatrix} 1 & 3 \\ -2 & 1 \end{bmatrix}, \begin{bmatrix} -2 & -5 \\ 3 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 4 \\ -3 & 4 \end{bmatrix} \right\}$
 - $S = \left\{ \begin{bmatrix} 1 & -1 \\ 3 & 0 \end{bmatrix}, \begin{bmatrix} 2 & -1 \\ 8 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 4 \\ 4 & -1 \end{bmatrix}, \begin{bmatrix} 3 & -4 \\ 5 & 6 \end{bmatrix} \right\}$
- Express the subspace \mathcal{W} of \mathbb{R}^4 of all 4-vectors of the form $[a + b, a + c, b + c, c]$ as the row space of a matrix \mathbf{A} .
 - Find the reduced row echelon form matrix \mathbf{B} for \mathbf{A} .
 - Use the matrix \mathbf{B} from part (b) to find a simplified form for the vectors in \mathcal{W} .
- Express the subspace \mathcal{W} of \mathbb{R}^5 of all 5-vectors of the form $[2a + 3b - 4c, a + b - c, -b + 7c, 3a + 4b, 4a + 2b]$ as the row space of a matrix \mathbf{A} .
 - Find the reduced row echelon form matrix \mathbf{B} for \mathbf{A} .
 - Use the matrix \mathbf{B} from part (b) to find a simplified form for the vectors in \mathcal{W} .
- Prove that the set $S = \{[1, 3, -1], [2, 7, -3], [4, 8, -7]\}$ spans \mathbb{R}^3 .
- Prove that the set $S = \{[1, -2, 2], [3, -4, -1], [1, -4, 9], [0, 2, -7]\}$ does not span \mathbb{R}^3 .

8. Show that the set $\{x^2 + x + 1, x + 1, 1\}$ spans \mathcal{P}_2 .
9. Prove that the set $\{x^2 + 4x - 3, 2x^2 + x + 5, 7x - 11\}$ does not span \mathcal{P}_2 .
10. (a) Let $S = \{[1, -2, -2], [3, -5, 1], [-1, 1, -5]\}$. Show that $[-4, 5, -13] \in \text{span}(S)$ by expressing it as a linear combination of the vectors in S .
 (b) Prove that the set S in part (a) does not span \mathbb{R}^3 .
- ★11. Consider the subset $S = \{x^3 - 2x^2 + x - 3, 2x^3 - 3x^2 + 2x + 5, 4x^2 + x - 3, 4x^3 - 7x^2 + 4x - 1\}$ of \mathcal{P} . Show that $3x^3 - 8x^2 + 2x + 16$ is in $\text{span}(S)$ by expressing it as a linear combination of the elements of S .
12. Prove that the set S of all vectors in \mathbb{R}^4 that have zeroes in exactly two coordinates spans \mathbb{R}^4 . (Hint: Find a subset of S that spans \mathbb{R}^4 .)
13. Let \mathbf{a} be any nonzero element of \mathbb{R} . Prove that $\text{span}(\{\mathbf{a}\}) = \mathbb{R}$.
14. ★(a) Suppose that S_1 is the set of symmetric 2×2 matrices and that S_2 is the set of skew-symmetric 2×2 matrices. Prove that $\text{span}(S_1 \cup S_2) = \mathcal{M}_{22}$.
 (b) State and prove the corresponding statement for $n \times n$ matrices.
15. Consider the subset $S = \{1 + x^2, x + x^3, 3 - 2x + 3x^2 - 12x^3\}$ of \mathcal{P} , and let $\mathcal{W} = \{ax^3 + bx^2 + cx + b \mid a, b, c \in \mathbb{R}\}$. Show that $\mathcal{W} = \text{span}(S)$.
16. Let $\mathbf{A} = \begin{bmatrix} -9 & -15 & 8 \\ -10 & -14 & 8 \\ -30 & -45 & 25 \end{bmatrix}$.
 ★(a) Find a set S of two fundamental eigenvectors for \mathbf{A} corresponding to the eigenvalue $\lambda = 1$. Multiply by a scalar to eliminate any fractions in your answers.
 (b) Verify that the set S from part (a) spans E_1 .
17. Let $S_1 = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a nonempty subset of a vector space \mathcal{V} . Let $S_2 = \{-\mathbf{v}_1, -\mathbf{v}_2, \dots, -\mathbf{v}_n\}$. Show that $\text{span}(S_1) = \text{span}(S_2)$.
18. Let \mathbf{u} and \mathbf{v} be two nonzero vectors in \mathbb{R}^3 , and let $S = \{\mathbf{u}, \mathbf{v}\}$. Show that $\text{span}(S)$ is a line through the origin if $\mathbf{u} = a\mathbf{v}$ for some real number a , but otherwise $\text{span}(S)$ is a plane through the origin.
19. Let \mathbf{u}, \mathbf{v} , and \mathbf{w} be three vectors in \mathbb{R}^3 and let \mathbf{A} be the matrix whose rows are \mathbf{u}, \mathbf{v} , and \mathbf{w} . Show that $S = \{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ spans \mathbb{R}^3 if and only if $|\mathbf{A}| \neq 0$. (Hint: To prove that $\text{span}(S) = \mathbb{R}^3$ implies $|\mathbf{A}| \neq 0$, suppose $\mathbf{x} \in \mathbb{R}^3$ such that $\mathbf{A}\mathbf{x} = \mathbf{0}$. First, show that \mathbf{x} is orthogonal to \mathbf{u}, \mathbf{v} , and \mathbf{w} . Then, express \mathbf{x} as a linear combination of \mathbf{u}, \mathbf{v} , and \mathbf{w} . Prove that $\mathbf{x} \cdot \mathbf{x} = 0$, and then use Theorem 2.5 and Corollary 3.6. To prove that $|\mathbf{A}| \neq 0$ implies $\text{span}(S) = \mathbb{R}^3$, show that \mathbf{A} is row equivalent to \mathbf{I}_3 and apply Theorem 2.8.)

20. Let $S = \{\mathbf{p}_1, \dots, \mathbf{p}_k\}$ be a finite subset of \mathcal{P} . Prove that there is some positive integer n such that $\text{span}(S) \subseteq \mathcal{P}_n$.
- 21. Prove Corollary 4.6.
22. (a) Prove that if S is a nonempty subset of a vector space \mathcal{V} , then S is a subspace of \mathcal{V} if and only if $\text{span}(S) = S$.
 (b) Use part (a) to show that every subspace \mathcal{W} of a vector space \mathcal{V} has a set of vectors that spans \mathcal{W} — namely, the set \mathcal{W} itself.
 (c) Describe the span of the set of the skew-symmetric matrices in \mathcal{M}_{33} .
23. Let S_1 and S_2 be subsets of a vector space \mathcal{V} . Prove that $\text{span}(S_1) = \text{span}(S_2)$ if and only if $S_1 \subseteq \text{span}(S_2)$ and $S_2 \subseteq \text{span}(S_1)$.
24. Let S_1 and S_2 be two subsets of a vector space \mathcal{V} .
 (a) Prove that $\text{span}(S_1 \cap S_2) \subseteq \text{span}(S_1) \cap \text{span}(S_2)$.
 ★(b) Give an example of distinct subsets S_1 and S_2 of \mathbb{R}^3 for which the inclusion in part (a) is actually an equality.
 ★(c) Give an example of subsets S_1 and S_2 of \mathbb{R}^3 for which the inclusion in part (a) is not an equality.
25. Let S_1 and S_2 be subsets of a vector space \mathcal{V} .
 (a) Show that $\text{span}(S_1) \cup \text{span}(S_2) \subseteq \text{span}(S_1 \cup S_2)$.
 (b) Prove that if $S_1 \subseteq S_2$, then the inclusion in part (a) is an equality.
 ★(c) Give an example of subsets S_1 and S_2 in \mathcal{P}_5 for which the inclusion in part (a) is not an equality.
26. Let S be a subset of a vector space \mathcal{V} , and let $\mathbf{v} \in \mathcal{V}$. Show that $\text{span}(S) = \text{span}(S \cup \{\mathbf{v}\})$ if and only if $\mathbf{v} \in \text{span}(S)$.
27. Let \mathbf{A} be an $n \times n$ matrix and λ be an eigenvalue for \mathbf{A} . Suppose S is a set of fundamental eigenvectors for \mathbf{A} corresponding to λ . Prove that S spans E_λ .
- 28. Finish the proof of Theorem 4.5 by providing the details necessary to show that $\text{span}(S)$ is closed under addition if S is an infinite subset of a vector space \mathcal{V} .
- ★29. True or False:
 (a) $\text{span}(S)$ is only defined if S is a finite subset of a vector space.
 (b) If S is a subset of a vector space \mathcal{V} , then $\text{span}(S)$ contains every finite linear combination of vectors in S .
 (c) If S is a subset of a vector space \mathcal{V} , then $\text{span}(S)$ is the smallest set in \mathcal{V} containing S .
 (d) If S is a subset of a vector space \mathcal{V} , and \mathcal{W} is a subspace of \mathcal{V} containing S , then we must have $\mathcal{W} \subseteq \text{span}(S)$.

- (e) The row space of a 4×5 matrix \mathbf{A} is a subspace of \mathbb{R}^4 .
- (f) A simplified form for the span of a finite set S of vectors in \mathbb{R}^n can be found by row reducing the matrix whose rows are the vectors of S .
- (g) The eigenspace E_λ for an eigenvalue λ of an $n \times n$ matrix \mathbf{A} is the row space of $\lambda \mathbf{I}_n - \mathbf{A}$.

4.4 LINEAR INDEPENDENCE

In this section, we explore the concept of a linearly independent set of vectors and examine methods for determining whether or not a given set of vectors is linearly independent. We will also see that there are important connections between the concepts of span and linear independence.

Linear Independence and Dependence

At first, we will define linear independence and linear dependence only for finite sets of vectors. We will extend the definition to infinite sets at the end of this section.

Definition Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a finite nonempty subset of a vector space \mathcal{V} . Then S is **linearly dependent** if and only if there exist real numbers a_1, \dots, a_n , not all zero, such that $a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n = \mathbf{0}$. That is, S is linearly dependent if and only if the zero vector can be expressed as a nontrivial linear combination of the vectors in S .

S is **linearly independent** if and only if it is *not* linearly dependent.

The empty set, $\{\}$, is linearly independent.

To understand this definition, we begin first with the simplest cases: sets having one or two elements.

Suppose $S = \{\mathbf{v}\}$, a one-element set. Then, by part (4) of Theorem 4.1, $a\mathbf{v} = \mathbf{0}$ implies that either $a = 0$ or $\mathbf{v} = \mathbf{0}$. Now, for S to be linearly dependent, we would have to have some nonzero a satisfy $a\mathbf{v} = \mathbf{0}$. This would imply that $\mathbf{v} = \mathbf{0}$. We conclude that if $S = \{\mathbf{v}\}$, a one-element set, then S is linearly dependent if and only if $\mathbf{v} = \mathbf{0}$. Equivalently, $S = \{\mathbf{v}\}$ is linearly independent if and only if $\mathbf{v} \neq \mathbf{0}$.

Example 1

Let $S_1 = \{[3, -1, 4]\}$. Since S_1 contains a single vector and this vector is nonzero, S_1 is a linearly independent subset of \mathbb{R}^3 . On the other hand, $S_2 = \{[0, 0, 0]\}$ is a linearly dependent subset of \mathbb{R}^4 .

Next, suppose $S = \{\mathbf{v}_1, \mathbf{v}_2\}$ is a linearly dependent set with two elements. Then there exist real numbers a_1 and a_2 , not both zero, such that $a_1\mathbf{v}_1 + a_2\mathbf{v}_2 = \mathbf{0}$. If $a_1 \neq 0$, this implies that $\mathbf{v}_1 = -\frac{a_2}{a_1}\mathbf{v}_2$. That is, \mathbf{v}_1 is a scalar multiple of \mathbf{v}_2 . Similarly, if $a_2 \neq 0$, we see that \mathbf{v}_2 is a scalar multiple of \mathbf{v}_1 . Thus, linearly dependent sets containing exactly two vectors are precisely those for which at least one of the vectors is a scalar multiple of the other. So, a set of exactly two vectors is linearly independent precisely when neither of the vectors is a scalar multiple of the other. That is, two linearly independent vectors are not parallel. They represent two different directions.

Example 2

The set of vectors $S_1 = \{[1, -1, 2], [-3, 3, -6]\}$ in \mathbb{R}^3 is linearly dependent since one of the vectors is a scalar multiple (and hence a linear combination) of the other. For example, $[1, -1, 2] = (-\frac{1}{3})[-3, 3, -6]$.

Also, the set $S_2 = \{[3, -8], [2, 5]\}$ is a linearly independent subset of \mathbb{R}^2 because neither of these vectors is a scalar multiple of the other. These two vectors are not parallel. They represent two different directions.

In general, because linear independence is defined as the negation of linear dependence, we can express linear independence as follows:

Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a finite nonempty subset of a vector space \mathcal{V} . Then S is **linearly independent** if and only if for any set of real numbers a_1, \dots, a_n , the equation $a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n = \mathbf{0}$ implies $a_1 = a_2 = \dots = a_n = 0$.

Example 3

The set of vectors $\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$ in \mathbb{R}^3 is linearly independent because $a\mathbf{i} + b\mathbf{j} + c\mathbf{k} = [a, b, c] = [0, 0, 0]$ if and only if $a = b = c = 0$. More generally, the set $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ in \mathbb{R}^n is linearly independent.

Example 4

Let S be any subset of a vector space \mathcal{V} containing the zero vector $\mathbf{0}$. If S contains no vector other than $\mathbf{0}$, then we have already seen that S is linearly dependent. If $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ contains at least two distinct vectors with one of them $\mathbf{0}$ (say $\mathbf{v}_1 = \mathbf{0}$), then $\mathbf{0}$ can be expressed as a nontrivial linear combination of the vectors in S since $1\mathbf{v}_1 + 0\mathbf{v}_2 + \dots + 0\mathbf{v}_n = 1 \cdot \mathbf{0} + \mathbf{0} + \dots + \mathbf{0} = \mathbf{0}$. Hence, by the definition, S is linearly dependent. Therefore, in all cases, any finite subset of a vector space that contains the zero vector $\mathbf{0}$ is linearly dependent.

The result we obtained in Example 4 is important enough to highlight:

Any finite subset of a vector space that contains the zero vector $\mathbf{0}$ is linearly dependent.

Example 5

Let $S = \{[2, 5], [3, -2], [4, -9]\}$. Notice that $[4, -9] = -[2, 5] + 2[3, -2]$. This shows that some vector in S can be expressed as a linear combination of other vectors in S . In other words, the vector $[4, -9]$ is a “destination” that can be reached using a linear combination of the other vectors in S . It does not strike out in a new, independent, direction. Notice that we can subtract $[4, -9]$ from both sides of the equation $[4, -9] = -[2, 5] + 2[3, -2]$ to obtain

$$\mathbf{0} = -[2, 5] + 2[3, -2] - [4, -9].$$

We have thus expressed the zero vector as a nontrivial linear combination of the vectors in S , and this implies that S is linearly dependent.

Example 6

Consider the subset $S = \{[1, -1, 0, 2], [0, -2, 1, 0], [2, 0, -1, 1]\}$ of \mathbb{R}^4 . We will investigate whether S is linearly independent.

We proceed by assuming that $a[1, -1, 0, 2] + b[0, -2, 1, 0] + c[2, 0, -1, 1] = [0, 0, 0, 0]$ and solve for a, b , and c to see whether all these coefficients must be zero. That is, we determine whether the following homogeneous system has only the trivial solution:

$$\begin{cases} a & + 2c = 0 \\ -a - 2b & = 0 \\ & b - c = 0 \\ 2a & + c = 0 \end{cases}.$$

Row reducing

$$\left[\begin{array}{ccc|c} a & b & c & 0 \\ 1 & 0 & 2 & 0 \\ -1 & -2 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 2 & 0 & 1 & 0 \end{array} \right], \quad \text{we obtain} \quad \left[\begin{array}{ccc|c} a & b & c & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right],$$

which shows that this system has only the trivial solution $a = b = c = 0$. Hence, S is linearly independent.

Using Row Reduction to Test for Linear Independence

Notice that in Example 6, the columns of the matrix to the left of the augmentation bar are just the vectors in S . In general, to test a finite set of vectors in \mathbb{R}^n for linear independence, we row reduce the matrix whose *columns* are the vectors in the set, and then check whether the associated homogeneous system has only the trivial solution. In practice it is not necessary to include the augmentation bar and the column of zeroes to its right, since this column never changes in the row reduction process. Thus, we have

Method to Test for Linear Independence Using Row Reduction (Independence Test Method)

Let S be a finite nonempty set of vectors in \mathbb{R}^n . To determine whether S is linearly independent, perform the following steps:

Step 1: Create the matrix \mathbf{A} whose *columns* are the vectors in S .

Step 2: Find \mathbf{B} , the reduced row echelon form of \mathbf{A} .

Step 3: If there is a pivot in every column of \mathbf{B} , then S is linearly independent. Otherwise, S is linearly dependent.

Example 7

Consider the subset $S = \{[3, 1, -1], [-5, -2, 2], [2, 2, -1]\}$ of \mathbb{R}^3 . Using the Independence Test Method, we row reduce

$$\begin{bmatrix} 3 & -5 & 2 \\ 1 & -2 & 2 \\ -1 & 2 & -1 \end{bmatrix} \text{ to obtain } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Since we found a pivot in every column, the set S is linearly independent. ■

Example 8

Consider the subset $S = \{[2, 5], [3, 7], [4, -9], [-8, 3]\}$ of \mathbb{R}^2 . Using the Independence Test Method, we row reduce

$$\begin{bmatrix} 2 & 3 & 4 & -8 \\ 5 & 7 & -9 & 3 \end{bmatrix} \text{ to obtain } \begin{bmatrix} 1 & 0 & -55 & 65 \\ 0 & 1 & 38 & -46 \end{bmatrix}.$$

Since we have no pivots in columns 3 and 4, the set S is linearly dependent. ■

In the last example, there are more columns than rows in the matrix we row reduced. Hence, there must ultimately be some column without a pivot, since each pivot is in a different row. In such cases, the original set of vectors must be linearly dependent. This motivates the following result, which we ask you to formally prove as Exercise 16:

Theorem 4.7 If S is any set in \mathbb{R}^n containing k distinct vectors, where $k > n$, then S is linearly dependent.

The Independence Test Method can be adapted for use on vector spaces other than \mathbb{R}^n , as in the next example. We will prove that the Independence Test Method is actually valid in such cases in Section 5.5.

Example 9

Consider the following subset of \mathcal{M}_{22} :

$$S = \left\{ \begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 6 & -1 \\ 3 & 2 \end{bmatrix}, \begin{bmatrix} -11 & 3 \\ -2 & 2 \end{bmatrix} \right\}.$$

We determine whether S is linearly independent using the Independence Test Method. First, we represent the 2×2 matrices in S as 4-vectors. Placing them in a matrix, using each 4-vector as a column, we get

$$\begin{bmatrix} 2 & -1 & 6 & -11 \\ 3 & 0 & -1 & 3 \\ -1 & 1 & 3 & -2 \\ 4 & 1 & 2 & 2 \end{bmatrix}, \text{ which reduces to } \begin{bmatrix} 1 & 0 & 0 & \frac{1}{2} \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -\frac{3}{2} \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

There is no pivot in column 4. Hence, S is linearly dependent. ■

Alternate Characterizations of Linear Independence

We have already seen that a set of two vectors is linearly dependent if one vector is a linear combination of the other. We now generalize this to larger sets as well. Notice in the last two examples that the final columns of the row reduced matrix indicate how to obtain the original vectors in the nonpivot columns from earlier columns. In Example 8, the third column of the row reduced matrix is $[-55, 38]$. The entries -55 and 38 represent the coefficients for a linear combination of the original first and second columns that produces the original third column; that is, $[4, -9] = -55[2, 5] + 38[3, 7]$. Similarly, the fourth column $[65, -46]$ of the row reduced matrix implies $[-8, 3] = 65[2, 5] - 46[3, 7]$. In Example 9, the entries of the fourth column of the row reduced matrix are $\frac{1}{2}, 3, -\frac{3}{2}, 0$, respectively. The first three of these are the coefficients for a linear combination of the first three matrices in S that produces the fourth matrix; that is,

$$\begin{bmatrix} -11 & 3 \\ -2 & 2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 2 & 3 \\ -1 & 4 \end{bmatrix} + 3 \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} - \frac{3}{2} \begin{bmatrix} 6 & -1 \\ 3 & 2 \end{bmatrix}.$$

We see that when vectors are linearly dependent, the Independence Test Method gives a natural way of expressing certain vectors as linear combinations of the others. More generally, we have

Theorem 4.8 Suppose S is a finite set of vectors having at least two elements. Then S is linearly dependent if and only if some vector in S can be expressed as a linear combination of the other vectors in S .

Proof. We start by assuming that S is linearly dependent. Therefore, we have coefficients a_1, \dots, a_n such that $\mathbf{0} = a_1 \mathbf{v}_1 + \dots + a_n \mathbf{v}_n$, with $a_i \neq 0$ for some i . Then,

$$\mathbf{v}_i = \left(-\frac{a_1}{a_i}\right) \mathbf{v}_1 + \dots + \left(-\frac{a_{i-1}}{a_i}\right) \mathbf{v}_{i-1} + \left(-\frac{a_{i+1}}{a_i}\right) \mathbf{v}_{i+1} + \dots + \left(-\frac{a_n}{a_i}\right) \mathbf{v}_n,$$

which expresses \mathbf{v}_i as a linear combination of the other vectors in S .

For the second half of the proof, we assume that there is a vector \mathbf{v}_i in S that is a linear combination of the other vectors in S . Without loss of generality, assume $\mathbf{v}_i = \mathbf{v}_1$; that is, $i = 1$. Therefore, there are real numbers a_2, \dots, a_n such that

$$\mathbf{v}_1 = a_2 \mathbf{v}_2 + a_3 \mathbf{v}_3 + \dots + a_n \mathbf{v}_n.$$

Letting $a_1 = -1$, we get $\mathbf{0} = a_1 \mathbf{v}_1 + \dots + a_n \mathbf{v}_n$. Since $a_1 \neq 0$, this shows that S is linearly dependent, completing the proof of the theorem. \square

Example 10

The set of vectors $S = \{[1, 2, -1], [0, 1, 2], [2, 7, 4]\}$ in \mathbb{R}^3 is linearly dependent because it is possible to express some vector in the set S as a linear combination of the others. For example, $[2, 7, 4] = 2[1, 2, -1] + 3[0, 1, 2]$. From a geometric point of view, the fact that $[2, 7, 4]$ can be expressed as a linear combination of the vectors $[1, 2, -1]$ and $[0, 1, 2]$ means that $[2, 7, 4]$ lies in the plane spanned by $[1, 2, -1]$ and $[0, 1, 2]$, assuming that all three vectors have their initial points at the origin (see Figure 4.5).




FIGURE 4.5

The vector $[2, 7, 4]$ in the plane spanned by $[1, 2, -1]$ and $[0, 1, 2]$

Example 11

Consider the subset $S = \{[1, 2, -1, 1], [2, 1, 0, 1], [2, -2, 1, 0], [11, 1, 1, 4]\}$ of \mathbb{R}^4 . Using the Independence Test Method, we row reduce

$$\begin{bmatrix} 1 & 2 & 2 & 11 \\ 2 & 1 & -2 & 1 \\ -1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 4 \end{bmatrix} \text{ to obtain } \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Because there is no pivot in column 4, S is linearly dependent. This means that at least one vector in S is a linear combination of the others. In particular, the first three entries of the fourth column of the row reduced matrix represent coefficients that express $[11, 1, 1, 4]$ as a linear combination of the other vectors:

$$[11, 1, 1, 4] = 1 \cdot [1, 2, -1, 1] + 3 \cdot [2, 1, 0, 1] + 2 \cdot [2, -2, 1, 0].$$

The characterization of linear dependence and linear independence in Theorem 4.8 can be expressed in alternate notation using the concept of span.

If \mathbf{v} is a vector in a set S , we use the notation $S - \{\mathbf{v}\}$ to represent the set of all (other) vectors in S except \mathbf{v} . Of course, in the special case where $S = \{\mathbf{v}\}$ itself, the set $S - \{\mathbf{v}\} = \{\}$, the empty set. Theorem 4.8 implies that a subset S of two or more vectors in a vector space \mathcal{V} is linearly independent precisely when no vector \mathbf{v} in S is in the span of the remaining vectors. That is,

A set S in a vector space \mathcal{V} is linearly independent if and only if there is no vector $\mathbf{v} \in S$ such that $\mathbf{v} \in \text{span}(S - \{\mathbf{v}\})$.

This statement holds even in the special cases when $S = \{\mathbf{v}\}$ or $S = \{\}$. You are asked to prove this in Exercise 21.

Equivalently, we have

A set S in a vector space \mathcal{V} is linearly dependent if and only if there is some vector $\mathbf{v} \in S$ such that $\mathbf{v} \in \text{span}(S - \{\mathbf{v}\})$.

Another useful characterization of linear independence is the following:

A nonempty set of vectors $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is linearly independent if and only if

- (1) $\mathbf{v}_1 \neq \mathbf{0}$, and
- (2) for each k , $2 \leq k \leq n$, $\mathbf{v}_k \notin \text{span}(\{\mathbf{v}_1, \dots, \mathbf{v}_{k-1}\})$.

This states that S is linearly independent if each vector in S can not be expressed as a linear combination of those vectors listed before it. You are asked to prove this in Exercise 22.

Uniqueness of Expression of a Vector as a Linear Combination

The next theorem serves as the foundation for the rest of this chapter because it gives an even more powerful connection between the concepts of span and linear independence.

Theorem 4.9 Let S be a nonempty finite subset of a vector space \mathcal{V} . Then S is linearly independent if and only if every vector $\mathbf{v} \in \text{span}(S)$ can be expressed *uniquely* as a linear combination of the elements of S .

Proof. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$.

Suppose first that S is linearly independent. Assume that $\mathbf{v} \in \text{span}(S)$ can be expressed both as $\mathbf{v} = a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n$ and as $\mathbf{v} = b_1\mathbf{v}_1 + \dots + b_n\mathbf{v}_n$. In order to show that the linear combination for \mathbf{v} is unique, we need to prove that $a_i = b_i$ for all i . But $\mathbf{0} = \mathbf{v} - \mathbf{v} = (a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n) - (b_1\mathbf{v}_1 + \dots + b_n\mathbf{v}_n) = (a_1 - b_1)\mathbf{v}_1 + \dots + (a_n - b_n)\mathbf{v}_n$. Since S is a linearly independent set, each $a_i - b_i = 0$, by the definition of linear independence, and thus $a_i = b_i$ for all i .

Conversely, assume every vector in $\text{span}(S)$ can be uniquely expressed as a linear combination of elements of S . Since $\mathbf{0} \in \text{span}(S)$, there is exactly one linear combination $a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n$ of elements of S that equals $\mathbf{0}$. But the fact that $\mathbf{0} = 0\mathbf{v}_1 + \dots + 0\mathbf{v}_n$ together with the uniqueness of expression for $\mathbf{0}$ means a_1, \dots, a_n are all zero. Thus, by the definition of linear independence, S is linearly independent. \square

By Theorem 4.9, S is linearly independent if there is precisely one way of reaching any “destination” in $\text{span}(S)$ using the given “directions” in S !

Example 12

Recall the subset $S = \{[1, -1, 0, 2], [0, -2, 1, 0], [2, 0, -1, 1]\}$ of \mathbb{R}^4 from Example 6. In that example, we proved that S is linearly independent. Now

$$[11, 1, -6, 10] = 3[1, -1, 0, 2] + (-2)[0, -2, 1, 0] + 4[2, 0, -1, 1]$$

so $[11, 1, -6, 10]$ is in $\text{span}(S)$. Then by Theorem 4.9, this is the *only* possible way to express $[11, 1, -6, 10]$ as a linear combination of the elements in S .

Recall the subset $S = \{[2, 5], [3, 7], [4, -9], [-8, 3]\}$ of \mathbb{R}^2 from Example 8. In that example, we proved that S is linearly dependent. Just before Theorem 4.8 we showed that $[4, -9] = -55[2, 5] + 38[3, 7]$. This means that $[4, 9] = -55[2, 5] + 38[3, 7] + 0[4, -9] + 0[-8, 3]$, but we can also express this vector as $[4, 9] = 0[2, 5] + 0[3, 7] + 1[4, -9] + 0[-8, 3]$. Since $[4, 9]$ is obviously in $\text{span}(S)$, we have found a vector in $\text{span}(S)$ for which the linear combination of elements in S is not unique, just as Theorem 4.9 asserts. \blacksquare

Linear Independence of Eigenvectors

We will prove in Section 5.6 that any set of fundamental eigenvectors for an $n \times n$ matrix produced by the Diagonalization Method is always linearly independent (also see Exercise 25). Let us assume this for the moment. Now, if the method produces n eigenvectors, then the matrix \mathbf{P} whose columns are these eigenvectors must row reduce to \mathbf{I}_n , by the Independence Test Method. This will establish the claim in Section 3.4 that \mathbf{P} is nonsingular.

Example 13

Consider the 3×3 matrix

$$\mathbf{A} = \begin{bmatrix} -2 & 12 & -4 \\ -2 & 8 & -2 \\ -3 & 9 & -1 \end{bmatrix}.$$

You are asked to show in Exercise 14 that $[4, 2, 3]$ is a fundamental eigenvector for the eigenvalue $\lambda_1 = 1$, and that $[3, 1, 0]$ and $[-1, 0, 1]$ are fundamental eigenvectors for the eigenvalue $\lambda_2 = 2$. We test their linear independence by row reducing

$$\mathbf{P} = \begin{bmatrix} 4 & 3 & -1 \\ 2 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix} \text{ to obtain } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

thus illustrating that this set of fundamental eigenvectors is indeed linearly independent *and* that \mathbf{P} is nonsingular. ■

Linear Independence of Infinite Sets

Most cases in which we check for linear independence involve a *finite* set S . However, we will occasionally want to discuss linear independence for infinite sets of vectors.

Definition An infinite subset S of a vector space \mathcal{V} is **linearly dependent** if and only if there is some finite subset T of S such that T is linearly dependent. S is **linearly independent** if and only if S is *not* linearly dependent.

Example 14

Consider the subset S of \mathcal{M}_{22} consisting of all nonsingular 2×2 matrices. We will show that S is linearly dependent.

Let $T = \{\mathbf{I}_2, 2\mathbf{I}_2\}$, a subset of S . Clearly, since the second element of T is a scalar multiple of the first element of T , T is a linearly dependent set. Hence, S is linearly dependent, since one of its finite subsets is linearly dependent. ■

We can also express the definition of linear independence using the negation of the definition of linear dependence:

An infinite subset S of a vector space \mathcal{V} is linearly independent if and only if every finite subset T of S is linearly independent.

From this, Theorem 4.8 implies that an infinite subset S of a vector space \mathcal{V} is linearly independent if and only if no vector in S is a finite linear combination of other vectors in S .

These characterizations of linear independence are obviously valid as well when S is a finite set.

Example 15

Let $S = \{1, 1 + x, 1 + x + x^2, 1 + x + x^2 + x^3, \dots\}$, an infinite subset of \mathcal{P} . We will show that S is linearly independent.

Suppose $T = \{\mathbf{p}_1, \dots, \mathbf{p}_n\}$ is a finite subset of S , with the polynomials written in order of increasing degree. Also suppose that

$$a_1 \mathbf{p}_1 + \dots + a_n \mathbf{p}_n = \mathbf{0}.$$

We need to show that $a_1 = a_2 = \dots = a_n = 0$. We prove this by contradiction.

Suppose at least one a_i is nonzero. Let a_k be the last nonzero coefficient in the series. Then,

$$a_1 \mathbf{p}_1 + \dots + a_k \mathbf{p}_k = \mathbf{0}, \text{ with } a_k \neq 0.$$

Hence,

$$\mathbf{p}_k = -\frac{a_1}{a_k} \mathbf{p}_1 - \frac{a_2}{a_k} \mathbf{p}_2 - \dots - \frac{a_{k-1}}{a_k} \mathbf{p}_{k-1}.$$

Because all the degrees of the polynomials in T are different and they were listed in order of increasing degree, this equation expresses \mathbf{p}_k as a linear combination of polynomials whose degrees are lower than that of \mathbf{p}_k . This can not happen, and so we get our desired contradiction. ■

The next theorem generalizes Theorem 4.9 to include both finite and infinite sets. You are asked to prove this in Exercise 27.

Theorem 4.10 Let S be a nonempty subset of a vector space \mathcal{V} . Then S is linearly independent if and only if every vector $\mathbf{v} \in \text{span}(S)$ can be expressed *uniquely* as a finite linear combination of the elements of S , if terms with zero coefficients are ignored.

Remember: a *finite* linear combination from an infinite set S involves only a finite number of vectors from S . The phrase “if terms with zero coefficients are ignored”

means that two finite linear combinations from a set S are considered the same when all their terms with nonzero coefficients agree. Adding more terms with zero coefficients to a linear combination is not considered to produce a different linear combination.

Example 16

Recall the set S of nonsingular 2×2 matrices discussed in Example 14. Because S is linearly dependent, some vector in $\text{span}(S)$ can be expressed in more than one way as a linear combination of vectors in S . For example,

$$\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix} = 2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + 2 \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = 1 \begin{bmatrix} 3 & 1 \\ 0 & 3 \end{bmatrix} + (-1) \begin{bmatrix} 1 & -1 \\ -2 & 1 \end{bmatrix}.$$

Summary of Results

This section includes several different, but equivalent, descriptions of linearly independent and linearly dependent sets of vectors. Several additional characterizations are described in the exercises. The most important results from both the section and the exercises are summarized in Table 4.1.

New Vocabulary

Independence Test Method	linearly independent (set of vectors)
linearly dependent (set of vectors)	redundant vector

Highlights

- A set of vectors is linearly dependent if there is a nontrivial linear combination of the vectors that equals $\mathbf{0}$.
- A set of vectors is linearly independent if the only linear combination of the vectors that equals $\mathbf{0}$ is the trivial linear combination (i.e., all coefficients = 0).
- A single element set $\{\mathbf{v}\}$ is linearly independent if and only if $\mathbf{v} \neq \mathbf{0}$.
- A two-element set $\{\mathbf{v}_1, \mathbf{v}_2\}$ is linearly independent if and only if neither vector is a scalar multiple of the other.
- The vectors $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ are linearly independent in \mathbb{R}^n , and the vectors $\{1, x, x^2, \dots, x^n\}$ are linearly independent in \mathcal{P}_n .
- Any set containing the zero vector is linearly dependent.
- The Independence Test Method determines whether a finite set is linearly independent by calculating the reduced row echelon form of the matrix whose *columns* are the given vectors.

Table 4.1 Equivalent conditions for a subset S of a vector space to be linearly independent or linearly dependent

Linear Independence of S	Linear Dependence of S	Source
If $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ and $a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n = \mathbf{0}$, then $a_1 = a_2 = \dots = a_n = 0$. (The zero vector requires zero coefficients.)	If $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$, then $a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n = \mathbf{0}$ for some scalars a_1, a_2, \dots, a_n , with some $a_i \neq 0$. (The zero vector does not require all coefficients to be zero.)	Definition
No vector in S is a finite linear combination of other vectors in S .	Some vector in S is a finite linear combination of other vectors in S .	Theorem 4.8 and Remarks after Example 14
For every $\mathbf{v} \in S$, we have $\mathbf{v} \notin \text{span}(S - \{\mathbf{v}\})$.	There is a $\mathbf{v} \in S$ such that $\mathbf{v} \in \text{span}(S - \{\mathbf{v}\})$.	Alternate characterization
For every $\mathbf{v} \in S$, $\text{span}(S - \{\mathbf{v}\})$ does not contain all the vectors of $\text{span}(S)$.	There is some $\mathbf{v} \in S$ such that $\text{span}(S - \{\mathbf{v}\}) = \text{span}(S)$.	Exercise 12
If $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$, then for each k , $\mathbf{v}_k \notin \text{span}(\{\mathbf{v}_1, \dots, \mathbf{v}_{k-1}\})$. (Each \mathbf{v}_k is not a linear combination of the previous vectors in S .)	If $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$, some \mathbf{v}_k can be expressed as $\mathbf{v}_k = a_1\mathbf{v}_1 + \dots + a_{k-1}\mathbf{v}_{k-1}$. (Some \mathbf{v}_k is a linear combination of the previous vectors in S .)	Exercise 22
Every vector in $\text{span}(S)$ can be uniquely expressed as a linear combination of the vectors in S .	Some vector in $\text{span}(S)$ can be expressed in more than one way as a linear combination of the vectors in S .	Theorem 4.9 and Theorem 4.10
Every finite subset of S is linearly independent.	Some finite subset of S is linearly dependent.	Definition when S is infinite

- If a subset of \mathbb{R}^n contains more than n vectors, then the subset is linearly dependent.
- A set of vectors is linearly dependent if some vector can be expressed as a linear combination of the others (i.e., is in the span of the other vectors). (Such a vector is said to be redundant.)
- A set of vectors is linearly independent if no vector can be expressed as a linear combination of the others (i.e., is in the span of the other vectors).
- A set of vectors is linearly independent if no vector can be expressed as a linear combination of those listed before it in the set.
- A set of fundamental eigenvectors produced by the Diagonalization Method is linearly independent (this will be justified in Section 5.6).

- An infinite set of vectors is linearly dependent if some finite subset is linearly dependent.
- An infinite set of vectors is linearly independent if every finite subset is linearly independent.
- A set S of vectors is linearly independent if and only if every vector in $\text{span}(S)$ is produced by a unique linear combination of the vectors in S .

EXERCISES FOR SECTION 4.4

- ★1. In each part, determine by quick inspection whether the given set of vectors is linearly independent. State a reason for your conclusion.
 - (a) $\{[0, 1, 1]\}$
 - (b) $\{[1, 2, -1], [3, 1, -1]\}$
 - (c) $\{[1, 2, -5], [-2, -4, 10]\}$
 - (d) $\{[4, 2, 1], [-1, 3, 7], [0, 0, 0]\}$
 - (e) $\{[2, -5, 1], [1, 1, -1], [0, 2, -3], [2, 2, 6]\}$
2. Use the Independence Test Method to determine which of the following sets of vectors are linearly independent:
 - ★(a) $\{[1, 9, -2], [3, 4, 5], [-2, 5, -7]\}$
 - ★(b) $\{[2, -1, 3], [4, -1, 6], [-2, 0, 2]\}$
 - (c) $\{[-2, 4, 2], [-1, 5, 2], [3, 5, 1]\}$
 - (d) $\{[5, -2, 3], [-4, 1, -7], [7, -4, -5]\}$
 - ★(e) $\{[2, 5, -1, 6], [4, 3, 1, 4], [1, -1, 1, -1]\}$
 - (f) $\{[1, 3, -2, 4], [3, 11, -2, -2], [2, 8, 3, -9], [3, 11, -8, 5]\}$
3. Use the Independence Test Method to determine which of the following subsets of \mathcal{P}_2 are linearly independent:
 - ★(a) $\{x^2 + x + 1, x^2 - 1, x^2 + 1\}$
 - (b) $\{x^2 - x + 3, 2x^2 - 3x - 1, 5x^2 - 9x - 7\}$
 - ★(c) $\{2x - 6, 7x + 2, 12x - 7\}$
 - (d) $\{x^2 + ax + b \mid |a| = |b| = 1\}$
4. Determine which of the following subsets of \mathcal{P} are linearly independent:
 - ★(a) $\{x^2 - 1, x^2 + 1, x^2 + x\}$
 - (b) $\{1 + x^2 - x^3, 2x - 1, x + x^3\}$
 - ★(c) $\{4x^2 + 2, x^2 + x - 1, x, x^2 - 5x - 3\}$
 - (d) $\{3x^3 + 2x + 1, x^3 + x, x - 5, x^3 + x - 10\}$

$$\star(\mathbf{e}) \{1, x, x^2, x^3, \dots\}$$

$$(\mathbf{f}) \{1, 1 + 2x, 1 + 2x + 3x^2, 1 + 2x + 3x^2 + 4x^3, \dots\}$$

5. Show that the following is a linearly dependent subset of \mathcal{M}_{22} :

$$\left\{ \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 3 & 2 \\ -6 & 1 \end{bmatrix}, \begin{bmatrix} 4 & -1 \\ -5 & 2 \end{bmatrix}, \begin{bmatrix} 3 & -3 \\ 0 & 0 \end{bmatrix} \right\}.$$

6. Prove that the following is linearly independent in \mathcal{M}_{32} :

$$\left\{ \begin{bmatrix} 1 & 2 \\ -1 & 1 \\ 3 & 0 \end{bmatrix}, \begin{bmatrix} 4 & 2 \\ -6 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & -1 \\ 2 & 2 \end{bmatrix}, \begin{bmatrix} 0 & 7 \\ 5 & 2 \\ -1 & 6 \end{bmatrix} \right\}.$$

7. Let $S = \{[1, 1, 0], [-2, 0, 1]\}$.

(a) Show that S is a linearly independent subset of \mathbb{R}^3 .

$\star(\mathbf{b})$ Find a vector \mathbf{v} in \mathbb{R}^3 such that $S \cup \{\mathbf{v}\}$ is also linearly independent.

$\star(\mathbf{c})$ Is the vector \mathbf{v} from part (b) unique, or could some other choice for \mathbf{v} have been made? Why or why not?

$\star(\mathbf{d})$ Find a nonzero vector \mathbf{u} in \mathbb{R}^3 such that $S \cup \{\mathbf{u}\}$ is linearly dependent.

8. Suppose that S is the subset $\{[2, -1, 0, 5], [1, -1, 2, 0], [-1, 0, 1, 1]\}$ of \mathbb{R}^4 .

(a) Show that S is linearly independent.

(b) Find a linear combination of vectors in S that produces $[-2, 0, 3, -4]$ (an element of $\text{span}(S)$).

(c) Is there a different linear combination of the elements of S that yields $[-2, 0, 3, -4]$? If so, find one. If not, why not?

9. Consider $S = \{2x^3 - x + 3, 3x^3 + 2x - 2, x^3 - 4x + 8, 4x^3 + 5x - 7\} \subseteq \mathcal{P}_3$.

(a) Show that S is linearly dependent.

(b) Show that every three-element subset of S is linearly dependent.

(c) Explain why every subset of S containing exactly two vectors is linearly independent. (Note: There are six possible two-element subsets.)

10. Let $\mathbf{u} = [u_1, u_2, u_3]$, $\mathbf{v} = [v_1, v_2, v_3]$, $\mathbf{w} = [w_1, w_2, w_3]$ be three vectors in \mathbb{R}^3 . Show that $S = \{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$ is linearly independent if and only if

$$\begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix} \neq 0.$$

(Hint: Consider the transpose and use the Independence Test Method.)
(Compare this exercise with Exercise 19 in Section 4.3.)

11. For each of the following vector spaces, find a linearly independent subset S containing exactly four elements:
- | | |
|---|---|
| <p>★(a) \mathbb{R}^4</p> <p>(b) \mathbb{R}^5</p> <p>★(c) \mathcal{P}_3</p> | <p>(d) \mathcal{M}_{23}</p> <p>★(e) \mathcal{V} = set of all symmetric matrices in \mathcal{M}_{33}.</p> |
|---|---|
12. Let S be a (possibly infinite) subset of a vector space \mathcal{V} . Prove that S is linearly dependent if and only if there is a vector $\mathbf{v} \in S$ such that $\text{span}(S - \{\mathbf{v}\}) = \text{span}(S)$. (We say that such a vector \mathbf{v} is **redundant** in S because the same set of linear combinations is obtained after \mathbf{v} is removed from S ; that is, \mathbf{v} is not needed.)
13. Find a redundant vector in each given linearly dependent set, and show that it satisfies the definition of a redundant vector given in Exercise 12.
- (a) $\{[4, -2, 6, 1], [1, 0, -1, 2], [0, 0, 0, 0], [6, -2, 5, 5]\}$
- ★(b) $\{[1, 1, 0, 0], [1, 1, 1, 0], [0, 0, -6, 0]\}$
- (c) $\{[x_1, x_2, x_3, x_4] \in \mathbb{R}^4 \mid x_i = \pm 1, \text{ for each } i\}$
14. Verify that the Diagonalization Method of Section 3.4 produces the fundamental eigenvectors given in the text for the matrix \mathbf{A} of Example 13.
15. Let $S_1 = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a subset of a vector space \mathcal{V} , let c be a nonzero real number, and let $S_2 = \{c\mathbf{v}_1, \dots, c\mathbf{v}_n\}$. Show that S_1 is linearly independent if and only if S_2 is linearly independent.
- 16. Prove Theorem 4.7. (Hint: Use the definition of linear dependence. Construct an appropriate homogeneous system of linear equations, and show that the system has a nontrivial solution.)
17. Let \mathbf{f} be a polynomial with at least two nonzero terms having different degrees. Prove that the set $\{\mathbf{f}(x), x\mathbf{f}'(x)\}$ (where \mathbf{f}' is the derivative of \mathbf{f}) is linearly independent in \mathcal{P} .
18. Let \mathcal{V} be a vector space, \mathcal{W} a subspace of \mathcal{V} , S a linearly independent subset of \mathcal{W} , and $\mathbf{v} \in \mathcal{V} - \mathcal{W}$. Prove that $S \cup \{\mathbf{v}\}$ is linearly independent.
19. Let \mathbf{A} be an $n \times m$ matrix, let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be a finite subset of \mathbb{R}^m , and let $T = \{\mathbf{A}\mathbf{v}_1, \dots, \mathbf{A}\mathbf{v}_k\}$, a subset of \mathbb{R}^n .
- (a) Prove that if T is a linearly independent subset of \mathbb{R}^n containing k distinct vectors, then S is a linearly independent subset of \mathbb{R}^m .
- ★(b) Find a matrix \mathbf{A} for which the converse to part (a) is false.
- (c) Show that the converse to part (a) is true if \mathbf{A} is square and nonsingular.
20. Prove that every subset of a linearly independent set is linearly independent.

21. Let S be a subset of a vector space \mathcal{V} . If $S = \{\mathbf{a}\}$ or $S = \{\}$, prove that S is linearly independent if and only if there is no vector $\mathbf{v} \in S$ such that $\mathbf{v} \in \text{span}(S - \{\mathbf{v}\})$.
22. Suppose $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a finite subset of a vector space \mathcal{V} . Prove that S is linearly independent if and only if $\mathbf{v}_1 \neq \mathbf{0}$ and, for each k with $2 \leq k \leq n$, $\mathbf{v}_k \notin \text{span}(\{\mathbf{v}_1, \dots, \mathbf{v}_{k-1}\})$. (Hint: Half of the proof is done by contrapositive. For this half, assume that S is linearly dependent, and use an argument similar to the first half of the proof of Theorem 4.8 to show some \mathbf{v}_k is in $\text{span}(\{\mathbf{v}_1, \dots, \mathbf{v}_{k-1}\})$. For the other half, assume S is linearly independent and show $\mathbf{v}_1 \neq \mathbf{0}$ and each $\mathbf{v}_k \notin \text{span}(\{\mathbf{v}_1, \dots, \mathbf{v}_{k-1}\})$.)
23. Let \mathbf{f} be an n th-degree polynomial in \mathcal{P} , and let $\mathbf{f}^{(i)}$ be the i th derivative of \mathbf{f} . Show that $\{\mathbf{f}, \mathbf{f}^{(1)}, \mathbf{f}^{(2)}, \dots, \mathbf{f}^{(n)}\}$ is a linearly independent subset of \mathcal{P} . (Hint: Reverse the order of the elements, and use Exercise 22.)
24. Let S be a nonempty (possibly infinite) subset of a vector space \mathcal{V} .
 - (a) Prove that S is linearly independent if and only if *some* vector \mathbf{v} in $\text{span}(S)$ has a unique expression as a linear combination of the vectors in S (ignoring zero coefficients).
 - (b) The contrapositive of both halves of the “if and only if” statement in part (a), when combined, gives a necessary and sufficient condition for S to be linearly dependent. What is this condition?
25. Suppose \mathbf{A} is an $n \times n$ matrix and that λ is an eigenvalue for \mathbf{A} . Let $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be a set of fundamental eigenvectors for \mathbf{A} corresponding to λ . Prove that S is linearly independent. (Hint: Consider that each \mathbf{v}_i has a 1 in a coordinate in which all the other vectors in S have a 0.)
26. Suppose T is a linearly independent subset of a vector space \mathcal{V} and that $\mathbf{v} \in \mathcal{V}$.
 - (a) Prove that if $T \cup \{\mathbf{v}\}$ is linearly dependent, then $\mathbf{v} \in \text{span}(T)$.
 - (b) Prove that if $\mathbf{v} \in \text{span}(T)$, then $T \cup \{\mathbf{v}\}$ is linearly independent. (Compare this to Exercise 18.)
- 27. Prove Theorem 4.10. (Hint: Generalize the proof of Theorem 4.9. In the first half of the proof, suppose that $\mathbf{v} \in \text{span}(S)$ and that \mathbf{v} can be expressed as both $a_1\mathbf{u}_1 + \dots + a_k\mathbf{u}_k$ and $b_1\mathbf{v}_1 + \dots + b_l\mathbf{v}_l$ for distinct $\mathbf{u}_1, \dots, \mathbf{u}_k$ and distinct $\mathbf{v}_1, \dots, \mathbf{v}_l$ in S . Consider the union $W = \{\mathbf{u}_1, \dots, \mathbf{u}_k\} \cup \{\mathbf{v}_1, \dots, \mathbf{v}_l\}$, and label the distinct vectors in the union as $\{\mathbf{w}_1, \dots, \mathbf{w}_m\}$. Then use the given linear combinations to express \mathbf{v} in two ways as a linear combination of the vectors in W . Finally, use the fact that W is a linearly independent set.)
- ★ 28. True or False:
 - (a) The set $\{[2, -3, 1], [-8, 12, -4]\}$ is a linearly independent subset of \mathbb{R}^3 .
 - (b) A set $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ in a vector space \mathcal{V} is linearly dependent if \mathbf{v}_2 is a linear combination of \mathbf{v}_1 and \mathbf{v}_3 .

- (c) A subset $S = \{\mathbf{v}\}$ of a vector space \mathcal{V} is linearly dependent if $\mathbf{v} = \mathbf{0}$.
- (d) A subset S of a vector space \mathcal{V} is linearly independent if there is a vector $\mathbf{v} \in S$ such that $\mathbf{v} \in \text{span}(S - \{\mathbf{v}\})$.
- (e) If $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a linearly independent set of vectors in a vector space \mathcal{V} , and $a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n = \mathbf{0}$, then $a_1 = a_2 = \dots = a_n = 0$.
- (f) If S is a subset of \mathbb{R}^4 containing six vectors, then S is linearly dependent.
- (g) Let S be a finite nonempty set of vectors in \mathbb{R}^n . If the matrix \mathbf{A} whose rows are the vectors in S has n pivots after row reduction, then S is linearly independent.
- (h) If $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is a linearly independent set of a vector space \mathcal{V} , then no vector in $\text{span}(S)$ can be expressed as two different linear combinations of $\mathbf{v}_1, \mathbf{v}_2$, and \mathbf{v}_3 .
- (i) If $S = \{\mathbf{v}_1, \mathbf{v}_2\}$ is a subset of a vector space \mathcal{V} , and $\mathbf{v}_3 = 3\mathbf{v}_1 - 2\mathbf{v}_2$, then $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ is linearly dependent.

4.5 BASIS AND DIMENSION

Suppose that S is a subset of a vector space \mathcal{V} and that \mathbf{v} is some vector in \mathcal{V} . We can ask the following two fundamental questions about S and \mathbf{v} :

Existence: Is there a linear combination of vectors in S equal to \mathbf{v} ?

Uniqueness: If so, is this the only such linear combination?

The interplay between existence and uniqueness questions is a pervasive theme throughout mathematics. Answering the existence question is equivalent to determining whether $\mathbf{v} \in \text{span}(S)$. Answering the uniqueness question is equivalent (by Theorem 4.10) to determining whether S is linearly independent.

We are most interested in cases where both existence and uniqueness occur. In this section, we tie together these concepts by examining those subsets of vector spaces that simultaneously span and are linearly independent. Such a subset is called a **basis**.

Definition of Basis

Definition Let \mathcal{V} be a vector space, and let B be a subset of \mathcal{V} . Then B is a **basis** for \mathcal{V} if and only if both of the following are true:

- (1) B spans \mathcal{V} .
- (2) B is linearly independent.

Example 1

We show that $B = \{[1, 2, 1], [2, 3, 1], [-1, 2, -3]\}$ is a basis for \mathbb{R}^3 by showing that it both spans \mathbb{R}^3 and is linearly independent.

First, we use the Simplified Span Method in Section 4.3 to show that B spans \mathbb{R}^3 . Expressing the vectors in B as rows and row reducing the matrix

$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & 1 \\ -1 & 2 & -3 \end{bmatrix} \text{ yields } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

which proves that $\text{span}(B) = \{a[1, 0, 0] + b[0, 1, 0] + c[0, 0, 1] \mid a, b, c \in \mathbb{R}\} = \mathbb{R}^3$.

Next, we must show that B is linearly independent. Expressing the vectors in B as columns, and using the Independence Test Method in Section 4.4, we row reduce

$$\begin{bmatrix} 1 & 2 & -1 \\ 2 & 3 & 2 \\ 1 & 1 & -3 \end{bmatrix} \text{ to obtain } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Hence, B is also linearly independent.

Since B spans \mathbb{R}^3 and is linearly independent, B is a basis for \mathbb{R}^3 . (B is not the only basis for \mathbb{R}^3 , as we show in the next example.)

Example 2

The vector space \mathbb{R}^n has $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ as a basis. Although \mathbb{R}^n has other bases as well, the basis $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ is the most useful for general applications and is therefore referred to as the **standard basis** for \mathbb{R}^n . Thus, we refer to $\{\mathbf{i}, \mathbf{j}\}$ and $\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$ as the standard bases for \mathbb{R}^2 and \mathbb{R}^3 , respectively.

Each of our fundamental examples of vector spaces also has a “standard basis.”

Example 3

The standard basis in \mathcal{M}_{32} is defined as the set

$$\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

More generally, we define the **standard basis** in \mathcal{M}_{mn} to be the set of $m \cdot n$ distinct matrices

$$\{\Psi_{ij} \mid 1 \leq i \leq m, 1 \leq j \leq n\},$$

where Ψ_{ij} is the $m \times n$ matrix with 1 in the (i, j) position and zeroes elsewhere. You should check that these $m \cdot n$ matrices are linearly independent and span \mathcal{M}_{mn} . In addition to the standard basis, \mathcal{M}_{mn} has many other bases as well. ■

Example 4

We define $\{1, x, x^2, x^3\}$ to be the standard basis for \mathcal{P}_3 . More generally, the **standard basis** for \mathcal{P}_n is defined to be the set $\{1, x, x^2, \dots, x^n\}$, containing $n + 1$ elements. Similarly, we define the infinite set $\{1, x, x^2, \dots\}$ to be the **standard basis** for \mathcal{P} . Again, note that in each case these sets both span and are linearly independent.

Of course, the polynomial spaces have other bases. For example, the following is also a basis for \mathcal{P}_4 :

$$\{x^4, x^4 - x^3, x^4 - x^3 + x^2, x^4 - x^3 + x^2 - x, x^3 - 1\}.$$

In Exercise 3, you are asked to verify that this is a basis. ■

Example 5

The empty set, $\{\}$, is a basis for the trivial vector space, $\{\mathbf{0}\}$. At the end of Section 4.3, we defined the span of the empty set to be the trivial vector space. That is, $\{\}$ spans $\{\mathbf{0}\}$. Similarly, at the beginning of Section 4.4, we defined $\{\}$ to be linearly independent. ■

A Technical Lemma

In Examples 1 through 4 we saw that \mathbb{R}^n , \mathcal{P}_n , and \mathcal{M}_{mn} each have some *finite* set for a basis, while \mathcal{P} has an infinite basis. We will mostly be concerned with those vector spaces that have finite bases. To begin our study of such vector spaces, we need to show that if a vector space has *one* basis that is finite, then *all* of its bases are finite, and all have the same size. Proving this requires some effort. We begin with Lemma 4.11.

In Lemma 4.11, and throughout the remainder of the text, we use the notation $|S|$ to represent the number of elements in a set S . For example, if B is the standard basis for \mathbb{R}^3 , $|B| = 3$.

Lemma 4.11 Let S and T be subsets of a vector space \mathcal{V} such that S spans \mathcal{V} , S is finite, and T is linearly independent. Then T is finite and $|T| \leq |S|$.

Proof. If S is empty, then $\mathcal{V} = \{\mathbf{0}\}$. Since $\{\mathbf{0}\}$ is not linearly independent, T is also empty, and so $|T| = |S|$.

Assume that $|S| = n \geq 1$. We will proceed with a proof by contradiction. Suppose that either T is infinite or $|T| > |S| = n$. Then, since every finite subset of T is also linearly independent (see Table 4.1 in Section 4.4), there is a linearly independent set $Y \subseteq T$ such that $|Y| = n + 1$. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ and let $Y = \{\mathbf{w}_1, \dots, \mathbf{w}_n, \mathbf{w}_{n+1}\}$. We will show that $\mathbf{w}_{n+1} \in \text{span}(\{\mathbf{w}_1, \dots, \mathbf{w}_n\})$, which will contradict the linear independence of Y .

Now since \mathcal{S} spans \mathcal{V} , there are scalars a_1, a_2, \dots, a_n such that

$$\mathbf{w}_{n+1} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \cdots + a_n \mathbf{v}_n.$$

Also, there are scalars c_{ij} , for $1 \leq i \leq n$ and $1 \leq j \leq n$, such that

$$\mathbf{w}_1 = c_{11} \mathbf{v}_1 + c_{12} \mathbf{v}_2 + \cdots + c_{1n} \mathbf{v}_n$$

$$\mathbf{w}_2 = c_{21} \mathbf{v}_1 + c_{22} \mathbf{v}_2 + \cdots + c_{2n} \mathbf{v}_n$$

$$\vdots \qquad \qquad \vdots$$

$$\mathbf{w}_n = c_{n1} \mathbf{v}_1 + c_{n2} \mathbf{v}_2 + \cdots + c_{nn} \mathbf{v}_n.$$

Let \mathbf{C} be the $n \times n$ matrix whose (i, j) entry is c_{ij} . Our first step is to prove that \mathbf{C}^T is nonsingular. To do this, we show that the homogeneous system $\mathbf{C}^T \mathbf{x} = \mathbf{0}$ has only the trivial solution. So, let \mathbf{u} represent a solution to the system $\mathbf{C}^T \mathbf{x} = \mathbf{0}$; that is, suppose $\mathbf{C}^T \mathbf{u} = \mathbf{0}$. Then, with $\mathbf{u} = [u_1, \dots, u_n]$, we have

$$\begin{aligned} u_1 \mathbf{w}_1 + u_2 \mathbf{w}_2 + \cdots + u_n \mathbf{w}_n &= u_1 (c_{11} \mathbf{v}_1 + c_{12} \mathbf{v}_2 + \cdots + c_{1n} \mathbf{v}_n) \\ &\quad + u_2 (c_{21} \mathbf{v}_1 + c_{22} \mathbf{v}_2 + \cdots + c_{2n} \mathbf{v}_n) \\ &\quad \vdots \\ &\quad + u_n (c_{n1} \mathbf{v}_1 + c_{n2} \mathbf{v}_2 + \cdots + c_{nn} \mathbf{v}_n) \\ &= c_{11} u_1 \mathbf{v}_1 + c_{12} u_1 \mathbf{v}_2 + \cdots + c_{1n} u_1 \mathbf{v}_n \\ &\quad + c_{21} u_2 \mathbf{v}_1 + c_{22} u_2 \mathbf{v}_2 + \cdots + c_{2n} u_2 \mathbf{v}_n \\ &\quad \vdots \\ &\quad + c_{n1} u_n \mathbf{v}_1 + c_{n2} u_n \mathbf{v}_2 + \cdots + c_{nn} u_n \mathbf{v}_n \\ &= (c_{11} u_1 + c_{21} u_2 + \cdots + c_{n1} u_n) \mathbf{v}_1 \\ &\quad + (c_{12} u_1 + c_{22} u_2 + \cdots + c_{n2} u_n) \mathbf{v}_2 \\ &\quad \vdots \\ &\quad + (c_{1n} u_1 + c_{2n} u_2 + \cdots + c_{nn} u_n) \mathbf{v}_n. \end{aligned}$$

But the coefficient of each \mathbf{v}_i in the last expression is just the i th entry of $\mathbf{C}^T \mathbf{u}$. Hence, the coefficient of each \mathbf{v}_i equals 0. Therefore,

$$u_1 \mathbf{w}_1 + u_2 \mathbf{w}_2 + \cdots + u_n \mathbf{w}_n = 0\mathbf{v}_1 + 0\mathbf{v}_2 + \cdots + 0\mathbf{v}_n = \mathbf{0}.$$

Now, $\{\mathbf{w}_1, \dots, \mathbf{w}_n\}$, a subset of \mathcal{Y} , is linearly independent. Hence, $u_1 = u_2 = \cdots = u_n = 0$. Thus, $\mathbf{u} = \mathbf{0}$, proving that the system $\mathbf{C}^T \mathbf{x} = \mathbf{0}$ has only the trivial solution. From this we conclude that \mathbf{C}^T is nonsingular.

Let $\mathbf{a} = [a_1, \dots, a_n]$, where a_1, \dots, a_n are as previously defined. Since \mathbf{C}^T is nonsingular, the system $\mathbf{C}^T \mathbf{x} = \mathbf{a}$ has a unique solution \mathbf{b} ; that is, there is a vector $\mathbf{b} = [b_1, \dots, b_n]$ such that $\mathbf{C}^T \mathbf{b} = \mathbf{a}$. Using a computation similar to the above, we get

$$\begin{aligned} b_1 \mathbf{w}_1 + b_2 \mathbf{w}_2 + \dots + b_n \mathbf{w}_n &= b_1 (c_{11} \mathbf{v}_1 + c_{12} \mathbf{v}_2 + \dots + c_{1n} \mathbf{v}_n) \\ &\quad + b_2 (c_{21} \mathbf{v}_1 + c_{22} \mathbf{v}_2 + \dots + c_{2n} \mathbf{v}_n) \\ &\quad \vdots \\ &\quad + b_n (c_{n1} \mathbf{v}_1 + c_{n2} \mathbf{v}_2 + \dots + c_{nn} \mathbf{v}_n) \\ &= (c_{11} b_1 + c_{21} b_2 + \dots + c_{n1} b_n) \mathbf{v}_1 \\ &\quad + (c_{12} b_1 + c_{22} b_2 + \dots + c_{n2} b_n) \mathbf{v}_2 \\ &\quad \vdots \\ &\quad + (c_{1n} b_1 + c_{2n} b_2 + \dots + c_{nn} b_n) \mathbf{v}_n. \end{aligned}$$

Now, the coefficient of each \mathbf{v}_i in the last expression equals the i th coordinate of $\mathbf{C}^T \mathbf{b}$, which equals a_i . Hence,

$$b_1 \mathbf{w}_1 + b_2 \mathbf{w}_2 + \dots + b_n \mathbf{w}_n = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_n \mathbf{v}_n = \mathbf{w}_{n+1}.$$

This proves that $\mathbf{w}_{n+1} \in \text{span}(\{\mathbf{w}_1, \dots, \mathbf{w}_n\})$, the desired contradiction, completing the proof of the lemma. \square

Example 6

Let $T = \{[1, 4, 3], [2, -7, 6], [5, 5, -5], [0, 3, 19]\}$, a subset of \mathbb{R}^3 . We already know from Theorem 4.7 that because $|T| > 3$, T is linearly dependent. However, Lemma 4.11 gives us the same conclusion because $\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$ is a spanning set for \mathbb{R}^3 containing three elements, and so the fact that $|T| > 3$ again shows that T is linearly dependent. \blacksquare

Dimension

We can now prove the main result of this section.

Theorem 4.12 Let \mathcal{V} be a vector space, and let B_1 and B_2 be bases for \mathcal{V} such that B_1 has finitely many elements. Then B_2 also has finitely many elements, and $|B_1| = |B_2|$.

Proof. Because B_1 and B_2 are bases for \mathcal{V} , B_1 spans \mathcal{V} and B_2 is linearly independent. Hence, Lemma 4.11 shows that B_2 has finitely many elements and $|B_2| \leq |B_1|$. Now, since B_2 is finite, we can reverse the roles of B_1 and B_2 in this argument to show that $|B_1| \leq |B_2|$. Therefore, $|B_1| = |B_2|$. \square

It follows from Theorem 4.12 that if a vector space \mathcal{V} has *one* basis containing a finite number of elements, then *every* basis for \mathcal{V} is finite, and all bases for \mathcal{V} have the

same number of elements. This allows us to unambiguously define the **dimension** of such a vector space, as follows:

Definition Let \mathcal{V} be a vector space. If \mathcal{V} has a basis B containing a finite number of elements, then \mathcal{V} is said to be **finite dimensional**. In this case, the **dimension** of \mathcal{V} , $\dim(\mathcal{V})$, is the number of elements in any basis for \mathcal{V} . In particular, $\dim(\mathcal{V}) = |B|$.
If \mathcal{V} has no finite basis, then \mathcal{V} is **infinite dimensional**.

Example 7

Because \mathbb{R}^3 has the (standard) basis $\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$, the dimension of \mathbb{R}^3 is 3. Theorem 4.12 then implies that every other basis for \mathbb{R}^3 also has exactly three elements. More generally, $\dim(\mathbb{R}^n) = n$, since \mathbb{R}^n has the basis $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$.

Example 8

Because the standard basis $\{1, x, x^2, x^3\}$ for \mathcal{P}_3 has four elements, $\dim(\mathcal{P}_3) = 4$. Every other basis for \mathcal{P}_3 , such as $\{x^3 - x, x^2 + x + 1, x^3 + x - 5, 2x^3 + x^2 + x - 3\}$, also has four elements. (Verify that this set is a basis for \mathcal{P}_3 .)

Also, $\dim(\mathcal{P}_n) = n + 1$, since \mathcal{P}_n has the basis $\{1, x, x^2, \dots, x^n\}$, containing $n + 1$ elements. Be careful! Many students *erroneously* believe that the dimension of \mathcal{P}_n is n because of the subscript n .

Example 9

The standard basis for \mathcal{M}_{22} contains four elements. Hence, $\dim(\mathcal{M}_{22}) = 4$. In general, from the size of the standard basis for \mathcal{M}_{mn} , we see that $\dim(\mathcal{M}_{mn}) = m \cdot n$.

Example 10

Let $\mathcal{V} = \{\mathbf{0}\}$ be the trivial vector space. Then $\dim(\mathcal{V}) = 0$ because the empty set, which contains no elements, is a basis for \mathcal{V} .

Example 11

Consider the following subsets of \mathbb{R}^4 :

$$S_1 = \{[1, 3, 1, 2], [3, 11, 5, 10], [-2, 4, 4, 4]\} \text{ and}$$

$$S_2 = \{[1, 5, -2, 3], [-2, -8, 8, 8], [1, 1, -10, -2], [0, 2, 4, -9], [3, 13, -10, -8]\}.$$

Since $\dim(\mathbb{R}^4) = 4$, $|S_1| = 3$, and $|S_2| = 5$, Theorem 4.12 shows us that neither S_1 nor S_2 is a basis for \mathbb{R}^4 . In particular, S_1 cannot span \mathbb{R}^4 because the standard basis for \mathbb{R}^4 would then be a linearly independent set that is larger than S_1 , contradicting Lemma 4.11. Similarly, S_2 cannot

be linearly independent because the standard basis would be a spanning set that is smaller than S_2 , again contradicting Lemma 4.11.

Notice, however, that in this case we can make no conclusions regarding whether S_1 is linearly independent or whether S_2 spans \mathbb{R}^4 based solely on the size of these sets. We must check for these properties separately using the techniques of Sections 4.3 and 4.4. ■

Sizes of Spanning Sets and Linearly Independent Sets

Example 11 illustrates the next result, which summarizes much of what we have learned regarding the sizes of spanning sets and linearly independent sets.

Theorem 4.13 Let \mathcal{V} be a finite dimensional vector space.

- (1) Suppose S is a finite subset of \mathcal{V} that spans \mathcal{V} . Then $\dim(\mathcal{V}) \leq |S|$. Moreover, $|S| = \dim(\mathcal{V})$ if and only if S is a basis for \mathcal{V} .
- (2) Suppose T is a linearly independent subset of \mathcal{V} . Then T is finite and $|T| \leq \dim(\mathcal{V})$. Moreover, $|T| = \dim(\mathcal{V})$ if and only if T is a basis for \mathcal{V} .

Proof. Let B be a basis for \mathcal{V} with $|B| = n$. Then $\dim(\mathcal{V}) = |B|$, by definition.

Part (1): Since S is a finite spanning set and B is linearly independent, Lemma 4.11 implies that $|B| \leq |S|$, and so $\dim(\mathcal{V}) \leq |S|$.

If $|S| = \dim(\mathcal{V})$, we prove that S is a basis for \mathcal{V} by contradiction. If S is not a basis, then it is not linearly independent (because it spans). So, by Exercise 12 in Section 4.4 (see Table 4.1), there is a redundant vector in S —that is, a vector \mathbf{v} such that $\text{span}(S - \{\mathbf{v}\}) = \text{span}(S) = \mathcal{V}$. But then $S - \{\mathbf{v}\}$ is a spanning set for \mathcal{V} having fewer than n elements, contradicting the fact that we just observed that the size of a spanning set is never less than the dimension.

Finally, suppose S is a basis for \mathcal{V} . By Theorem 4.12, S is finite, and $|S| = \dim(\mathcal{V})$ by the definition of dimension.

Part (2): Using B as the spanning set S in Lemma 4.11 proves that T is finite and $|T| \leq \dim(\mathcal{V})$.

If $|T| = \dim(\mathcal{V})$, we prove that T is a basis for \mathcal{V} by contradiction. If T is not a basis for \mathcal{V} , then T does not span \mathcal{V} (because it is linearly independent). Therefore, there is a vector $\mathbf{v} \in \mathcal{V}$ such that $\mathbf{v} \notin \text{span}(T)$. Hence, by part (b) of Exercise 26 in Section 4.4, $T \cup \{\mathbf{v}\}$ is also linearly independent. But $T \cup \{\mathbf{v}\}$ has $n + 1$ elements, contradicting the fact we just proved—that a linearly independent subset must have size $\leq \dim(\mathcal{V})$.

Finally, if T is a basis for \mathcal{V} , then $|T| = \dim(\mathcal{V})$, by the definition of dimension. □

Example 12

Recall the subset $B = \{[1, 2, 1], [2, 3, 1], [-1, 2, -3]\}$ of \mathbb{R}^3 from Example 1. In that example, after showing that B spans \mathbb{R}^3 , we could have immediately concluded that B is a basis for \mathbb{R}^3 without having proved linear independence by using part (1) of Theorem 4.13 because B is a spanning set with $\dim(\mathbb{R}^3) = 3$ elements.

Similarly, consider $T = \{3, x + 5, x^2 - 7x + 12, x^3 + 4\}$, a subset of \mathcal{P}_3 . T is linearly independent from Exercise 22 in Section 4.4 (see Table 4.1) because each vector in T is not in the span of those before it. Since $|T| = 4 = \dim(\mathcal{P}_3)$, part (2) of Theorem 4.13 shows that T is a basis for \mathcal{P}_3 . ■

Maximal Linearly Independent Sets and Minimal Spanning Sets

Theorem 4.13 shows that in a finite dimensional vector space, a large enough linearly independent set is a basis, as is a small enough spanning set. The “borderline” size is the dimension of the vector space. No linearly independent sets are larger than this, and no spanning sets are smaller. The next two results illustrate this same principle without explicitly using the dimension. Thus, they are useful in cases in which the dimension is not known or for infinite dimensional vector spaces. Outlines of their proofs are given in Exercises 18 and 19.

Theorem 4.14 Let \mathcal{V} be a vector space with spanning set S (so, $\text{span}(S) = \mathcal{V}$), and let B be a maximal linearly independent subset of S . Then B is a basis for \mathcal{V} .

The phrase “ B is a **maximal linearly independent subset** of S ” means that both of the following are true:

- B is a linearly independent subset of S .
- If $B \subset C \subseteq S$ and $B \neq C$, then C is linearly dependent.

Theorem 4.14 asserts that if there is no way to include another vector from S in B without making B linearly dependent, then B is a basis for $\text{span}(S) = \mathcal{V}$. The converse to Theorem 4.14 is also true (see Exercise 20).

Example 13

Consider the subset $S = \{[1, -2, 1], [3, 1, -2], [5, -3, 0], [5, 4, -5], [0, 0, 0]\}$ of \mathbb{R}^3 and the subset $B = \{[1, -2, 1], [5, -3, 0]\}$ of S . We show that B is a maximal linearly independent subset of S and hence, by Theorem 4.14, it is a basis for $\mathcal{V} = \text{span}(S)$.

Now, B is a linearly independent subset of S . The following equations show that if any of the remaining vectors of S are added to B , the set is no longer linearly independent:

$$[3, 1, -2] = -2[1, -2, 1] + [5, -3, 0]$$

$$[5, 4, -5] = -5[1, -2, 1] + 2[5, -3, 0]$$

$$[0, 0, 0] = 0[1, -2, 1] + 0[5, -3, 0].$$

Thus, B is a maximal linearly independent subset of S and so is a basis for $\text{span}(S)$. ■

Another consequence of Theorem 4.14 is that any vector space \mathcal{V} having a finite spanning set S must be finite dimensional. This is because a maximal linearly independent subset of S , which must also be finite, is a basis for \mathcal{V} (see Exercise 24).

We also have the following result for spanning sets:

Theorem 4.15 Let \mathcal{V} be a vector space, and let B be a minimal spanning set for \mathcal{V} . Then B is a basis for \mathcal{V} .

The phrase “ B is a **minimal spanning set** for \mathcal{V} ” means that both of the following are true:

- B is a subset of \mathcal{V} that spans \mathcal{V} .
- If $C \subset B$ and $C \neq B$, then C does not span \mathcal{V} .

The converse of Theorem 4.15 is true as well (see Exercise 21).

Example 14

Consider the subsets S and B of \mathbb{R}^3 given in Example 13. We can use Theorem 4.15 to give another justification that B is a basis for $\mathcal{V} = \text{span}(S)$. Recall from Example 13 that every vector in S is a linear combination of vectors in B , so $S \subseteq \text{span}(B)$. This fact along with $B \subseteq S$ and Corollary 4.6 shows that $\text{span}(B) = \text{span}(S) = \mathcal{V}$. Also, neither vector in B is a scalar multiple of the other, so that neither vector alone can span \mathcal{V} (why?). Hence, B is a minimal spanning set for \mathcal{V} , and by Theorem 4.15, B is a basis for $\text{span}(S)$. ■

Dimension of a Subspace

We conclude this section with the result that every subspace of a finite dimensional vector space is also finite dimensional. This is important because it tells us that the theorems we have developed about finite dimension apply to all *subspaces* of our basic examples \mathbb{R}^n , \mathcal{M}_{mn} , and \mathcal{P}_n .

Theorem 4.16 Let \mathcal{V} be a finite dimensional vector space, and let \mathcal{W} be a subspace of \mathcal{V} . Then \mathcal{W} is also finite dimensional with $\dim(\mathcal{W}) \leq \dim(\mathcal{V})$. Moreover, $\dim(\mathcal{W}) = \dim(\mathcal{V})$ if and only if $\mathcal{W} = \mathcal{V}$.

The proof of Theorem 4.16 is left for you to do, with hints, in Exercise 22. The only subtle part of this proof involves showing that \mathcal{W} actually has a basis.⁴

⁴ Although it is true that *every* vector space has a basis, we must be careful here, because we have not proven this. In fact, Theorem 4.16 establishes that every subspace of a finite dimensional vector space *does* have a basis and that this basis is finite. Although every finite dimensional vector space has a finite basis by definition, the proof that every infinite dimensional vector space has a basis requires advanced set theory and is beyond the scope of this text.

Example 15

Consider the nested sequence of subspaces of \mathbb{R}^3 given by $\{\mathbf{0}\} \subset \{\text{scalar multiples of } [4, -7, 0]\} \subset xy\text{-plane} \subset \mathbb{R}^3$. Their respective dimensions are $0, 1, 2$, and 3 (why?). Hence, the dimensions of each successive pair of these subspaces satisfy the inequality given in Theorem 4.16. ■

Example 16

It can be shown that $B = \{x^3 + 2x^2 - 4x + 18, 3x^2 + 4x - 4, x^3 + 5x^2 - 3, 3x + 2\}$ is a linearly independent subset of \mathcal{P}_3 . Therefore, by part (2) of Theorem 4.13, B is a basis for \mathcal{P}_3 . However, we can also reach the same conclusion from Theorem 4.16. For, $\mathcal{W} = \text{span}(B)$ has B as a basis (why?), and hence, $\dim(\mathcal{W}) = 4$. But since \mathcal{W} is a subspace of \mathcal{P}_3 and $\dim(\mathcal{P}_3) = 4$, Theorem 4.16 implies that $\mathcal{W} = \mathcal{P}_3$. Hence, B is a basis for \mathcal{P}_3 . ■

New Vocabulary

basis	maximal linearly independent set
dimension	minimal spanning set
finite dimensional (vector space)	standard basis (for $\mathbb{R}^n, \mathcal{M}_{mn}, \mathcal{P}_n$)
infinite dimensional (vector space)	

Highlights

- A basis is a subset of a vector space that both spans and is linearly independent.
- If a finite basis exists for a vector space, the vector space is said to be finite dimensional.
- For a finite dimensional vector space, all bases have the same number of vectors, and this number is known as the dimension of the vector space.
- The standard basis for \mathbb{R}^n is $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$; $\dim(\mathbb{R}^n) = n$.
- The standard basis for \mathcal{P}_n is $\{1, x, x^2, \dots, x^n\}$; $\dim(\mathcal{P}_n) = n + 1$.
- The standard basis for \mathcal{M}_{mn} is $\{\Psi_{ij}\}$, where each Ψ_{ij} has a 1 in the (i, j) entry, and zeroes elsewhere; $\dim(\mathcal{M}_{mn}) = m \cdot n$.
- The basis for the trivial vector space $\{\mathbf{0}\}$ is the empty set $\{\}$; $\dim(\{\mathbf{0}\}) = 0$.
- If no finite basis exists for a vector space, the vector space is said to be infinite dimensional. \mathcal{P} is an infinite dimensional vector space, as is the set of all real-valued functions (under normal operations).
- In a vector space \mathcal{V} with dimension n , the size of a spanning set S is always $\geq n$. If $|S| = n$, then S is a basis for \mathcal{V} .
- In a vector space \mathcal{V} with dimension n , the size of a linearly independent set T is always $\leq n$. If $|T| = n$, then T is a basis for \mathcal{V} .

- A maximal linearly independent set in a vector space is a basis.
- A minimal spanning set in a vector space is a basis.
- In a vector space \mathcal{V} with dimension n , the dimension of a subspace \mathcal{W} is always $\leq n$. If $\dim(\mathcal{W}) = n$, then $\mathcal{W} = \mathcal{V}$.

EXERCISES FOR SECTION 4.5

1. Prove that each of the following subsets of \mathbb{R}^4 is a basis for \mathbb{R}^4 by showing both that it spans \mathbb{R}^4 and that is linearly independent:
 - (a) $\{[2, 1, 0, 0], [0, 1, 1, -1], [0, -1, 2, -2], [3, 1, 0, -2]\}$
 - (b) $\{[6, 1, 1, -1], [1, 0, 0, 9], [-2, 3, 2, 4], [2, 2, 5, -5]\}$
 - (c) $\{[1, 1, 1, 1], [1, 1, 1, -1], [1, 1, -1, -1], [1, -1, -1, -1]\}$
 - (d) $\{[\frac{15}{2}, 5, \frac{12}{5}, 1], [2, \frac{1}{2}, \frac{3}{4}, 1], [-\frac{13}{2}, 1, 0, 4], [\frac{18}{5}, 0, \frac{1}{5}, -\frac{1}{5}]\}$
2. Prove that the following set is a basis for \mathcal{M}_{22} by showing that it spans \mathcal{M}_{22} and is linearly independent:

$$\left\{ \begin{bmatrix} 1 & 4 \\ 2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} -3 & 1 \\ -1 & 0 \end{bmatrix}, \begin{bmatrix} 5 & -2 \\ 0 & -3 \end{bmatrix} \right\}.$$

3. Show that the subset $\{x^4, x^4 - x^3, x^4 - x^3 + x^2, x^4 - x^3 + x^2 - x, x^3 - 1\}$ of \mathcal{P}_4 is a basis for \mathcal{P}_4 .
4. Determine which of the following subsets of \mathbb{R}^4 form a basis for \mathbb{R}^4 :
 - ★(a) $S = \{[7, 1, 2, 0], [8, 0, 1, -1], [1, 0, 0, -2]\}$
 - (b) $S = \{[1, 3, 2, 0], [-2, 0, 6, 7], [0, 6, 10, 7]\}$
 - ★(c) $S = \{[7, 1, 2, 0], [8, 0, 1, -1], [1, 0, 0, -2], [3, 0, 1, -1]\}$
 - (d) $S = \{[1, 3, 2, 0], [-2, 0, 6, 7], [0, 6, 10, 7], [2, 10, -3, 1]\}$
 - ★(e) $S = \{[1, 2, 3, 2], [1, 4, 9, 3], [6, -2, 1, 4], [3, 1, 2, 1], [10, -9, -15, 6]\}$
5. (a) Show that $B = \{[2, 3, 0, -1], [-1, 1, 1, -1]\}$ is a maximal linearly independent subset of $S = \{[1, 4, 1, -2], [-1, 1, 1, -1], [3, 2, -1, 0], [2, 3, 0, -1]\}$.
 - ★(b) Calculate $\dim(\text{span}(S))$.
 - ★(c) Does $\text{span}(S) = \mathbb{R}^4$? Why or why not?
 - (d) Is B a minimal spanning set for $\text{span}(S)$? Why or why not?
6. (a) Show that $B = \{x^3 - x^2 + 2x + 1, 2x^3 + 4x - 7, 3x^3 - x^2 - 6x + 6\}$ is a maximal linearly independent subset of $S = \{x^3 - x^2 + 2x + 1, x - 1, 2x^3 + 4x - 7, x^3 - 3x^2 - 22x + 34, 3x^3 - x^2 - 6x + 6\}$.
 - (b) Calculate $\dim(\text{span}(S))$.

(c) Does $\text{span}(S) = \mathcal{P}_3$? Why or why not?

(d) Is B a minimal spanning set for $\text{span}(S)$? Why or why not?

7. Let \mathcal{W} be the solution set to the matrix equation $\mathbf{A}\mathbf{X} = \mathbf{O}$, where

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & 1 & 0 & -1 \\ 2 & -1 & 0 & 1 & 3 \\ 1 & -3 & -1 & 1 & 4 \\ 2 & 9 & 4 & -1 & -7 \end{bmatrix}.$$

(a) Show that \mathcal{W} is a subspace of \mathbb{R}^5 .

(b) Find a basis for \mathcal{W} .

(c) Show that $\dim(\mathcal{W}) + \text{rank}(\mathbf{A}) = 5$.

8. Prove that every proper nontrivial subspace of \mathbb{R}^3 can be thought of, from a geometric point of view, as either a line through the origin or a plane through the origin.

9. Let \mathbf{f} be a polynomial of degree n . Show that the set $\{\mathbf{f}, \mathbf{f}^{(1)}, \mathbf{f}^{(2)}, \dots, \mathbf{f}^{(n)}\}$ is a basis for \mathcal{P}_n (where $\mathbf{f}^{(i)}$ denotes the i th derivative of \mathbf{f}). (Hint: See Exercise 23 in Section 4.4.)

10. (a) Let \mathbf{A} be a 2×2 matrix. Prove that there are real numbers a_0, a_1, \dots, a_4 , not all zero, such that $a_4\mathbf{A}^4 + a_3\mathbf{A}^3 + a_2\mathbf{A}^2 + a_1\mathbf{A} + a_0\mathbf{I}_2 = \mathbf{O}_2$. (Hint: You can assume that $\mathbf{A}, \mathbf{A}^2, \mathbf{A}^3, \mathbf{A}^4$, and \mathbf{I}_2 are all distinct because if they are not, opposite nonzero coefficients can be chosen for any identical pair to demonstrate that the given statement holds.)

(b) Suppose \mathbf{B} is an $n \times n$ matrix. Show that there must be a nonzero polynomial $\mathbf{p} \in \mathcal{P}_{n^2}$ such that $\mathbf{p}(\mathbf{B}) = \mathbf{O}_n$.

11. (a) Show that $B = \{(x-2), x(x-2), x^2(x-2), x^3(x-2), x^4(x-2)\}$ is a basis for $\mathcal{V} = \{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p}(2) = 0\}$.

★(b) What is $\dim(\mathcal{V})$?

★(c) Find a basis for $\mathcal{W} = \{\mathbf{p} \in \mathcal{P}_5 \mid \mathbf{p}(2) = \mathbf{p}(3) = 0\}$.

★(d) Calculate $\dim(\mathcal{W})$.

★12. Let \mathcal{V} be a finite dimensional vector space.

(a) Let S be a subset of \mathcal{V} with $\dim(\mathcal{V}) \leq |S|$. Find an example to show that S need not span \mathcal{V} .

(b) Let T be a subset of \mathcal{V} with $|T| \leq \dim(\mathcal{V})$. Find an example to show that T need not be linearly independent.

13. Let S be a subset of a finite dimensional vector space \mathcal{V} such that $|S| = \dim(\mathcal{V})$. If S is not a basis for \mathcal{V} , prove that S neither spans \mathcal{V} nor is linearly independent.

14. Let \mathcal{V} be an n -dimensional vector space, and let S be a subset of \mathcal{V} containing exactly n elements. Prove that S spans \mathcal{V} if and only if S is linearly independent.
15. Let \mathbf{A} be a nonsingular $n \times n$ matrix, and let B be a basis for \mathbb{R}^n .
 - (a) Show that $B_1 = \{\mathbf{A}\mathbf{v} | \mathbf{v} \in B\}$ is also a basis for \mathbb{R}^n . (Treat the vectors in B as column vectors.)
 - (b) Show that $B_2 = \{\mathbf{v}\mathbf{A} | \mathbf{v} \in B\}$ is also a basis for \mathbb{R}^n . (Treat the vectors in B as row vectors.)
 - (c) Letting B be the standard basis for \mathbb{R}^n , use the result of part (a) to show that the columns of \mathbf{A} form a basis for \mathbb{R}^n .
 - (d) Prove that the rows of \mathbf{A} form a basis for \mathbb{R}^n .
16. Prove that \mathcal{P} is infinite dimensional by showing that no finite subset S of \mathcal{P} can span \mathcal{P} , as follows:
 - (a) Let S be a finite subset of \mathcal{P} . Show that $S \subseteq \mathcal{P}_n$, for some n .
 - (b) Use part (a) to prove that $\text{span}(S) \subseteq \mathcal{P}_n$.
 - (c) Conclude that S cannot span \mathcal{P} .
17.
 - (a) Prove that if a vector space \mathcal{V} has an infinite linearly independent subset, then \mathcal{V} is not finite dimensional.
 - (b) Use part (a) to prove that any vector space having \mathcal{P} as a subspace is not finite dimensional.
18. The purpose of this exercise is to prove Theorem 4.14. Let \mathcal{V}, S , and B be as given in the statement of the theorem. Suppose $B \neq S$, and $\mathbf{w} \in S$ with $\mathbf{w} \notin B$.
 - (a) Explain why it is sufficient to prove that B spans \mathcal{V} .
 - (b) Prove that if $S \subseteq \text{span}(B)$, then B spans \mathcal{V} .
 - (c) Let $C = B \cup \{\mathbf{w}\}$. Prove that C is linearly dependent.
 - (d) Use part (c) to prove that $\mathbf{w} \in \text{span}(B)$. (Also see part (a) of Exercise 26 in Section 4.4.)
 - (e) Tie together all parts to finish the proof.
19. The purpose of this exercise is to prove Theorem 4.15.
 - (a) Explain why it is sufficient to prove the following statement: Let S be a spanning set for a vector space \mathcal{V} . If S is a minimal spanning set for \mathcal{V} , then S is linearly independent.
 - (b) State the contrapositive of the statement in part (a).
 - (c) Prove the statement from part (b). (Hint: Use Exercise 12 from Section 4.4.)
20. Let B be a basis for a vector space \mathcal{V} . Prove that B is a maximal linearly independent subset of \mathcal{V} . (Note: You may *not* use $\dim(\mathcal{V})$ in your proof, since \mathcal{V} could be infinite dimensional.)

21. Let B be a basis for a vector space \mathcal{V} . Prove that B is a minimal spanning set for \mathcal{V} . (Note: You may *not* use $\dim(\mathcal{V})$ in your proof, since \mathcal{V} could be infinite dimensional.)
22. The purpose of this exercise is to prove Theorem 4.16. Let \mathcal{V} and \mathcal{W} be as given in the theorem. Consider the set A of nonnegative integers defined by $A = \{k \mid \text{a set } T \text{ exists with } T \subseteq \mathcal{W}, |T| = k, \text{ and } T \text{ linearly independent}\}$.
- (a) Prove that $0 \in A$. (Hence, A is nonempty.)
 - (b) Prove that $k \in A$ implies $k \leq \dim(\mathcal{V})$. (Hint: Use Theorem 4.13.) (Hence, A is finite.)
 - (c) Let n be the largest element of A . Let T be a linearly independent subset of \mathcal{W} such that $|T| = n$. Prove T is a maximal linearly independent subset of \mathcal{W} .
 - (d) Use part (c) and Theorem 4.14 to prove that T is a basis for \mathcal{W} .
 - (e) Conclude that \mathcal{W} is finite dimensional and use part (b) to show $\dim(\mathcal{W}) \leq \dim(\mathcal{V})$.
 - (f) Prove that if $\dim(\mathcal{W}) = \dim(\mathcal{V})$, then $\mathcal{W} = \mathcal{V}$. (Hint: Let T be a basis for \mathcal{W} and use part (2) of Theorem 4.13 to show that T is also a basis for \mathcal{V} .)
 - (g) Prove the converse of part (f).
23. Let \mathcal{V} be a subspace of \mathbb{R}^n with $\dim(\mathcal{V}) = n - 1$. (Such a subspace is called a **hyperplane** in \mathbb{R}^n .) Prove that there is a nonzero $\mathbf{x} \in \mathbb{R}^n$ such that $\mathcal{V} = \{\mathbf{v} \in \mathbb{R}^n \mid \mathbf{x} \cdot \mathbf{v} = 0\}$. (Hint: Set up a homogeneous system of equations whose coefficient matrix has a basis for \mathcal{V} as its rows. Then notice that this $(n - 1) \times n$ system has at least one nontrivial solution, say \mathbf{x} .)
24. Let \mathcal{V} be a vector space and let S be a finite spanning set for \mathcal{V} . Prove that \mathcal{V} is finite dimensional.
- ★25. True or False:
- (a) A set B of vectors in a vector space \mathcal{V} is a basis for \mathcal{V} if B spans \mathcal{V} and B is linearly independent.
 - (b) All bases for \mathcal{P}_4 have four elements.
 - (c) $\dim(\mathcal{M}_{43}) = 7$.
 - (d) If S is a spanning set for \mathcal{W} and $\dim(\mathcal{W}) = n$, then $|S| \leq n$.
 - (e) If T is a linearly independent set in \mathcal{W} and $\dim(\mathcal{W}) = n$, then $|T| = n$.
 - (f) If T is a linearly independent set in a finite dimensional vector space \mathcal{W} and S is a finite spanning set for \mathcal{W} , then $|T| \leq |S|$.
 - (g) If \mathcal{W} is a subspace of a finite dimensional vector space \mathcal{V} , then $\dim(\mathcal{W}) < \dim(\mathcal{V})$.
 - (h) Every subspace of an infinite dimensional vector space is infinite dimensional.

- (i) If T is a maximal linearly independent set for a vector space \mathcal{V} and S is a minimal spanning set for \mathcal{V} , then $S = T$.
- (j) If \mathbf{A} is a nonsingular 4×4 matrix, then the rows of \mathbf{A} are a basis for \mathbb{R}^4 .

4.6 CONSTRUCTING SPECIAL BASES

In this section, we present additional methods for finding a basis for a given finite dimensional vector space, starting with either a spanning set or a linearly independent subset.

Using Row Reduction to Construct a Basis

Recall the Simplified Span Method from Section 4.3. Using that method, we were able to simplify the form of $\text{span}(S)$ for a subset S of \mathbb{R}^n . This was done by creating a matrix \mathbf{A} whose rows are the vectors in S , and then row reducing \mathbf{A} to obtain a reduced row echelon form matrix \mathbf{C} . We discovered that a simplified form of $\text{span}(S)$ is given by the set of all linear combinations of the nonzero rows of \mathbf{C} . Now, each nonzero row of the matrix \mathbf{C} has a (pivot) 1 in a column in which all other rows have zeroes, so the nonzero rows of \mathbf{C} must be linearly independent. Thus, the nonzero rows of \mathbf{C} not only span S but are linearly independent as well, and so they form a basis for $\text{span}(S)$. Therefore, whenever we use the Simplified Span Method on a subset S of \mathbb{R}^n , we are actually creating a basis for $\text{span}(S)$.

Example 1

Let $S = \{[2, -2, 3, 5, 5], [-1, 1, 4, 14, -8], [4, -4, -2, -14, 18], [3, -3, -1, -9, 13]\}$, a subset of \mathbb{R}^5 . We can use the Simplified Span Method to find a basis B for $\mathcal{V} = \text{span}(S)$. We construct the matrix

$$\mathbf{A} = \begin{bmatrix} 2 & -2 & 3 & 5 & 5 \\ -1 & 1 & 4 & 14 & -8 \\ 4 & -4 & -2 & -14 & 18 \\ 3 & -3 & -1 & -9 & 13 \end{bmatrix},$$

whose rows are the vectors in S . The reduced row echelon form matrix for \mathbf{A} is

$$\mathbf{C} = \begin{bmatrix} 1 & -1 & 0 & -2 & 4 \\ 0 & 0 & 1 & 3 & -1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Therefore, the desired basis for \mathcal{V} is the set $B = \{[1, -1, 0, -2, 4], [0, 0, 1, 3, -1]\}$ of nonzero rows of \mathbf{C} , and $\dim(\mathcal{V}) = 2$.