

Prerequisite for Linear Algebra

S. K. Panda

IIT Kharagpur

August 29, 2020

For this course, we have to have ideas about partial ordered set(poset), maximal and minimal elements of poset, lower bound and upper bound, Zorn's lemma and fields.

I suggest you to please recall the definitions of vector space, subspace, linear combination, linear span, linearly independent set, linearly dependent set, basis and dimensions.

Definition 0.1. [Relation] Let X and Y be two nonempty sets. A relation R from X to Y is a subset of $X \times Y$, i.e., it is a collection of certain ordered pairs. We write xRy to mean $(x, y) \in R \subseteq X \times Y$.

Example 0.1. 1. Let X be any nonempty set and consider the set $\mathcal{P}(X)$. Define a relation R on $\mathcal{P}(X)$ by $R = \{(S, T) \in \mathcal{P}(X) \times \mathcal{P}(X) : S \subseteq T\}$.

2. Let $A = \{a, b, c, d\}$. Some relations R on A are:

- (a) $R = A \times A$.
- (b) $R = \{(a, a), (b, b), (c, c), (d, d), (a, b), (a, c), (b, c)\}$.
- (c) $R = \{(a, a), (b, b), (c, c)\}$.

Definition 0.2. Let A be a nonempty set. Then, a relation R on A is said to be

- 1. **reflexive:** if for each $a \in A$, $(a, a) \in R$.
- 2. **symmetric:** if for each pair of elements $a, b \in A$, $(a, b) \in R$ implies $(b, a) \in R$.
- 3. **transitive:** if for each triple of elements $a, b, c \in A$, $(a, b), (b, c) \in R$ imply $(a, c) \in R$.
- 4. **antisymmetric** if for each pair of elements $a, b \in A$, $(a, b) \in R$ and $(b, a) \in R$ imply $a = b$.

Definition 0.3. Let X be a nonempty set. A relation R on X is called partially order relation if R is reflexive, anti-symmetric and transitive.

Example 0.2. 1. Let $X = \mathbb{R}$ and xRy if and only if x is less than or equal to y . Then R is a partially order relation.

2. Let $X = \mathbb{N}$ and xRy if and only if x divides y . Then R is a partial order relation.

3. Let X be a non-empty set. Let R is a relation on $\mathcal{P}(X)$ defined by ARB if and only if $A \subseteq B$. Then R is a partial order relation on $\mathcal{P}(X)$.

A partial order relation is denoted by ' \leq '.

Let ' \leq ' be a partial order relation on X . Then (X, \leq) is called partially ordered set.

Definition 0.4. Let (X, \leq) be a partially ordered set. Let $x, y \in X$. Then x and y are comparable if $x \leq y$ or $y \leq x$.

Remark: Let (X, \leq) be a partially ordered set. Let $x, y \in X$. If $x \leq y$, then we say x is less than equal to y or y is greater than equal to x .

Definition 0.5. Let (X, \leq) be a partially ordered set and $A \subseteq X$. Then A is called **totally ordered set** if any two elements in X are comparable. A **chain** in X is a totally ordered subset.

Definition 0.6. Let (X, \leq) be a partially ordered set. Let $A \subseteq X$. An element $a \in A$ is called **maximal element** of A if a is not smaller than any other element of A . That is there is no $b \in A - \{a\}$ such that $a \leq b$.

Let (X, \leq) be a partially ordered set. Let $A \subseteq X$. An element $a \in A$ is called **minimal element** of A if a is not smaller than any other element of A . That is there is no $b \in A - \{a\}$ such that $a \leq b$.

Remarks:

1. Maximal (minimal) element of a set if it exists must be an element of that set.
2. Maximal (minimal) element of a set may not be unique.

Example 0.3. 1. Let $X = \{1, 2, 3\}$. $R = \{(1, 1), (2, 2), (3, 3)\}$. You can easily check that R is a partial order relation on X . Let $A = \{1, 2\}$. Then 1, 2 and 3 are the maximal elements of A .

2. Let (\mathbb{R}, \leq) be a poset where the relation \leq is usual less than or equal on \mathbb{R} . Let $A = (0, 1)$. Then A does not have any maximal element.

3. Let $X = \{1, 2, 3\}$. $R = \{(1, 1), (2, 2), (3, 3)\}$. You can easily check that R is a partial order relation on X . Let $A = \{1, 2\}$. Then 1, 2 and 3 are the minimal elements of A .

4. Let (\mathbb{R}, \leq) be a poset where the relation \leq is usual less than or equal on \mathbb{R} . Let $A = (0, 1)$. Then A does not have minimal element.

Definition 0.7. Let (X, \leq) be a partially ordered set. Let $A \subseteq X$ be a totally ordered subset. An element $a \in X$ is called **upper bound** of A if $b \leq a$ for all $b \in A$.

Let (X, \leq) be a partially ordered set. Let $A \subseteq X$ be a totally ordered subset. An element $a \in X$ is called **lower bound** of A if $b \leq a$ for all $b \in A$.

Example 0.4. 1. Let (\mathbb{R}, \leq) be a poset where the relation \leq is usual less than or equal on \mathbb{R} . Let $A = (0, 1)$. Then $[1, \infty)$ is the set of all upper bounds of A .

2. Let (\mathbb{N}, \leq) be a poset where the relation $x \leq y$ iff x divides y . Let $A = \{2, 4, 6, 8, \dots\}$. Then 2 and 1 are the lower bounds of A .

Zorn's Lemma: Let (X, \leq) be a partially ordered set. Let every chain have upper bound. Then X has a maximal element.

Definition 0.8. [Group] A non-empty set G is said to form a **group** with respect to binary operation \circ , if

1. G is closed under \circ ,
2. \circ is associative,
3. there exists an element e in G such that $e \circ a = a \circ e = a$ for all in G ,
4. for each element a in G , there exists an element a' in G such that $a' \circ a = a \circ a' = e$.

The group is denoted by the symbol (G, \circ) .

Definition 0.9. [Field] A nonempty set R is said to be a **field** if in R there are defined two binary operations, denoted by $+$ and \cdot , respectively, such that

1. $(R, +)$ is abelian group, the identity element is denoted by zero.
2. $(R - \{0\}, \cdot)$ is abelian group, the identity element is denoted 1.
3. $a \cdot (b + c) = a \cdot b + a \cdot c$ and $(b + c) \cdot a = b \cdot a + c \cdot a$ for all $a, b, c \in R$