

3) $\dim V = 2$, $|F| = 2$

$\therefore F$ has only two elements, additive and multiplicative identity i.e. 0 and 1.

\therefore There are only 3 possible basis:

i) $\{(0,1), (1,0)\}$

ii) $\{(1,1), (0,1)\}$

iii) $\{(1,1), (1,0)\}$

Ans) 3c) V has exactly 3 bases.

$$4) \quad V = M_{3 \times 2}(\mathbb{R}) \quad , \quad x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$W = \{ A \in V : Ax = 0 \}$$

$$\text{Let } A = \begin{bmatrix} a & b \\ c & d \\ e & f \end{bmatrix} \quad \therefore \begin{bmatrix} a & b \\ c & d \\ e & f \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0$$

$$\Rightarrow \begin{bmatrix} a+b \\ c+d \\ e+f \end{bmatrix} = 0 \Rightarrow \begin{matrix} a = -b \\ c = -d \\ e = -f \end{matrix}$$

\therefore Standard basis of W :

$$\left\{ \begin{bmatrix} 1 & -1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & -1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & -1 \end{bmatrix} \right\}$$

Ans): 4. c) Dimension of W is 3.

5) If v_1, v_2, v_3 are L

$$c_1 v_1 + c_2 v_2 + c_3 v_3 = 0 \quad \text{has only trivial soln. } c_1 = c_2 = c_3 = 0$$

$$\Rightarrow c_1(u_1 + 2u_2 + 3u_3) + c_2(au_2 + 5u_3) + c_3 \cdot 2u_3 = 0$$

$$\Rightarrow (c_1)u_1 + (2c_1 + ac_2)u_2 + (3c_1 + 5c_2 + 2c_3)u_3 = 0$$

u_1, u_2, u_3 are L.

$$\therefore c_1 = 0$$

$$\therefore 2c_1 + ac_2 = 0$$

$$3c_1 + 5c_2 + 2c_3 = 0$$

This system of homogeneous linear eqn's has trivial soln. iff $D \neq 0$
($c_1 = c_2 = c_3 = 0$)

$$\Rightarrow \begin{vmatrix} 1 & 0 & 0 \\ 2 & a & 0 \\ 3 & 5 & 2 \end{vmatrix} \neq 0$$

$$\Rightarrow 1(2a) \neq 0 \Rightarrow a \neq 0$$

Ans) 5b) v_1, v_2 and v_3 are linearly independent iff $a \neq 0$.

6) $V = M_{n \times n}(\mathbb{R})$, $\dim(V) = n^2$ (no. of places to put 1 to form standard basis)

$W = \{A \in V : A \text{ is an upper triangular and } \text{trace}(A) = 0\}$
 $2x + n = n^2 \Rightarrow x = \frac{n^2 - n}{2}$

$\dim W = \frac{n^2 - n}{2} + n - 1 = \frac{n^2}{2} + \frac{n}{2} - 1$
 (no. of places to put 1 above the principal diagonal, all places below diagonal will have 0)
 (no. of places to put 1 in the diagonal, $\because \text{trace} = 0$ we can consider a_{11} as $a_{11} = -(a_{22} + \dots + a_{nn})$)

$V = W \oplus W^\perp \Rightarrow \dim V = \dim W + \dim W^\perp$
 $\Rightarrow \dim W^\perp = n^2 - \left(\frac{n^2}{2} + \frac{n}{2} - 1\right)$
 $= \frac{n^2}{2} - \frac{n}{2} + 1 = \frac{n^2 - n + 2}{2}$

6 Ans) c) Dimensions of orthogonal complement of W is $\frac{n^2 - n + 2}{2}$.

7) $\|x\| = \sqrt{\langle x, x \rangle}$

$\|x + y\|^2 = \langle x + y, x + y \rangle$
 $= \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle$
 $= \|x\|^2 + \|y\|^2 + \langle x, y \rangle + \langle y, x \rangle$

Given, $\|x + y\|^2 = \|x\|^2 + \|y\|^2$

$\Rightarrow \langle x, y \rangle = \langle y, x \rangle = 0$

$\Rightarrow \overline{\langle y, x \rangle} = \langle y, x \rangle = 0$ (conjugate symmetry of IP)

$\Rightarrow x$ is orthogonal to y & $\langle x, y \rangle \in \mathbb{R}$

Ans) 7c) $\|x + y\|^2 = \|x\|^2 + \|y\|^2 \Rightarrow x$ is orthogonal to y if $\mathbb{K} = \mathbb{R}$

$$8) \quad W = \{ (x_1, x_2, \dots, x_n) : \sum_{i=1}^n x_i = 0 \} \text{ subspace of } \mathbb{R}^n$$

W is non-trivial

$$\frac{\dim(\mathbb{R}^n)}{2} = \frac{n}{2}$$

$$\dim(W) = n-1 \quad (\text{taking } x_1 = -(x_2 + \dots + x_n))$$

$$\therefore \dim(W) \geq \frac{\dim(\mathbb{R}^n)}{2} \quad \left[\begin{array}{l} n-1 \geq \frac{n}{2} \\ \because n \geq 2 \end{array} \right]$$

\therefore Ans) 8a) W has two virtually disjoint complements.

$$9) \quad \{\phi\}^\perp = \{u \in V \mid \langle u, v \rangle = 0 \forall v \in \{\phi\}\}$$

$\therefore \{\phi\}$ is an empty set.

$$\therefore \forall u \in V, \langle u, v \rangle = 0, v \in \{\phi\}$$

$$\therefore \{\phi\}^\perp = V$$

Ans) 9a) $\{\phi\}^\perp = V, \{\phi\}$ is an empty set.

10) $C[-1, 1]$ is an infinite dimensional inner product space.

$$\therefore C[-1, 1] \neq U \oplus U^\perp \quad \left(\begin{array}{l} \text{equality} \\ \text{holds only} \\ \text{for finite-} \\ \text{dimensional IPS} \end{array} \right)$$

$$U = \{ f \in C[-1, 1] : f(0) = 0 \}$$

$$\bullet \text{ Let } f(x) = x \in U$$

$$\text{Let } g(x) = 1$$

$$\therefore \langle f, g \rangle = \int_{-1}^1 x \cdot 1 dx = \frac{x^2}{2} \Big|_{-1}^1 = 0$$

$$\therefore g(x) \in U^\perp$$

$$\therefore 1 \in U^\perp \Rightarrow U^\perp \neq \{0\}$$

\therefore Ans) 10c) U^\perp is a proper and non-trivial subspace of $C[-1, 1]$.